

# OABSTRACTBAND NON

Science Get Together Paracelsus Science Summer

www.pmu.ac.at/forschung-innovation

#### Vorwort

Mit dem alljährlich stattfindenden Paracelsus Science Get Together #SGT durften wir auch in diesem Jahr wieder über 100 Besucher\*innen, Forscher\*innen und Forschungsinteressierte an der Paracelsus Medizinischen Universität (dieses Mal in Nürnberg) begrüßen und gemeinsam einen Tag der Forschung feiern.

Forschende der PMU Salzburg und Nürnberg, des Uniklinikums Salzburg, des Klinikums Nürnberg, sowie Kooperationspartner der PMU reichten insgesamt mehr als 100 Poster ein. Sie stellten ihre aktuellen Forschungsarbeiten vor – diese sind auch beim Virtual Science Summer online über die PMU-Website <u>https://sgt.pmu.ac.at/</u> verfügbar. Die gezeigten Arbeiten umfassen eine enorme thematische Bandbreite aus den verschiedensten Wissenschaftsbereichen, darunter Medizin, Naturund Sozialwissenschaften, von Grundlagen- bis zu klinischer Forschung.

Die eingereichten Poster wurden durch eine Fachjury bewertet und Preisträger\*innen in den Kategorien "Best Poster", "Best PhD Poster" und "Best Student Poster" ausgewählt.

Die ausgewählten Preisträger\*innen sind:

Kategorie Best Poster: **Bernd Neumann** (Nürnberg): "WGS-based characterization of putative hypervirulent Klebsiella pneumoniae identified in a tertiary care hospital in Germany"

Kategorie Best PhD Poster: **Selina Ismair** (Salzburg): ,,Within and between effects of mothers' and fathers' attachment, mentalizing and caregiving" und **Mahdi Safdarian** (Salzburg): ,,Burden of spinal cord injury, 1990 2019: a systematic analysis for the Global Burden of Disease Study 2019"

Kategorie Best Student Poster: **Melina Wirtz** (Nürnberg): "A new approach for management of aneurysmal subarachnoid hemorrhage based on Manual and Artificial Intelligence blood volume measurement in Non-Contrast Head CT scans"

Herzliche Gratulation an alle Gewinner\*innen der Poster Awards!

Die Preisträger\*innen stellten ihre Arbeiten am 28. Juni im Rahmen der Posterausstellung vor.

Ein herzlicher Dank gilt unseren Organisator\*innen aus dem Forschungsmanagement und Services Nürnberg, sowie den zahlreichen Unterstützer\*innen, die auch in diesem Jahr wieder die Durchführung des SGT mitermöglicht haben. Die fünf internationalen Sponsoren Cochlear, Terumo Aortic, Janssen, Bristol Myers Squibb und insbesondere der Hauptsponsor Der Westerhof / Berghotel Sudelfeld (Inhaber: Dr. Andreas Greither), der zum zweiten Mal die Preise der Research and Innovation Preisträger\*innen gesponsert hat, ermöglichten das umfangreiche standortübergreifende Netzwerken.

Wir freuen uns, Ihnen zum Abschluss des Paracelsus Science Get Together und dem bis 15.10.2024 verfügbaren Virtual Science Summer, diesen Abstractband zum besseren Überblick und als Nachschlagewerk übermitteln zu dürfen - wiederum mit einer ISBN-Nummer versehen und damit zitierfähig.

Der besondere Dank der Universität gilt neben den Mitarbeiter\*innen des Forschungsmanagement und Services Nürnberg und Forschungsmanagement Salzburg, der gesamten IT, die das virtuelle Format bestens betreute. So wurde der Netzwerkgedanke des SGT nicht nur von den Teilnehmenden, sondern auch von der Organisation gelebt – ganz im Sinne des Leitbildes der Universität.

Es ist uns ein großes Anliegen, den Vernetzungsgedanken weiter zu verankern und auch gemeinsame Forschungsprojekte zwischen den beiden Universitätsstandorten Salzburg und Nürnberg im Sinne eines Brückenschlages innerhalb der PMU weiter voranzutreiben und zu fördern und auch die Standorthochschulpartner miteinzubeziehen.

Bleiben Sie gesund und neugierig und freuen Sie sich mit uns auf den Science Get Together 2025 in Salzburg, bei dem in einem etwas anderen Format das großartige Portfolio der Forschung an der PMU präsentiert werden soll.

Es grüßen Sie herzlich

Univ.-Prof. Dr. Wolfgang Sperl

Rektor

a.o. Univ.-Prof.<sup>in</sup> Mag.<sup>a</sup> Dr.<sup>in</sup> Barbara Kofler Vizerektorin für Forschungsangelegenheiten

#### Inhaltsverzeichnis

Vo	rwort
----	-------

#### Sponsor\*innen

#### Abstractsammlung

Nr. 1:		A retrospective analysis of an AI-algorithms performance compared to radiologists and the departments acceptance over timeS. 1
Nr. 2:	:	Synergistic effects of Lenvatinib and Dinaciclib combinational treatment in differentiated thyroid cancer cellsS. 2
Nr. 3:	:	Identification of patients with clinically relevant sleep disordered breathing using a contactless device in neurological rehabilitationS. 3
Nr. 4:		Automated diabetic retinopathy screening in out-patient diabetes careS. 4
Nr. 5:	:	Validation of a physiological relevant biomechanical test setup for the investigation of complex pelvic ring fracturesS. 5
Nr. 6:	:	Application of the 5th WHO guidelines for the diagnosis of small lung biopsies in a tertiary care center: is insecurity of pathologists for the right diagnosis justified?S. 6
Nr. 7:	:	What Keeps You Up at Night? Moral Distress in Nurse Leaders in the U.S., Germany, Austria, and   Switzerland.
Nr. 8:		Multimodal evaluation of cryotherapy to reduce incidence of chemotherapy-induced peripheral neurotoxicity in breast cancer patientsS. 8
Nr. 9:	:	Clinical and demographic factors affecting trough levels of isavuconazole in critically ill patients with or without COVID-19S. 9
Nr. 10	0:	Procnostic scoring in Stevens-Johnson-syndrome & toxic epidermal necrolysis : A 17-year analysis at a high-volume centerS. 10
Nr. 1′	1:	Demographic and morphologic factors influencing Transcranial Doppler Sonography (TCD) performanceS. 11
Nr. 12	2:	Influence of Vitamin K2 on the Response of Chondrocytes from Diabetic and Non-diabetic RatsS. 12
Nr. 13	3:	Development of a new protective solution for vein grafts in bypass surgeryS. 13

I

IX

Nr. 14:	Does vitamin K2 (MK7) influence cardiac fibrosis in diabetes mellitus type 2 (T2DM) associated cardiomyopathy progression - insights from a rat diabetes modelS. 14
Nr. 15:	Laboratory and Anamnestic Risk Indicator for Necrotizing FasciitisS. 15
Nr. 16:	Cost-effectiveness of Arthroscopic knee surgery in degenerative meniscal tearsS. 16
Nr. 17:	How do Parkinson's patients from south tyrol sleep?S. 17
Nr. 18:	Generation of a Tendon-specific Knock-Out Mouse Model to Deeply Probe the Role of SPARC in Tendon Health and DiseaseS. 18
Nr. 19:	Structural diversity and biophysical properties of voltage-gated proton channels (Hv)S. 19
Nr. 20:	Sodium Valproate Modulates Cortical Morphology in Juvenile Myoclonic EpilepsyS. 20
Nr. 21:	Small-Molecule Library Screen Identifies Tanespimycin as a YAP-modulator in SH-SY5Y CellsS. 21
Nr. 22:	Effect of Probiotic Supplementation on the Intestinal Microbiome of Preterm InfantsS. 22
Nr. 23:	Sunflower Syndrome – Creating Awareness & International Medication SurveyS. 23
Nr. 24:	Prenatal Psychosocial Risk: Association with Stress, Social Support and ResilienceS. 24
Nr. 25:	From laptop to top-lab: a combined in silico in vitro approach to identify novel microtubule stabilizers as potential drug candidatesS. 25
Nr. 26:	Safe Forests: The psychological wellbeing of harvester and forwarder drivers – a cross-sectional studyS. 26
Nr. 27:	A comparative study of three different bioactive glass scaffolds tailored for cartilage tissue engineeringS. 27
Nr. 28:	Chondrogenesis of hMSCs in 3D printed bioglass/hydrogel scaffolds for cartilage regenerationS. 28
Nr. 29:	Hiatal hernia after transthoracic esophagectomy for cancer: the key role of omentectomyS. 29
Nr. 30:	Impact of Anastomotic Leak vs Pneumonia on failure to rescue after transthoracic Esophagectomy for cancerS. 30
Nr. 31:	Mixed Methods Evaluation of Early Home-Based Interventions for Children at RiskS. 31
Nr. 32:	First insights into SESAM: "SonoElastography: Sarcopenia Associated Muscle Change"S. 32
Nr. 33:	Parents Media Guide: How parental Smartphone use influences family lifeS. 33
Nr. 34:	Role of galanin receptor 2/3 in inflammation associated with peritonitisS. 34
Nr. 35:	Therapeutic Drug Monitoring in critically ill patients with urosepsisS. 35

#### IV

Nr. 36:	First results from a multimodal day-care psychosomatic post-COVID treatment approachS. 36
Nr. 37:	The ubiquitin-proteasome system component Zbtb preferentially interacts with pathogenic variants of SLC26A4 (pendrin)S. 37
Nr. 38:	Comparison of image quality in CT neck imaging between different protocols with linearly reduced radiation doses and contrast agent amountsS. 38
Nr. 39:	Within and between effects of mothers' and fathers' attachment, mentalizing and caregiving 🛎 S. 39
Nr. 40:	Deckung komplexer pararektaler Beckendefekte: Rolle der freien myokutanen Musculus vastus lateralis LappenplastikS. 40
Nr. 41:	Influencing factors on SARS-CoV-2-specific IgG antibody development following vaccinationS. 41
Nr. 42:	Collection efficiency of mononuclear cells in offline extracorporeal photopheresis: can processing time be shortened?S. 42
Nr. 43:	Characterization of patients with classical myeloproliferative neoplasms: A retrospective, single- center analysisS. 43
Nr. 44:	A personalised and customisable serious gaming system for children with movement disordersS. 44
Nr. 45:	Relationship between Education and Steatotic liver diseaseS. 45
Nr. 46:	Caring for a relative with dementia and the lived experiences of boundaries in caregiving – a qualitative study in a rural area of SalzburgS. 46
Nr. 47:	Interatrial block improves prediction of new-onset atrial fibrillation after cardiac surgeryS. 47
Nr. 48:	Novel genetic variants lead to sensorineural hearing loss and Enlarged Vestibular Aqueduct (EVA) S. 48
Nr. 49:	Oxidative stress-related cellular aging causes dysfunction of the Kv3.1/KCNC1 channel reverted by melatoninS. 49
Nr. 50:	Fructose appears to be able to influence the immune systemS. 50
Nr. 51:	Influence of the isolation method on the measurement of the respiratory burstS. 51
Nr. 52:	Telemedicine for the management of behavioral disorders in nursing home residents with dementiaS. 52
Nr. 53:	Impact of sugars on reactive oxygen species production during the respiratory burst of phagocytes S. 53
Nr. 54:	Embroidered silk fibroin scaffolds for ACL tissue engineeringS. 54
Nr. 55:	Transcriptomic analysis unveils the regulatory network of Sturge-Weber syndrome-associated Gαq protein during neural differentiationS. 55

١	/

Nr.	56:	EZH1 - a new therapeutic target for biliary tract cancer?	S. 56
Nr.	57:	Prevalence of metabolic dysfunction-associated fatty liver disease (MAFLD) and effects of glue lowering medications in patients with type 2 diabetes mellitus	cose- S. 57
Nr.	58:	Risk of Interprosthetic Femur Fracture Is Associated with Implant Spacing - A Biomechanical S	Study S. 58
Nr.	59:	Trauma-focused art therapy in patients with miscarriage or stillbirth: Psychometric treatment ef of the randomized controlled MALT! study	fects S. 59
Nr.	60:	The voltage-gated proton channel discovers its family and function	S. 60
Nr.	61:	Pathogenic Pendrin (SLC26a4) variants are functionally rescued by Carfilzomib	S. 61
Nr.	62:	WGS-based characterization of putative hypervirulent Klebsiella pneumoniae identified in a te care hospital in Germany	rtiary S. 62
Nr.	63:	Impact of Mesenchymal Stem Cells Derived Laminin-Binding Extracellular Vesicles on Schu Cells and in Peripheral Nerve Regeneration Processes	wann S. 63
Nr.	64:	Concentrations of antiseizure medications during pregnancy and breastfeeding in women epilepsy: evaluation of acute side effects and systematic follow-up of development of breasted ir A international multicenter study.	with 1fant. S. 64
Nr.	65:	Digital assessment of real-world walking activity in pulmonary hypertension: A prospective bic clinical trial	enter S. 65
Nr.	66:	Patient's Perspective and Expectations in Dupuytren's Disease Treatment	S. 66
Nr.	67:	Barrieren des Zugangs zu Palliative Care im Krankenhaus	S. 67
Nr.	68:	The assessment of biological maturation I academy soccer players: a comparison of two met	hods: S. 68
Nr.	69:	Minimally invasive ascending aortic surgery: operative and mid-term outcomes	S. 69
Nr.	70:	ATR-FTIR as tool for quality control of extracellular vesicles	S. 70
Nr.	71:	Ozanimod – Safety and Efficacy in a real-world-MS-cohort	S. 71
Nr.	72:	The Influence of Chronotype on Interictal Epileptiform Discharges and Sleep Quality in Individual with Epilepsy: A Study of Sleep-EEG Findings	duals S. 72
Nr.	73:	Risk factors for prolonged length of stay after first single-level lumbar microdiscectomy	S. 73
Nr.	74:	Burden of spinal cord injury, 1990 2019: a systematic analysis for the Global Burden of Dis Study 2019	ease S. 74

Nr. 75:	Validation of the arthroplasty register of south Tyrol between 2010-2020 – data analysis of the SABES hospitalsS. 75
Nr. 76:	How primary care physicians deal with the climate crisis: preliminary results of a scoping reviewS. 76
Nr. 77:	Sex differences in pain courses in patients with hip or knee osteoarthritis after multimodal spa therapy in the Austrian valley of GasteinS. 77
Nr. 78:	Comparison of naturally produced EVs and artificial CDVsS. 78
Nr. 79:	Physicians' Perspectives, Well-being, and Fulfilment in TelemedicineS. 79
Nr. 80:	Characterisation of variants in the SLC26A4 gene identified in a Slovak cohort with non-syndromic enlarged vestibular aqueduct and Pendred syndromeS. 80
Nr. 81:	Prognostic and predictive factors (clinical data, histological and molecular data, laboratory values) under palliative first-line therapy for squamous cell carcinoma of the lung with chemotherapy plus immunotherapyS. 81
Nr. 82:	Metabolic Footprint of Ketosis-inducing Diets in MiceS. 82
Nr. 83:	Theranostic EV biomarkers from non-invasive liquid biopsiesS. 83
Nr. 84:	Nanoindentation on Tendon TissueS. 84
Nr. 85:	The clinical relevance of low-grade infected nonunionS. 85
Nr. 86:	Bioanalytic methods in psychosomatic medicineS. 86
Nr. 87:	Detecting high risk neighborhoods and socioeconomic determinants for common oral diseases in Germany with geostatistics and machine learning modelsS. 87
Nr. 88:	The Impact of Educational Status on the Occurrence of Colonic DiverticulaS. 88
Nr. 89:	Optimised intraoperative radiotherapy treatment workflow using machine learning methodsS. 89
Nr. 90:	PKC activators orchestrate neuronal immune modulation: Unveiling microglial dynamics in NF-KB activation and phagocytosisS. 90
Nr. 91:	Preliminary Data of a 3d Single-Limb-Squat Assessment of Laterality in Exercise-proficient PhysiotherapistsS. 91
Nr. 92:	The degree of cyclic stretching influences lapine Achilles tenocytes and hMSCs in 2D culture and under 3D conditions on a Dura mater cell carrierS. 92
Nr. 93:	Lessel-Kubisch syndrome as a model to decipher the Achilles heel of p53S. 93
Nr. 94:	The effect of UVC irradiation on germ density on breast implants and epidermisS. 94
Nr. 95:	DIAFILD – a study aiming to expedite DIAgnosis of Fibrosing Interstitial Lung DiseaseS. 95

r. 96: A new approach for management of aneurysmal subarachnoid hemorrhage based on Manual Artificial Intelligence blood volume measurement in Non-Contrast Head CT scans	and 3. 96
r. 97: Wound healing and self organization of skin cells is guided by protein corona-bearing extracel vesiclesS	llular 3. 97
r. 98: Beyond Stress: The Role of Attachment and Mentalizing for the Well-being among Elemen	ntary
EducatorsS	5. 98
r. 99: Short-term CD8+ T cells ablation reduces microgliosis in the hippocampus of old APP/PS1 anir	mals
S	3. 99
r. 100: 3D Scanning and Printing Technology for Breast Reconstruction: Increased Aesthetic Outcomes	and
Quality of LifeS.	100
r. 101: Impact of 2-hydroxypropyl-β-cyclodextrin inclusion complex formation on dopamine receptor-lig	gand
interaction – a case studyS.	. 101
r. 102: BEATS – MIGRAINE: Biomarker Establishment of Activated Targets in Skin for MigraineS.	. 102

Liste der Einrichtungen (korrespondierende Autor*innen)	X
---	---

XI

#### Verzeichnis aller Autor\*innen

VIII

Der Paracelsus Science Get Together 2024 wurde mit freundlicher Unterstützung folgender Unternehmen ermöglicht:









PHARMACEUTICAL COMPANIES OF Johnson Afohnson

# Bristol Myers Squibb<sup>®</sup>





Jnteres Sudelfeld 4 • 83735 Bayrischzell • Tel. (+49) 8023 81 99-0 info@berghotel-sudelfeld.de • www.berghotel-sudelfeld.de



el. (+49) 08022 - 188 988 • www.derwesterhof.de



# Ihr Weg ist unsere Inspiration

Unser Anliegen ist es, Produkte, Services und Versorgungslösungen zu bieten, die Ihren Patienten das Leben leichter machen.

Wir entwickeln zahlreiche Produkte einschließlich Cochlea- und Knochenleitungsimplantaten für verschiedene Arten von Hörverlust.

Ganz gleich, ob diese Hörlösungen heute oder vor vielen Jahren implantiert wurden – wir möchten unseren Trägern das bestmögliche Hörerlebnis bieten.



Cochlear<sup>™</sup> Nucleus<sup>®</sup> System



Cochlear<sup>™</sup> Baha<sup>®</sup> System

Cochlear<sup>™</sup> Osia<sup>®</sup> System

#### www.cochlear.at 🕜 🖸 🖸

Dieses Material ist für Fachkräfte im Gesundheitswesen vorgesehen. Wenn Sie Kunde bzw. Kundin sind, informieren Sie sich bei Ihrem Arzt über die Möglichkeiten der Behandlung von Hörverlust. Ergebnisse können abweichen; Ihr Arzt berät Sie bezüglich der Faktoren, die Ihr Ergebnis beeinflussen könnten. Lesen Sie stets das Benutzerhandbuch. Nicht alle Produkte sind in allen Ländern erhähltlich. Für Produktinformationen wenden Sie sich bitte an Ihren Vertreter von Cochlear vor Ort. Cochlear, Hear now. And always, Nucleus, Kanso, Baha, Osia, das elliptische Logo und mit dem Symbol © oder ™ versehene Marken sind entweder Marken oder eingetragene Marken von Cochlear Limited oder Cochlear Bone Anchored Solutions AB (sofern nicht anders angegeben).

© Cochlear Limited 2022. D2024642 V1 2022-08 German (Austria) Translation of D1978899 V1 2022-04



# NP WEITER

In der Onkologie haben wir schon viel erreicht. Menschen mit Krebs überleben heute dank moderner Therapien oft länger. Doch Leben ist mehr als Überleben. Leben braucht Perspektiven und Qualität. In jeder Indikation. In jedem Stadium. Dafür arbeiten wir. Jeden Tag. Mit USD 25 Millionen Investitionen pro Tag. Mit Sprunginnovationen, die Grenzen verschieben. Mit beschleunigten Zulassungsverfahren und mit neuen Partnerschaften und Kooperationen.

Unser Anspruch ist Leben.

BIS HIER

Janssen Oncology

©2023 Janssen-Cilag GmbH Bei den abgebildeten Personen handelt es sich um Models, nicht um reale Ärzt:innen.

1\_Bishierundweiter\_Janssen\_Oncology\_Dachmarke\_Anzeigen-HCP\_210x279mm.indd 1

CP-403660

11.11.23 16:21



# Advancing Fenestrated Horizons



To find out more, visit: terumoaortic.com/fenestrated-treo

Visit our website for more information on use, indications, contraindications, warnings/precautions and availability within your market.

Custom made devices are specifically made in accordance with a written prescription of any person authorised by national law by virtue of that person's professional qualifications; which gives (1) specific design characteristics provided under that person's responsibility and (2) is intended for the sole use of a particular patient exclusively to meet their individual conditions and needs. Custom made devices are not available in the US and availability is subject to local regulatory approval.

Manufactured by: Bolton Medical Inc, 799 International Parkway, Sunrise, Florida 33325, United States

OUS\_0247-A Feb23 • PM-06701

Abstractsammlung

## A retrospective analysis of an Al-algorithms performance compared to radiologists and the departments acceptance over time

#### Selim Abed

Radiology, Paracelsus Medical University, Salzburg, Austria; Contact: selim.abed@stud.pmu.ac.at

#### Objective

There is an increasing number of scans being done in hospitals, which is leading to an increase in workload of radiologists. In the emergency setting the detection of intracranialhaemorrhage (ICH) is essential to start treatment as soon as possible. Artificial intelligence(AI) algorithms have shown use in several different medical settings, including the detection of ICHs. In our radiology department, we implemented an FDA-approved and CE-markeddeep learning algorithm to aid in the detection of ICHs.

#### Methods

This study evaluated the performance of the algorithm compared to general radiologists, while using a boardcertified neuro radiologist as "ground truth". We performed two threeweek retrospective analyses six months apart. We also conducted two surveys six monthsapart to gauge the acceptance of the algorithm among radiologists. False negatives and false positives were defined as cases where either the original interpreting radiologist or the AI algorithm misinterpreted the presence of ICH. Discrepancies wereidentified when the AI algorithm flagged cases that were not initially reported as ICH by theradiologist. As stated, before the neuroradiologist served as "ground truth".

#### Results

Our findings demonstrate that the algorithm had a comparable performance to radiologists indetecting ICHs. The sensitivity and specificity that we calculated were in the range reported in previous studies with comparable or the same algorithm. The survey results indicated ahigh level of acceptance of the algorithm among radiologists, which increased slightly overtime. A significant proportion of radiologists found that an additional resident post wouldprovide more value, given they have a similar yearly cost.

#### Conclusions

In conclusion, the implementation of the algorithm to detect ICHs showed promising results. The algorithm' performed comparable to radiologists, and its' integration, especially overtime, led to a high and increasing acceptance among the department's staff. The use of Al inradiology in general has the potential to reprioritise acute finding, improve workflow, reduceerrors in interpretation by serving as second look, and facilitate early treatment for conditionssuch as ICHs, which require immediate attention. We encourage further research into theperformance of Al-algorithm, their potential uses and their acceptance among medicalprofessionals.

- 1. J. C. Hemphill et al., 'Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association', Stroke, vol. 46, no. 7, pp. 2032–2060, Jul. 2015, doi: 10.1161/STR.000000000000069.
- C. J. van Asch, M. J. Luitse, G. J. Rinkel, I. van der Tweel, A. Algra, and C. J. Klijn, 'Incidence, case fatality, and functional outcome of intracerebral haemorrhage over time, according to age, sex, and ethnic origin: a systematic review and meta-analysis', Lancet Neurol., vol. 9, no. 2, pp. 167–176, Feb. 2010, doi: 10.1016/S1474-4422(09)70340-0.
- 3. S. A. Mayer, 'Ultra-early hemostatic therapy for intracerebral hemorrhage', Stroke, vol. 34, no. 1, pp. 224–229, Jan. 2003, doi: 10.1161/01.str.0000046458.67968.e4.

#### Synergistic effects of Lenvatinib and Dinaciclib combinational treatment in differentiated thyroid cancer cells

Celina Ablinger<sup>1</sup>, Petra Huber-Cantonati<sup>1</sup>, Raffael Höllerer<sup>1</sup>, Teresa Kiener<sup>2</sup>, Gundula Rendl<sup>2</sup>, Christian Pirich<sup>2</sup>, Johanna Pachmayr<sup>1</sup>

<sup>1</sup>Institute of Pharmacy, Department of Pharmaceutical Biology and Clinical Pharmacy, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Division of Molecular Imaging and Theranostics, Department of Nuclear Medicine, University Hospital, Paracelsus Medical University, Salzburg, Austria;

Contact: celina.ablinger@pmu.ac.at

#### Objective

Thyroid cancer is the eighth most frequently cancer worldwide, whereas differentiated thyroid carcinomas (DTCs) represent over 90% of thyroid cancers (1). Although thyroidectomy and administration of radioactive iodine results in a good prognosis, some patients develop therapy resistance (RR-DTC) (2). In this case Lenvatinib was approved for the treatment of RR-DTC. Due to severe side effects and/or resistance to therapy, there is an urgent need to develop new treatment strategies (3). Dinaciclib, a Cdk-inhibitor, has shown promising anti-tumor effects in different tumor entities (4). Therefore, the aim of this study is to investigate potential synergistic effects of combinational treatment of Lenvatinib and Dinaciclib in DTC cell lines.

#### **Methods**

To assess the combination effects of Lenvatinib and Dinaciclib, Lenvatinib-resistant K1 and radio-iodine resistant TT2609-C02 cells were treated for 72 h and analyzed via CCK8 kit. The migratory ability of K1 was investigated via a wound healing assay and the Tecan Spark plate reader over 24 h. To elucidate colony formation, DTC cells were pre-treated for 24 h and viable cells were further re-seeded and allowed to form colonies before they were stained with crystal violet. For 3D cell culture, spheroids were treated with Dinaciclib/Lenvatinib for 96 h and spheroid viability was assessed with the CellTiter-Glo® 3D kit.

#### Results

The combinational treatment of Lenvatinib and Dinaciclib significantly reduces the 2D and 3D cell viability in K1 and TT2609-C02 cells either in a synergistic or additive manner. Additionally, for K1 cells, the combination treatment inhibits the migration potential, whereas monotherapies do not lead to a change in the migration behavior. We demonstrated that the simultaneous treatment synergistically decreases the colony formation capacity in both DTC cell lines, while cells receiving monotherapy are still capable of forming colonies.

#### Conclusions

We show that the Cdk-inhibitor Dinaciclib synergistically enhances the efficacy of Lenvatinib treatment. additional experiments will give further insight into the molecular mode of action to improve future therapy options for DTC patients.

#### Acknowledgements

This project is supported by the PMU-FFF research fund (R-19/03/120 ARD).

- 1. Sherman, S. I., (2003) Thyroid Carcinoma. Lancet 361, 501-511
- Busaidy, N.L., and Cabanillas, M. E. (2012) Differentiated thyroid cancer: management of patients with radioiodine 2. nonresponsive disease. J Thyroid Res 2012, 618985
- Schlumberger, M., Tahara, M., Wirth, L. J., Robinson, B., Brose, M. S., Elisei, R., Habra, M. A., Newbold, K. (2015) 3. Lenvatinib versus placebo in raioiodine-refractory thyroid cancer. N Engl J Med 372, 621-630
- 4. Lin SF, Lin JD, Hsueh C, Chou TC, Wong RJ. A cyclin-dependent kinase inhibitor, dinaciclib in preclinical treatment models of thyroid cancer

## Identification of patients with clinically relevant sleep disordered breathing using a contactless device in neurological rehabilitation

Jens Georg Acker<sup>1,2</sup>, Andreas Gantenbein<sup>3,4</sup>, Kneginja Richter<sup>5,6,7</sup>

<sup>1</sup>Clinic of Sleep Medicine, ZURZACH Care, Bad Zurzach, Switzerland; <sup>2</sup>University Clinic for Psychiatry and Psychotherapy, Klinikum Nuernberg, Paracelsus Medical University, Nuremberg, Germany; <sup>3</sup>Pain and Research Department, ZURZACH Care, Bad Zurzach, Switzerland; <sup>4</sup>Department of Neurology, University Hospital Zurich (USZ), Zurich, Switzerland; <sup>5</sup>Faculty for Social Work, Technical University for Applied Sciences, Nuremberg, Germany; <sup>6</sup>Faculty for Medical Sciences, Goce Delcev University, Stip, Macedonia; <sup>7</sup>CuraMed Tagesklinik Nürnberg GmbH, Nuremberg, Germany; Contact: jens.acker@stud.pmu.ac.at

#### Objective

Stroke is one of the five leading causes of death in the Western world. Sleep disturbances are common in ischaemic stroke patients and are associated with poorer stroke outcomes. To improve the quality of medical care for post-stroke patients, we assessed neurological patients in a rehabilitation clinic simultaneously with a non-contact method (Sleepiz One, based on Doppler radar technology and advanced signal processing) and a standard diagnostic polygraphy (PG) measurement.

#### Methods

We studied patients who had suffered a stroke within >5 days but <180 days before inclusion, were older than 18 years, were able to give informed consent, and were able to undergo routine electrophysiological sleep measurement. Of 230 consecutive patients, we recruited 47 participants (the main reasons for exclusion were dementia, aphasia and inability to give written informed consent). We simultaneously performed a standard PG measurement (device set-up by trained carers) and a non-contact Sleepiz-One measurement; 6 measurements were excluded due to insufficient data.

#### Results

Data analysis was performed with 41 complete measurements, simultaneous registration of classic polygraphy and non-contact sleep diagnostics. 58% of the patients suffered from severe sleep apnoea (RDI >15), 33% from severe sleep apnoea (RDI > 25). The comparison of the raw scores showed a correlation of r = 0.79between the conventional (PG) and the non-contact method (Sleepiz One). In 30 cases there was a classification (sleep apnoea (0/1)) with both variables (Respiratory Disturbance Index (per hour) RDI PG >= 15 and RDI Sleepiz One >=15. No significant difference in the prevalences was found (McNemar test, p = 1.0). Both examination methods had a similar error rate.

#### Conclusions

In neurological care, a relevant number of patients with sleep-disordered breathing can be detected with a simple screening examination. In this study, no recognisable difference in the quality of recordings was found between conventional polygraphy (PG) and an innovative contactless device (Sleepiz). Easy-to-use technical procedures such as Sleepiz can reduce the burden on patients and staff.

- 1. Seiler, Andrea; Camilo, Millene; Korostovtseva, Lyudmila; Haynes, Alan G.; Brill, Anne-Kathrin; Horvath, Thomas et al. (2019): Prevalence of sleep-disordered breathing after stroke and TIA: A meta-analysis. In: Neurology 92 (7), e648-e654. DOI: 10.1212/WNL.00000000006904
- Kapur, Vishesh K.; Auckley, Dennis H.; Chowdhuri, Susmita; Kuhlmann, David C.; Mehra, Reena; Ramar, Kannan; Harrod, Christopher G. (2017): Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea. An American Academy of Sleep Medicine Clinical Practice Guideline. In: Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine 13 (3), S. 479–504. DOI: 10.5664/jcsm.6506

#### Automated diabetic retinopathy screening in out-patient diabetes care

Florian Bauer<sup>1</sup>, Annette Sauerbeck<sup>2</sup>, Nick Piravej<sup>1</sup>, Josef Schmidbauer<sup>1</sup>

<sup>1</sup>Department of Opthalmology, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Department of Internal Medicine Endocrinology, Paracelsus Medical University, Nuremberg, Germany; Contact: florian\_maria.bauer@yahoo.com

#### Objective

The aim of this study is to evaluate the performance of an autonomous screening system for diabetic retinopathy (DR) in a real-world setting at the diabetes out-patient center Nuremberg. Digital retinal photographs of the retina were taken by a mobile and a table-top non mydriatic fundus camera. The images were graded for quality and DR by an ophthalmologist and an artificial intelligence (AI) software (1,2).

#### Methods

This is a prospective study. From August 2023 till November 2023 one hundred fifty patients with diabetes type 1 or type 2 were included. Each patient underwent fundus photography in Miosis with a standard 45° camera NW400 (Topcon Medical Systems,Oakland, USA) and then with a hand-held 50° camera Aurora (Optomed Plc, Oulu, Finnland). Two images per eye, macula and optic nerve head centered were taken (figure 1). All images were graded by an ophthalmologist and the artificial intelligence software RetCAD Version 2.2.0 (Thirona Retina, Nijmegen, Netherlands) (3). The DR was graded according to the International Clinical Diabetic Retinopathy Severity Scale (ICDR) (figure 2). More than mild DR (mtmDR) means DR grade 2 or more and was defined as referable DR. The patient was graded taking the worst of two eyes. The images were analyzed by the ophthalmologist first and then by the AI software RetCAD Version 2.2.0 (Thirona Retina, Nijmegen, Netherlands).

#### Results

A total of 150 participants were enrolled, of which 138 completed fundus imaging with both cameras. The average age was 56 years (range, 21-98 years). The mean HbA1c was 7,4 mmol/l (range, 4,8-12,3 mmol/l) (table 1). In 95% of patients DR screening without pupil dilation was possible. The prevalence of mtmDR was 7.3% (10/138). The Al software RetCAD Version 2.2.0 (Thirona Retina, Nijmegen, Netherlands) obtained a sensitivity of 70% and specificity of 98% for recognizing referable DR in 552 fundus images taken with the table top camera NW 400 (Topcon Medical Systems,Oakland, USA). The image quality was rated sufficient in 98%. The Al software RetCAD Version 2.2.0 (Thirona Retina, Nijmegen, Netherlands) reached a sensitivity of 90% and specificity of 100% for correctly identifying referable DR in 552 digital photographs taken with the handheld camera Aurora (Optomed Plc, Oulu,Finnland). The image quality was rated sufficient in 89% (table 2). There was no significant difference in the number of mtm DR patients detected by the ophthalmologist compared to the Al independent of the camera system used.

#### Conclusions

The analysis of retinal images for DR by the AI software RetCAD Version 2.2.0 (Thirona Retina, Nijmegen, Netherlands) showed a high agreement to the evaluation by an ophthalmologist. The handheld fundus camera Aurora (Optomed Plc, Oulu, Finnland) reached similar results compared to the table top camera NW 400 (Topcon Medical Systems,Oakland, USA) despite lower image quality scores (figure 3). The tested automated diabetic retinopathy screening system seems to be suitable for a diabetes out-patient center or primary care setting and offers patients a fast access to routine DR screening.

#### Acknowledgements

Robert Habel (Topcon Company) Hans-Joerg Groeber (Löwe Medizintechnik) Diederik Sakkers (Thirona Retina)

- 1. Abràmoff MD, Lavin PT, Birch M et al. (2018) Pivotal trial of an autonomous Al-based diagnostic system for detection of diabetic retinopathy in primary care offices. NPJ Digit Med 1:39. https://doi.org/10.1038/s41746-018-0040-6
- Joseph S, Selvaraj J, Mani I et al. (2024) Diagnostic Accuracy of Artificial Intelligence-Based Automated Diabetic Retinopathy Screening in Real-World Settings: A Systematic Review and Meta-Analysis. Am J Ophthalmol 263:214– 230. https://doi.org/10.1016/j.ajo.2024.02.012
- Skevas C, Weindler H, Levering M et al. (2022) Simultaneous screening and classification of diabetic retinopathy and age-related macular degeneration based on fundus photos-a prospective analysis of the RetCAD system. Int J Ophthalmol 15:1985–1993. https://doi.org/10.18240/ijo.2022.12.14

## Validation of a physiological relevant biomechanical test setup for the investigation of complex pelvic ring fractures

<u>Dirk Baumeister</u><sup>1,2</sup>, Jonas Mock<sup>1,3</sup>, Nhat Nguyen<sup>1,4</sup>, Bony<sup>1,5</sup>, Markus Greinwald<sup>1</sup>, Martin Winkler<sup>1</sup>, Peter Augat<sup>1,2</sup>

<sup>1</sup>Institute for Biomechanics, BG Unfallklinik Murnau, Germany; <sup>2</sup>Institute for Biomechanics, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Kempten University of Applied Sciences, Germany; <sup>4</sup>Technical University of Munich, Germany; <sup>5</sup>Polytech Engineering School, Marseille, France; Contact: dirk.baumeister@bgu-murnau.de

#### Objective

Fractures of the human pelvis can lead to disruptions of the ring-shaped structure which results in mechanical destabilization of the pelvic ring. A variety of different methods for an osteosynthetic treatment are available, but there is still a lack of knowledge about the mechanical stability of these methods. The aim of this project is to build a test setup able to determine the stability of different fixation methods for complex pelvic ring fractures under a valid loading situation simulating human gait.

#### Methods

In order to be able to evaluate the force distribution on the entire pelvis for the validation of the new test setup, a 6-component load cell was mounted on an embedding pot in addition to the other load cells already installed. With an increasing load protocol, 58 load cycles were applied and the forces normalized to the resulting hip joint force were compared with data from the literature [1,2].

#### Results

In accordance with the literature, the largest force component at the acetabulum acts in the vertical direction at the time point of contralateral heel strike. The forces on the anterior and posterior muscle pull and the force applied to the sacrum correspond to approximately one third of the resulting hip joint force.

#### Conclusions

To date, biomechanical experiments on pelvises have been performed primarily on single-leg stance or doubleleg stance models. Because these do not fully take into account the complex loading situation during gait, the new test setup was developed. Simplifications were made to account for the applied loads, especially the muscle forces. The new test setup can be used to evaluate the fatigue strength of osteosyntheses for complex pelvic ring fractures under dynamic and realistic loading.

- Soliman A, Ricci PL, Kedziora S, Kelm J, Gerich T, Maas S. Developing a Biomechanical Testing Setup of the Pelvis-Part I: Computational Design of Experiments. J Biomech Eng. 2023 Oct 1;145(10):101001. doi: 10.1115/1.4062538. PMID: 37199569.
- Bergmann, G., Bender, A., Dymke, J., Duda, G., & Damm, P. (2016). Standardized loads acting in hip implants. PloS one, 11(5), e0155612.

## Application of the 5th WHO guidelines for the diagnosis of small lung biopsies in a tertiary care center: is insecurity of pathologists for the right diagnosis justified?

#### Manuela Beckert

Institute of Pathology, Paracelsus Medical University, University Hospital Nuremberg, Germany; Contact: manuela.beckert@klinikum-nuernberg.de

#### Objective

The diagnosis of lung cancer based on small lung biopsies is part of the diagnostic routine for many pathologists. According to the recent WHO guidelines, lung cancer in small biopsies should be made by morphology when possible. If this is not possible, a limited panel of additional testing, namely PAS, TTF-1 and p40 should be performed. In daily routine, however, according to our experience numerous ancillary examinations are often conducted to further confirm the morphological diagnosis or to exclude metastases. The aim of this study was to compare the histological diagnosis made by applying the guidelines suggested by the recent WHO classification of lung cancer with the diagnosis made by an extensive use of ancillary techniques.

#### Methods

We have carried out a retrospective analysis of 288 lung biopsies diagnosed in the Department of Pathology, Nuremberg Hospital between September 2020 and May 2021. Two experienced pathologists re-examined all samples according to the guidelines suggested by WHO. We also compared the results of the two immunohistochemical markers used routinely for the diagnosis of squamous cell differentiation, namely p63 and p40.

#### Results

The histological classification was made using a mean number of 1 ancillary technique (SD  $\pm$  1,4) compared with a mean number of 6 techniques (SD $\pm$ 3) in the original diagnostic procedure. By comparing the diagnoses obtained, we were able to show that a comparable diagnostic certainty can be achieved by applying the WHO guidelines in the case of definite primary disease, reaching a concordance rate of 97,3% compared to the extensive use of additional testing. Only in the presence of metastases, the application of limited additional testing could lead to a diagnosis of a primary non-small cell lung carcinoma not otherwise specified (NSCLC NOS) (14/47 metastases). In this setting, clear advantage in diagnostic accuracy using more extensive immunohistochemical examinations is proven. We also found a comparable sensitivity of p40 and p63 for the diagnosis of squamous cell carcinoma, but higher expression of p63 in adenocarcinomas (p<0,001) or NSCLC NOS (p=0,007).

#### Conclusions

The uncertainty of pathologists regarding the diagnosis of lung cancer in small biopsies is justified, in cases where the diagnosis of NSCLC NOS is reached.

#### References

1. Thoracic Tumours WHO Classification of Tumours, 5th Edition, Volume 5. Thoracic Tumours WHO Classification of Tumours, 5th Edition, Volume 5

## What Keeps You Up at Night? Moral Distress in Nurse Leaders in the U.S., Germany, Austria, and Switzerland.

#### Margitta Beil-Hildebrand<sup>1</sup>, Firuzan Sari Kundt<sup>1</sup>, Patrick Kutschar<sup>1</sup>, Lorri Birkholz<sup>2</sup>

<sup>1</sup>Institute of Nursing Science and Practice, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>School of Nursing, Chatham University, Pittsburgh, PA, USA; Contact: m.beil-hildebrand@pmu.ac.at

#### Objective

Nurse leaders are challenged by ethical issues in today's complex health care settings. The purpose of this research was to describe and analyze key elements of moral distress identified by nurse leaders from health care systems in the U.S., Germany, Austria, and Switzerland. The aim was to develop an understanding of distressing ethical issues nurse leaders face in the U.S. and three German-speaking European countries.

#### Methods

This descriptive cross-sectional study surveyed a convenience sample of nurse leaders in the U.S., Germany, Austria, and Switzerland. The voluntary, anonymous survey also included qualitative questions and was distributed using the Qualtrics® platform. A thematic analysis of the qualitative data in each country was carried out and a comparative analysis identified similarities and differences between the groups of nurse leaders comparing the U.S. data to that from three German-speaking European countries.

#### Results

The survey was completed by 316 nurse leaders: Germany, Austria, and Switzerland (n=225) and the U.S. (n=91). Similar themes identified as causing all nurse leaders moral distress included a lack of individual and organizational integrity, hierarchical and interprofessional issues, lack of nursing professionalism, patient care/patient safety concerns, finances negatively impacting care, and issues around social justice. Within these six themes, there were also differences between the U.S. and the three German-speaking European countries.

#### Conclusions

This study demonstrates that it is vital to develop and retain ethically competent nurse leaders in the U.S. and three German-speaking European countries where patient care, professional development, and organizational integrity are paramount. Despite the geo-political differences between these countries, there were reported similarities in the ethical issues that trouble nurse leaders. This study can be a segue into a health care organization's self-assessment to survey its nurse leaders and work collectively to develop supportive leadership development programs, both practical and theoretical, with the goal of reducing moral distress and improving the well-being of these leading employees within complex health care systems.

#### References

 Beil-Hildebrand, M.B., Sari Kundt, F., Kutschar, P. & Birkholz, L. (2024). What Keeps You Up at Night? Moral Distress in Nurse Leaders in the U.S., Germany, Austria, and Switzerland. Leadership in Health Services. 37. DOI: 10.1108/LHS-09-2023-0075

### Multimodal evaluation of cryotherapy to reduce incidence of chemotherapy-induced peripheral neurotoxicity in breast cancer patients

#### Mattia Bergamo<sup>1</sup>, Paola Alberti<sup>2</sup>, Thorsten Fischer<sup>3</sup>, Sonia Prader<sup>1</sup>

<sup>1</sup>Gynecology and Obstetrics department, Südtiroler Sanitätsbetrieb - SABES, Brixen, Italy; <sup>2</sup>School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; <sup>3</sup>Department of Obstetrics and Gynecology, Salzburger Landeskrankenhaus (SALK), Paracelsus Medical University, Salzburg, Austria; Contact: mattia.bergamo@stud.pmu.ac.at

#### Objective

Chemotherapy-induced peripheral neurotoxicity (CIPN) is a common side effect of the drug Paclitaxel, used as adjuvant therapy against breast cancer, which can affect the outcome of cancer therapy, reduce patients' quality of life, and has an important economic and social burden. Although many possible interventions have been proposed in this context, to date there is no standard treatment for the prevention of CIPN (1,4). Cryotherapy, which through cold-induced acromial vasoconstriction reduces the exposure of peripheral nerve fibers to the neurotoxic agent, has been proposed as a promising solution. It can be administered in various ways, including the use of refrigerated gel gloves and stockings or the use of dedicated devices. Many studies support the use of cryotherapy, but to date, there is a lack of consensus on its efficacy and how to perform it (2).

#### Methods

This randomized controlled trial aims to evaluate the efficacy of the two main cryotherapy methods, compared to a control group, using a multimodal assessment approach. For this purpose, 144 breast cancer patients, treated with adjuvant Paclitaxel-weekly in the day-hospitals of Brixen and Meran, will be recruited in the present study and divided into three groups of equal numbers. Group A will receive cryotherapy via a dedicated device (HilothermTM), group B via gloves and refrigerated gel stockings, and group C (control) will not receive cryotherapy. To objectively assess the effectiveness of cryotherapy, robust clinical and neurophysiological outcome measures will be used. Together with a clinical neurological evaluation, following the TNSn score as a guide, and the collection of the FACT/GOG-NTX questionnaire, nerve conduction studies of the dorsal sural nerve, nerve excitability testing of the ulnar nerve, neurological clinical evaluation and measurement of light chain neurofilaments in serum will be performed (3). These evaluation procedures will be collected prospectively during the course of Paclitaxel therapy up to 12-month follow-up.

#### Results

At this time no data are available.

#### Conclusions

Giving the scientific community solid results in favor of the efficacy of cryotherapy in preventing CIPN would encourage its more widespread use. This would ideally reduce the incidence of CIPN, decrease the rate of reduction or discontinuation of chemotherapy, increase the success rate of chemotherapy, and thus increase the life expectancy and quality of life of patients. Furthermore, the development and validation of an effective method to prevent the onset of CIPN, which constitutes a heavy socio-economic burden for patients and public health, would reduce the costs associated with CIPN and thus provide a major contribution to the efforts to build sustainable healthcare for the future.

- 1. Brewer JR, Morrison G, Dolan ME, Fleming GF. Chemotherapy-induced peripheral neuropathy: Current status and progress. Gynecol Oncol. 2016 Jan;140(1):176-83. doi: 10.1016/j.ygyno.2015.11.011. Epub 2015 Nov 7. PMID: 26556766; PMCID: PMC4698212.
- Bailey AG, Brown JN, Hammond JM. Cryotherapy for the prevention of chemotherapy-induced peripheral neuropathy: A systematic review. J Oncol Pharm Pract. 2021 Jan;27(1):156-164. doi: 10.1177/1078155220959431. Epub 2020 Sep 21. PMID: 32955997.
- Alberti P, Bernasconi DP, Cornblath DR, Merkies ISJ, Park SB, Velasco R, Bruna J, Psimaras D, Koeppen S, Pace A, Dorsey SG, Argyriou AA, Kalofonos HP, Briani C, Schenone A, Faber CG, Mazzeo A, Grisold W, Valsecchi M, Cavaletti G; Cl-PeriNomS group. Prospective Evaluation of Health Care Provider and Patient Assessments in Chemotherapy-Induced Peripheral Neurotoxicity. Neurology. 2021 Aug 17;97(7):e660-e672. doi: 10.1212/WNL.000000000012300. Epub 2021 Jun 2. PMID: 34078718; PMCID: PMC10365895.
- Burgess J, Ferdousi M, Gosal D, Boon C, Matsumoto K, Marshall A, Mak T, Marshall A, Frank B, Malik RA, Alam U. Chemotherapy-Induced Peripheral Neuropathy: Epidemiology, Pathomechanisms and Treatment. Oncol Ther. 2021 Dec;9(2):385-450. doi: 10.1007/s40487-021-00168-y. Epub 2021 Oct 16. PMID: 34655433; PMCID: PMC8593126.

#### A new mouse model of DFNB4/Pendred syndrome: CBA/J-SIc26a4p.R409H/p.R409H

Emanuele Bernardinelli<sup>1</sup>, Sebastian Roesch<sup>2</sup>, Martin Jakab<sup>3</sup>, Silvia Dossena<sup>1,4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, Paracelsus Medical University, Salzburg; <sup>2</sup>Department of Otorhinolaryngology, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Center of Physiology, Pathophysiology and Biophysics, Institute of Physiology and Pathophysiology Salzburg, Paracelsus Medical University, Salzburg, Austria; <sup>4</sup>FIZ - Regenerative Medicine & Novel Therapies (RM & NT); Contact: e.bernardinelli@pmu.ac.at

#### Objective

Pendrin (SLC26A4) is an anion exchanger mainly expressed in the inner ear, kidney, and thyroid. Defects in the ion transport function consequent to genetic mutation are associated with syndromic (Pendred syndrome) and non-syndromic forms of deafness (1). The generation of a consistent animal model would help in understanding the pathophysiology of pendrin-related hearing loss and the investigation of potential therapeutic approaches.

#### Methods

A mouse model expressing the pendrin protein variant p.R409H has been produced by CRISPR-Cas with Cyagen-Taconic (Santa Clara, CA, USA). For fast genotyping, a silent mutation resulting in a restriction site for BsrGI has also been introduced. After gDNA extraction and amplification of a 500bp fragment, the presence of the mutation is identified by enzymatic digestion. A homozygous WT produces only a 500bp band on an agarose gel, a heterozygous a 500bp band and a 250bp band, and a homozygous mutant only a 250bp band. To investigate the hearing and vestibular function of the mice, open-field and startle reflex tests were performed. The mice were placed individually in an empty box and their movements were recorded for 15 minutes with a camera and analyzed with Ethovision XT (Noldus, NL).

#### Results

F1 heterozygous mice have been mated in our facility and produced the first F2 litter of 8 pups, 7 males and 1 female. The pups were genotyped with tissue samples from the ear punch. From the restriction analysis, 3 mice were homozygous WT, 3 homozygous mutant, and 2 heterozygous. The vestibular function test was performed at P30 with all 8 mice from the first litter. Several parameters have been evaluated, including the total distance, the velocity, the area occupied, and the rotational movements of the mice. Mouse #7 (MUT) behaved anomalously, remaining the whole time of the experiment in one corner, showing a very agitated state and it was therefore excluded from the analysis.

#### Conclusions

We could successfully produce and mate mice carrying a pathogenic variant in the pendrin gene. From a pilot experiment investigating the vestibular function of the first litter, we could observe that the homozygous mutant mice behaved differently from the WT and heterozygous mice and displayed the typical hallmarks of vestibular dysfunction. Even with the limited number of animals, we could show a significant difference in most of the observed parameters (distance moved, velocity, rotations). Hearing measurements have not been performed yet, but we could observe a lack of startle reflex to a loud clap in all three mutant mice, denoting hearing loss.

#### Acknowledgements

This research was funded by projects 2022-lif-004-DOSSENA, FIZ RM&NT Talent Pool SR 042023, and PMU-FFF R-20/04/136-DOS to Silvia Dossena. The Present address of SR is: Department of Otorhinolaryngology, University Hospital Regensburg, Regensburg, Germany

#### References

1. Smith RJH, et al. Pendred Syndrome / Nonsyndromic Enlarged Vestibular Aqueduct. In: Adam MP, Everman DB, et al., editors. GeneReviews(®). Seattle (WA): University of Washington, Seattle; 1993. PMID: 20301640.

## PROGNOSTIC SCORING IN STEVENS-JOHNSON-SYNDROME & TOXIC EPIDERMAL NECROLYSIS : A 17-YEAR ANALYSIS AT A HIGH-VOLUME CENTER

Moritz Billner<sup>1</sup>, David Breidung<sup>1</sup>, Konrad Karcz<sup>1</sup>, Denis Ehrl<sup>1</sup>

<sup>1</sup>Department of Plastic, Reconstructive and Hand Surgery, Burn Unit, Klinikum Nürnberg, Paracelsus Medical University, Breslauer Str. 201, 90471 Nuremberg, Germany Contact: moritz.billner@klinikum-nuernberg.de

#### Objective

Toxic Epidermal Necrolysis (TEN) and Stevens-Johnson Syndrome (SJS) are rare yet severe dermatologic conditions with potentially life- threatening consequences due to extensive skin and mucous membrane involvement. Despite their low incidence, they pose significant challenges in clinical management and carry high mortality rates. This retrospective study aimed to characterize SJS/TEN cases and evaluate the clinical applicability of existing prognostic scoring systems to predict patient outcomes.

#### Methods

Data from 68 SJS/TEN patients treated over a 17-year period at a specialized burn center were meticulously analyzed. Demographic, clinical, and laboratory information, alongside prognostic scores (SCORTEN, Re-SCORTEN, ABCD-10), and the newly devised Nuremberg Classification Tree for Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis (NCT-SJS/TEN), underwent thorough evaluation regarding their ability to predict mortality.

#### Results

The study identified metamizole as the most frequently associated trigger for SJS/TEN. The overall mortality rate was 51%. Survivors exhibited significantly lower body surface area involvement, platelet counts, and serum blood urea nitrogen levels compared to non-survivors. Among the prognostic scoring systems, only Re-SCORTEN demonstrated significant differentiation between survivors and non- survivors. Notably, the newly developed NCT- SJS/TEN model exhibited a remarkable 89% accuracy in predicting mortality risk.

#### Conclusions

This study underscores the limitations of current prognostic scoring systems in effectively predicting mortality in SJS/TEN cases. The parameters identified of the NCT-SJS/TEN model may offer a potential avenue for enhanced risk stratification and clinical decision-making in managing these complex conditions. However, further validation through larger-scale studies is essential to enhance the reliability and generalizability of predictive models for SJS/TEN. Such advancements hold potential to significantly improve patient outcomes and optimize therapeutic strategies in the clinical management of SJS/TEN.

- 1. Harr T, French LE. Toxic epidermal necrolysis and Stevens-Johnson syndrome. Orphanet J Rare Dis. 2010;5:39. doi:10.1186/1750-1172-5-39
- 2. Borchers AT, Lee JL, Naguwa SM, Cheema GS, Gershwin ME. Stevens-Johnson syndrome and toxic epidermal necrolysis. Autoimmun Rev. 2008;7(8):598-605. doi:10.1016/j.autrev.2008.06.004
- 3. Figure 1.jpg Sehgal VN, Srivastava G. Toxic epidermal necrolysis (TEN) Lyell's syndrome. J Dermatolog Treat. 2005;16(5-6):278-286. doi:10.1080/09546630500375684

## Demographic and morphologic factors influencing Transcranial Doppler Sonography (TCD) performance.

Roberta Bitti<sup>1</sup>, Barbara Schmid<sup>2</sup>, Panagiota Manava<sup>1</sup>, Michael Lell<sup>1</sup>, Josephine Ammon<sup>3</sup>

<sup>1</sup>Department of Radiology, Neuroradiology and Nuclear Medicine, Paracelsus Medical University, Nuremberg, Germany.; <sup>2</sup>Department of Neurology, Paracelsus Medical University, Nuremberg, Germany.; <sup>3</sup>Institute of Medical Physics, Paracelsus Medical University, Nuremberg, Germany; Contact: Roberta.Bitti@klinikum-nuernberg.de

#### Objective

Transcranial Doppler sonography (TCD) is a cost-effective examination that can be performed bedside on the patient to assess the presence of different intracranial pathologies, from brain edema to subdural hemorrhage. TCD can also be used to visualize the middle cerebral artery (MCA) and its pathological changes due mainly to atherosclerosis or vasospasm, the latter being a frequent complication after subarachnoid hemorrhage. The aim of this study was to evaluate the performance of TCD in patients with acute neurological symptoms by testing its efficacy in assessing the MCA as well as three other structures, including the contralateral temporal bone, the mesencephalon, and the ipsilateral sphenoid bone.

#### Methods

Patients who referred to the emergency room of the Klinikum Nürnberg Süd (Nürnberg, Germany) after the onset of acute neurological symptoms between September and November 2021 were eligible for enrollment. Each patient received a computed tomography (CT) scan of the skull and brain to rule out acute intracranial pathologies such as stroke or bleeding. TCD was later performed with a linear array transducer by a trained neurologist as a follow-up examination and to rule out vasospasm. The ultrasound was performed on the temporal bone on both sides of the head. One score was used to evaluate the MCA (0= no visible, 1= poorly visible, 2= visible in its vast majority, and 3= completely visible). Another score, the Transcranial color-coded duplex sonography (Tccs) score was used to assess the visibility of the contralateral temporal bone, the mesencephalon, and the ipsilateral sphenoid bone (0= none of the above-mentioned structures is visible, 1= one structure is visible, 2= two structures are visible, 3= all three of the structures are visible). Furthermore, the area on the temporal bone where the transducer was placed during the TCD was divided into 9 portions and temporal bone thickness (TBT) was measured in millimeters (mm) in 9 different points per side using imaging data from the CT-scans. The two scores were then correlated to age, sex, side and skull thickness and the Pearson coefficient (r) was calculated. Linear regression was used to assess whether TBT significantly predicted the outcome of the MCA and Tccs scores and the R<sup>2</sup> coefficient was calculated.

#### Conclusions

This study shows that the visibility of the intracranial structures at TCD is correlated to age and sex, and particularly that poor diagnostic performance and lower MCA and Tccs scores can be more frequently associated with female sex and older age. Furthermore, a moderate negative correlation between skull thickness and MCA and Tccs scores was found, and regression analyses suggested that with increasing TBT the visibility of intracranial structures decreases substantially. These findings are in agreement with previous literature showing that TCD performance is influenced by sex and age, thus strongly corroborating these notions with data from a vast unselected cohort of patients. Adding to this, we also identified TBT as a further relevant parameter to determine the performance of TCD, suggesting that patient-specific characteristics also influence the visibility of intracranial structures by TCD. Whether this is due to bone thickness alone or even to the mineral density of the bone remains to be determined and has not been assessed during this study. Thus, further validation studies are needed to further define the field of application of TCD.

#### References

1. Blanco P, Abdo-Cuza A. Transcranial Doppler ultrasound in neurocritical care. J Ultrasound. 2018;21(1):1-16. doi:10.1007/s40477-018-0282-9

#### Influence of Vitamin K2 on the Response of Chondrocytes from Diabetic and Non-diabetic Rats

<u>Tobias Braun</u><sup>1</sup>, Vivienne Rüdig<sup>1</sup>, Alexander Reinhardt<sup>1</sup>, Julius Wehrmann<sup>1</sup>, Nils Fleischmann, Maria Kokozidou<sup>1</sup>, Christian Werner<sup>1</sup>, Clemens Gögele<sup>1</sup>, Gundula Schulze-Tanzil<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cellbiology, Paracelsus Medical University, Nuremberg, Germany; Contact: to.braun22@gmail.com

#### Objective

Osteoarthritis (OA) is a chronic inflammatory joint disease and often coexists with type 2 diabetes mellitus (T2DM)[1,2]. The chronic state of hyperglycemia and inflammation in the context of T2DM is believed to play a crucial role in the onset and progression of OA, yet the complex pathogenetic connections are not fully understood [3]. Due to its anti-inflammatory and antioxidative properties, Vitamin K2 (Vit.K2) appears to counteract chronic inflammatory diseases [4], but there is still limited research regarding OA. Therefore, this study aims to investigate the interplay between OA and T2DM, as well as the potentially protective influence of Vit.K2 on articular chondrocytes.

#### Methods

Chondrocytes were isolated from the knee joints of non-diabetic and diabetic rats (Zucker Diabetic Fatty Leptfa/Crl rats), some of which were supplemented with Vit.K2 (Menaquinone-7, 100 mg/kg per day, Kappa Bioscience AS), resulting in four experimental groups: control without (Ctrl w/o Vit.K2) or with Vit.K2 supplementation (Ctrl w/ Vit.K2), and diabetic without (T2DM w/o Vit.K2) or with Vit.K2 supplementation (T2DM w/ Vit.K2). After additional stimulation with tumor necrosis factor  $\alpha$  (TNF $\alpha$ , 10 ng/ml) and anaphylatoxin C5a (25 ng/ml) to induce an inflammatory milieu, as well as culturing in normo- (NG, 1.0 g/l) or hyperglycemic (HG, 4.5 g/l) medium to mimic advanced T2DM conditions, chondrocytes were examined in various assays. In addition to assessing metabolic and proliferative activity, the directed migration/wound healing capacity, as well as undirected motility of chondrocytes, were determined. Furthermore, gene expression analyses were conducted for the cytokine feedback inhibitors 1 and 3 (Socs1 and Socs3) and the stress response gene Heme Oxygenase 1 (Hmox1) using Real-Time detection PCR.

#### Results

In the comparison between chondrocytes from non-diabetic and diabetic rats, no differences were observed in terms of metabolic activity, while higher DNA quantities were detected in chondrocytes from animals with T2DM in terms of cell proliferation. The supplementation of rats with Vit.K2 tended to increase metabolic activity and significantly enhance chondrocyte proliferation. T2DM tended to increase the undirected motility of chondrocytes but significantly decreased directed migration/wound healing capacity. Chondrocytes from rats supplemented with Vit.K2 showed a trend towards higher motility and migration. No differences were found in the gene expression of Socs1, Socs3, and Hmox1 between chondrocytes from non-diabetic and diabetic rats. Supplementation with Vit.K2 tended to decrease the gene expression of Socs1, Socs3, and Hmox1 in chondrocytes.

#### Conclusions

The increase in cell proliferation and undirected motility of chondrocytes induced by T2DM, coupled with a decrease in wound healing capacity, suggests a loss of cell quality that may potentially predispose to OA development. Elevated metabolic and proliferative activity, as well as increased motility and wound healing capacity, indicate a general anabolic effect of Vit.K2 on chondrocytes. The lower expression of Hmox1, Socs1, and Socs3 suggests the anti-inflammatory and antioxidative properties of the vitamin within the joint. The anabolic effects and protective properties support the potential preventive role of Vit.K2 in OA

#### Acknowledgements

We thank Stiftung Endoprothetic for financial support.

- 1. Berenbaum, F., Osteoarthritis as an inflammatory disease (osteoarthritis is not osteoarthrosis!). Osteoarthritis Cartilage, 2013. 21(1): p. 16-21.
- 2. Louati, K., et al., Association between diabetes mellitus and osteoarthritis: systematic literature review and meta-analysis. RMD Open, 2015. 1(1): p. e000077.
- 3. Courties, A. and J. Sellam, Osteoarthritis and type 2 diabetes mellitus: What are the links? Diabetes Res Clin Pract, 2016. 122: p. 198-206.
- 4. Simes, D.C., et al., Vitamin K as a Powerful Micronutrient in Aging and Age-Related Diseases: Pros and Cons from Clinical Studies. Int J Mol Sci, 2019. 20(17).

#### Development of a new protective solution for vein grafts in bypass surgery

<u>Tobias Braun</u><sup>1</sup>, Theodor Fischlein<sup>1</sup>, Gundula Schulze-Tanzil<sup>2</sup>, Clemens Gögele<sup>2</sup>, Maria Kokozidou<sup>2</sup>, Julia Mladek<sup>2</sup>, Rainald Seitelberger<sup>3</sup>

<sup>1</sup>Cardiac Surgery, Klinikum Nürnberg, Germany; <sup>2</sup>Institute for Anatomy and Cellbiology, Paracelsus Medical University, Nuremberg, Germany; <sup>3</sup>Cardiac Surgery, Paracelsus Medical University, Salzburg, Austria; Contact: tobias.braun2@klinikum-nuernberg.de

#### Objective

Bypass operations are often the only treatment option for atherosclerosis [1]. However, saphenous vein grafts (SVGs) frequently fail shortly after surgery. SVGs are stored in preservative solutions, which can lead to functional decline due to the type of preservation solution used [2, 3], surgical trauma, and other factors. Research has shown that saline, the most common solution, is inadequate for storing blood vessels, causing partial detachment of the endothelium [2]. This damage makes the graft more prone to platelet adhesion, macrophage infiltration, plaque development, and intimal hyperplasia, resulting in graft failure [5].

#### Methods

The aim is to develop a preservation solution that protects endothelial cells (ECs) from cellular stress during bypass grafting. In our previous study, we compared several commonly used solutions. Our goal is to further optimize these solutions to protect endothelial integrity better than standard solutions. Remnants of SVGs were used for (immuno-)histological staining to detect endothelial detachment or to isolate ECs for in vitro experiments (e.g., cell viability assays, metabolic activity measurements, migration capability tests, and activity assessments of endothelium-specific genes). Additionally, ex vivo experiments with the vein remnants are planned.

#### Results

Our findings indicate that full electrolyte solutions with an extracellular composition are more suitable for preserving SVGs compared to saline solution (that showed the overall lowest protection) or solutions with a low sodium content. Additionally, metabolizable buffer substances like lactate show potential in maintaining endothelial integrity by serving as an energy source and increasing migration capabilities.

#### Conclusions

The research aims to create a perfusion solution that meets the stringent requirements of cardiac and vascular surgery, improving long-term graft patency and patient outcomes. Such a solution could significantly reduce the failure rate of SVGs and enhance the success of bypass operations.

#### Acknowledgements

This Project is funded by "Brückenschlag Förderung" and in Cooperation with Cardiacsurgery Salzburg. We thank Prof. Dr. Minnich for the oportunity to perform SEM-Images.

- 1. R.N. Singh, J.A. Sosa, G.E. Green, Internal mammary artery versus saphenous vein graft. Comparative performance in patients with combined revascularisation, Br Heart J 50(1) (1983) 48-58.
- D.R. Weiss, G. Juchem, M. Eblenkamp, B.M. Kemkes, B. Gansera, M. Geier, S. Nees, Search for optimized conditions for sealing and storage of bypass vessels: influence of preservation solution and filling pressure on the degree of endothelialization, Int J Clin Exp Med 3(1) (2010) 10-27.
- H.R. Zerkowski, M. Knocks, M.A. Konerding, N. Doetsch, G. Roth, K. Hakim, J.C. Reidemeister, Endothelial damage of the venous graft in CABG. Influence of solutions used for storage and rinsing on endothelial function, Eur J Cardiothorac Surg 7(7) (1993) 376-82.
- 4. Jeffrey V. Manchio, BS, Junyan Gu, MD, Linda Romar, RN, James Brown, MD, James Gammie, MD, Richard N. Pierson III, MD, Bartley Griffith, MD, and Robert S. Poston, MD Disruption of Graft Endothelium Correlates With Early Failure After Off-Pump Coronary Artery Bypass Surgery; Division of Cardiac Surgery, University of Maryland School of Medicine, Baltimore, Maryland
- 5. Christopher D. Owens, Warren J. Gasper, Amreen S. Rahman, Michael S. Conte; Vein graft failure; Journal of Vascular Surgery; Volume 61, Issue 1; 2015; Pages 203-216; ISSN 0741-5214, https://doi.org/10.1016/j.jvs.2013.08.019.

## Does vitamin K2 (MK7) influence cardiac fibrosis in diabetes mellitus type 2 (T2DM) associated cardiomyopathy progression - insights from a rat diabetes model

<u>Tobias Braun</u><sup>1</sup>, Robin Santoro<sup>1</sup>, Maria Kokozidou<sup>1</sup>, Christian Werner<sup>1</sup>, Lukas Motloch<sup>2</sup>, Clemens Gögele<sup>1</sup>, Vera Paar<sup>2</sup>, Gundula Schulze-Tanzil<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cell Biology; <sup>2</sup>Department of Internal Medicine II, Paracelsus Medical University, Salzburg; Contact: Tobias.Braun2@klinikum-nuernberg.de

#### Objective

Diabetic cardiomyopathy is a severe impairment in chronic diabetes mellitus type 2 (T2DM). Vitamin K2 (Vit.K2) is commonly known to provide vasculo-protective effects, but its impact on diabetic cardiomyopathy is unknown. The homozygous Zucker Diabetes Fatty (ZDF fa/fa) rats develop T2DM and cardiomyopathy under diabetogenic high protein diet (Purina 5008), whereas heterozygous rats (ZDF fa/+) can be used under similar conditions as non-diabetic controls. Cardiomyopathy is associated with cardiac fibrosis. Factors contributing to cardiac fibrosis are increased transforming growth factor (TGF) $\beta$ 1 expression, collagen deposition, elevated numbers of myofibroblasts, expressing smooth muscle actin (ACTA2) and upregulated inflammatory mediators (1).This project aims to get a deeper understanding of the pathogenesis of cardiac fibrosis in the T2DM ZDF rat model and to define the influence of Vit.K2 on the heart.

#### Methods

Folowing sacrifice, rat hearts were explanted and a part of the left ventricle was shock frozen to be used for RNA isolation and Real time PCR analysis. The mid third of the remaining heart was fixed and transverse sections were prepared for histopathological analyses (hematoxylin Eosin, HE stain) or immunostained for collagen type 1 and alpha smooth muscle actin. Morphometric analysis was done using Precipoint software. Picrosirius red stain (SR) combined with imageJ based analyses reflected the total collagen content.

#### Results

T2DM manifestation was proven by significantly elevated blood glucose levels. Vit.K2 supplemented rats showed a trend of higher body weights and left ventricle wall thicknesses. Histopathological features of cardiomyopathy could be demonstrated in T2DM rats. The total collagen deposition was in mean higher in diabetic rats and lower in those with Vit.K2 supplementation. Fibronectin gene expression was significantly increased in the left ventricles of T2DM rats with no influence of Vit.K2 supplementation. In addition, a trend of more total collagen in hearts of T2DM rats compared to those of healthy controls became evident. This collagen deposition was lesser, if the rats were fed with Vit.K2 in both, T2DM and non T2DM rats. The collagen type 3 gene expression followed this trend of being higher in T2DM hearts but not the collagen type 1 immunoreactivity. The cytokine regulator gene suppressor of cytokine signaling (Socs) showed also elevated activity in left ventricles of T2DM compared to those of non T2DM rats.

#### Conclusions

The results suggest degenerative and early fibrotic changes in T2DM hearts but Vit.K2 supplementation did not significantly regulate the investigated molecular markers.

#### Acknowledgements

The work is financially supported by a Brückenschlag project founding. Tobias Braun's help in this project was founded by the budget of the Anatomical Center.

#### References

1. Kong et al., 2014, Cell Mol Life Sci 71:549-574. PMID: 23649149

#### Laboratory and Anamnestic Risk Indicator for Necrotizing Fasciitis

<u>David Breidung</u><sup>1</sup>, Gerrit Grieb<sup>2</sup>, Asja T. Malsagova<sup>1</sup>, Sarina Delavari<sup>1</sup>, Simonas Lapinskas<sup>1</sup>, Phillip von Imhoff<sup>1</sup>, Felix Ortner<sup>1</sup>, Maresa D. Berns<sup>1</sup>, Dayana Tsolova<sup>1</sup>, André A. Barth<sup>1</sup>, Alejandro Marti Edo<sup>1</sup>, Lena Schumann<sup>1</sup>, Tamas Püski<sup>1</sup>, Michaela Tutmann-Scheld<sup>1</sup>, Anke Wanninger<sup>1</sup>, Moritz Billner<sup>1</sup>, Wolfgang Hitzl<sup>3</sup>, Bert Reichert<sup>1</sup>, Ioannis-Fivos Megas<sup>1</sup>, Konrad Karcz<sup>1</sup>, Denis Ehrl<sup>1</sup>

<sup>1</sup>Department of Plastic, Reconstructive and Hand Surgery, Burn Unit, Klinikum Nürnberg, Paracelsus Medical University, Breslauer Str. 201, 90471 Nuremberg, Germany; <sup>2</sup>Department of Plastic Surgery and Hand Surgery, Gemeinschaftskrankenhaus Havelhoehe, Berlin, Germany; <sup>3</sup>Research and Innovation Management (RIM), Paracelsus Medical University Salzburg, Department of Ophthalmology and Optometry, Salzburg, Austria; Contact: david.breidung@icloud.com

#### Objective

Necrotizing fasciitis is a life-threatening soft tissue infection where the timing of surgical intervention significantly impacts clinical outcomes. The Laboratory Risk Indicator for Necrotizing Fasciitis (LRINEC) score, the Site other than the lower limb, Immunosuppression, Age <60 years, Renal impairment (creatinine >141), and Inflammatory markers (CRP  $\geq$ 150, WCC  $\geq$ 25) (SIARI) score, and procalcitonin levels are intended to assist in differentiating between necrotizing fasciitis and soft tissue infections.

#### Methods

A retrospective analysis of cases treated at Klinikum Nürnberg was conducted to evaluate necrotizing fasciitis and soft tissue infections. The diagnostic value of the two scores and procalcitonin levels was assessed, leading to the creation of a new score, the Laboratory and Anamnestic Risk Indicator for Necrotizing Fasciitis (LARINF), based on the findings.

#### Results

Procalcitonin, LRINEC score, and SIARI score exhibited inadequate discriminatory ability in our study. The newly devised LARINF score integrated laboratory parameters of the LRINEC score (hemoglobin and C-reactive protein) with procalcitonin and three comorbidities, resulting in a sensitivity of 84% and a specificity of 75%, with the highest area under the receiver operating characteristic (ROC) curve of 0.83.

#### Conclusions

The LARINF score presents a novel decision-supporting tool. The determination of when to initiate surgical exploration remains a clinical one. However, the score appears to offer an improved foundation for identifying this rare clinical condition.

#### References

 Wong CH, Khin LW, Heng KS, Tan KC, Low CO. The LRINEC (Laboratory Risk Indicator for Necrotizing Fasciitis) score: a tool for distinguishing necrotizing fasciitis from other soft tissue infections. Crit Care Med. 2004 Jul;32(7):1535-41. doi: 10.1097/01.ccm.0000129486.35458.7d. PMID: 15241098.. 2. Cribb BI, Wang MTM, Kulasegaran S, Gamble GD, MacCormick AD. The SIARI Score: A Novel Decision Support Tool Outperforms LRINEC Score in Necrotizing Fasciitis. World J Surg. 2019 Oct;43(10):2393-2400. doi: 10.1007/s00268-019-05061-4. PMID: 31214830.

## COST-EFFECTIVENESS OF ARTHROSCOPIC KNEE SURGERY IN DEGENERATIVE MENISCAL TEARS

Martin Breitwieser<sup>1</sup>, Thomas Freude<sup>1</sup>, Bernhard Rupp<sup>2</sup>, Katja Emmanuel<sup>1</sup>

<sup>1</sup>Department of Orthopaedic Surgery & Traumatology, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Healthcare & Sanitation, Chamber of Labor Lower Austria, St. Pölten; Contact: martin.breitwieser@stud.pmu.ac.at

#### Objective

Osteoarthritis [OA] of the knee is one of the most common orthopedic diagnoses in patients over the age of 40 years and represents a substantial burden of disease with clinical implications and accumulation of costs in the health care sector. Degenerative tears of the meniscus of the knee in these patients is a very common occurrence and should be treated conservatively or if this treatment plan fails with full or partial arthroplasty of the knee . Recent opposing studies show that knee arthroscopy can ease the pain for patients that fail to improve after conservative treatment and find relief when undergoing surgery (1). Therefore, knee arthroscopy might be more cost-efficient than conservative therapy in working adults with degenerative meniscus lesions.

#### Methods

A cohort of patients over the age of 40 years, who presented to the orthopedic clinic in Salzburg from 2010 to 2022 with degenerative meniscal tear undergoing either conservative or surgical treatment was modeled from clinical records. According administrative databases allowed the calculation of costs with exhaustive timedriven activity-based costing and a decision-analytic Markov disease progression model was constructed, using strategy-specific failure rates and treatment-specific probabilities for subsequent partial or total knee replacement to run a validated Monte Carlo simulation with a 5-year time horizon with a common endpoint of total or partial knee replacement surgery.

#### Results

Relative to conservative therapy, arthroscopic surgery is expected to show cost-effectiveness in more than 65% of simulations at willingness-to-pay thresholds of both  $\in$  50,000 per QALY and  $\in$  100,000 per QALY in sensitivity analysis. Arthroscopic surgery could be associated with an increased failure rate, but meaningful reductions in total or partial knee replacement therapy in the model-based analysis. Over a 10- year horizon, arthroscopic surgery is expected to show savings of up to  $\in$  3000 per patient, making it the dominant index procedure strategy.

#### Conclusions

Both interventions are associated with a clinically meaningful improvement regarding patients' most important functional limitations. Although arthroscopic surgery is associated with a statistically larger improvement, the difference compared with conservative therapy is small and clinically not meaningful in other clinical trials and studies. Other cost-effectiveness analysis suggest that arthroscopic surgery, despite substantially higher failure rates, is associated with improved long-term outcomes and cost savings relative to conservative treatment in the majority of patients, making it the dominant treatment strategy.

#### References

1. Rotini M, Papalia G, Setaro N, Luciani P, Marinelli M, Specchia N, et al. Arthroscopic surgery or exercise therapy for degenerative meniscal lesions: A systematic review of Systematic Reviews. MUSCULOSKELETAL SURGERY. 2022 Sept 3;107(2):127–41. doi:10.1007/s12306-022-00760-z
# HOW DO PARKINSON'S PATIENTS FROM SOUTH TYROL SLEEP?

## Susanne Buechner<sup>1,2</sup>, Agatha Eisendl<sup>1</sup>

<sup>1</sup>Department of Neurology, Provincial Hospital of Bolzano (SABES-ASDAA), Bolzano-Bozen, Italy <sup>2</sup>Teaching Hospital of the Paracelsus Medical University Salzburg, Bolzano, Italy Contact: susanne.buechner@sabes.it

### Objective

Sleep disturbances (SDs) are among the most common non-motor symptoms in patients with Parkinson's disease (PD).<sup>1</sup> The aim of the present study was to describe daytime sleepiness and sleep quality in a small cohort of South Tyrolean Parkinson's patients. This investigation is part of a large descriptive study aimed at characterizing parkinsonian patients living in South Tyrol.

### Methods

So far, a total of 22 Parkinson's patients (10 females and 12 males; Hoehn&Yahr (H&Y) stages 1-4; age 52-83 years) of the approximately 400 planned participants (recruitment until summer 2025) have completed the Epworth Sleepiness Scale (ESS) questionnaire, which assesses daytime sleepiness, and the Parkinson's Disease Sleep Scale (PDSS-2), which measures specific parkinsonian sleep problems.<sup>2,3</sup>

### Results

The mean ESS score was 8.14 (range 3-14), indicating a "higher normal daytime sleepiness"; there was no statistically significant difference between women and men (p=0.067). The highest probability of falling asleep was in the afternoon when patients lay down to rest. The mean PDSS-2 score was 16.18 (range 0-44), indicating a fairly good sleep quality, again with no significant difference between male and female patients (p=0.378). The most common nocturnal problems were sleeping through the night and getting up to urinate. Only seven patients (4 women with a mean age of 70.5 years and a mean H&Y stage of 2.37 and 3 men with a mean age of 74 years and a mean H&Y stage of 2.16) had a PDSS-2 score  $\geq$  18, indicating poor sleep quality; only two of them (one man and one woman) were taking melatonin. Only one other patient (a 74-year-old man with a H&Y stage of 3) was on specific sleep medication, namely with triazolam, which had resulted in an ESS score of 4 and a PDSS-2 score of 10.

# Conclusions

Preliminary findings indicate that no severe SDs were detected in the enrolled PD patients, regardless of gender. Overall, there was no evidence of excessive daytime sleepiness and only a few patients suffered from poor sleep quality, which did not clearly correlate with the PD's stage. Only three patients in total (13.64%) were taking sleeping pills, two of them with a PDSS-2 score above 18; both were taking melatonin. A comprehensive understanding, and a more robust picture of daytime sleepiness, sleep quality and therapy behaviour of the parkinsonian patients living in South Tyrol will only emerge once all 400 planned participants have been surveyed. In a second step, the final results will be compared with data from similar studies of other PD populations to assess the influence of geographical and cultural factors on sleep.

- Suzuki K, Miyamoto M, Miyamoto T, Iwanami M, Hirata K. Sleep disturbances associated with Parkinson's disease. Parkinsons Dis. 2011; 2011:219056. doi: 10.4061/2011/219056. Epub 2011 Aug 17. PMID: 21876839; PMCID: PMC3159123.
- 2. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. Sleep. 1991 Dec;14(6):540-5. doi: 10.1093/sleep/14.6.540. PMID: 1798888.
- Trenkwalder C, Kohnen R, Högl B, Metta V, Sixel-Döring F, Frauscher B, Hülsmann J, Martinez-Martin P, Chaudhuri KR. Parkinson's disease sleep scale--validation of the revised version PDSS-2. Mov Disord. 2011 Mar;26(4):644-52. doi: 10.1002/mds.23476. Epub 2011 Feb 10. PMID: 21312275.

# Generation of a Tendon-specific Knock-Out Mouse Model to Deeply Probe the Role of SPARC in Tendon Health and Disease

<u>Nevra Pelin Cesur<sup>1,2</sup></u>, Andrea Wagner<sup>1,2</sup>, Renate Gehwolf<sup>1,2</sup>, Andreas Traweger<sup>1,2</sup>

<sup>1</sup>Institute of Tendon and Bone Regeneration, Paracelsus Medical Private University, Salzburg, Austria; <sup>2</sup> Austrian Cluster for Tissue Regeneration, Vienna, Austria; Contact: nevra.cesur@pmu.ac.at

# Objective

Injured tendons mostly heal inadequately, leading to the formation of a stiff scar tissue with an altered extracellular matrix (ECM). These changes trigger degenerative cellular responses, ultimately driving tendon disease. Recently, the critical role of the ECM protein Sparc (Secreted protein acidic and rich in cysteine; BM-40 or osteonectin) in tendons has been described by our group (1,2). Sparc-null mice exhibit severely hypotrophic tendons, resulting in significantly reduced biomechanical strength and a higher incidence of spontaneous Achilles tendon rupture. Overall, our studies demonstrate that loss of Sparc results in a deteriorated, catabolic tendon phenotype, with altered sensitivity to mechanical stress and an impaired ECM. However, as these studies were conducted using a global Sparc KO model we cannot fully exclude, that the functional impairment is in part caused by loss of Sparc in e.g. bone or muscle in the tendon unit. Further, we are currently deciphering the potential role of Sparc during metabolic re-wiring in tendon degeneration. Therefore, we generated tendon-specific Sparc-null mice using the constitutive Scleraxis-Cre mouse line (Sparcfl/fl:Scx-Cre+).

# Methods

The successful loss of Sparc in tendons was examined by genotyping, gene expression analysis and Western blotting. Tendon phenotype was examined by gross morphology (e.g. Achilles length and width, tail tendon diameter), descriptive histology (e.g. Herovici's and Alcian blue staining), immunohistological stainings (e.g. Sparc, collagens) and biomechanical testing (e.g. tendon stiffness, maximal load).

# Results

We could confirm a significant reduction of Sparc in tendon tissues, both on the transcriptional and protein level. Additionally, Achilles tendon width is significantly smaller in Sparcfl/fl:Scx-Cre+ animals when compared to age-matched controls, whereas the tendon length remains unchanged. In addition, Sparcfl/fl:Scx-Cre+ load-bearing tendons display a severe hypoblastic phenotype, similar to Sparc-/- mice (1,2).

# Conclusions

In conclusion, Sparcfl/fl:Scx-Cre+mice are a valuable novel tendinopathy mouse model, avoiding the need for inducing tendon degeneration through ECM-degrading enzymes or subjecting mice to sustained mechanical overload. Therefore, this mouse model is an essential tool for investigating the cellular, molecular, and metabolic changes underlying degenerative tendon diseases and to further investigate the central role of Sparc in tendon biology.

- 1. Gehwolf R. et al; (2016); Sci Rep. 2016 Sep 2;6:32635
- 2. Wang T, Wagner A, Gehwolf R et al. (2021); Sci Transl Med. 2021 Feb 24;13(582):eabe5738

# Structural diversity and biophysical properties of voltage-gated proton channels (Hv)

### Gustavo Chaves<sup>1</sup>

<sup>1</sup>Center of Physiology, Pathophysiology and Biophysics, Paracelsus Medical University, Nuremberg, Germany; Contact: gustavo.chaves@pmu.ac.at

### Objective

The main objective of this study was to investigate the structural diversity and functional properties of voltagegated proton channels (Hv) within a single species, challenging the conventional dogma of a single Hv channel gene per species.

### Methods

GenBank (nr), transcriptional survey sequences (TSA), and expressed sequence tags (EST) databases were screened withthe TBLASTN algorithm for mollusc sequences with significant homology to known Hv1 sequences. Homology models for each paralog were constructed by homolgy modelling of the transmembrane domain of Ciona VSP. Molecular dynamics simulations for AcHv3 paralog were done. Hv gene paralogs were syntetized, cloned and heterologously expressed in tsA201 cells to further electrophysiological characterization.

### Results

Our study revealed the presence of novel paralogs of the typical Hv1 channel within a single species, indicating previously unrecognized diversity of a Hv family. These newly discovered Hv channels exhibit typical biophysical characteristics but also display subtle differences in functioning, potentially attributed to small but significant structural components. Structural analysis highlighted notable examples such as Hv2, which activates negatively to the Nernst potential for protons, enabling proton influx into the cell and membrane depolarization. Additionally, Hv3 and Hv4 paralogs exhibited distinct structural features compared to Hv1 or Hv2, such as variations in the voltage-sensing motif and disrupted channel behavior.

### Conclusions

1. The discovery of novel paralogs within the Hv channels challenges the traditional view of a unique Hvchannel gene per species, highlighting for the first time the existence of a voltage-gated proton channels family. 2. Structural analysis of these newly discovered paralogs, such as Hv2, Hv3, and Hv4, revealed distinct features and functional properties, including variations in gating charge carriers and altered voltage dependence, expanding our understanding of Hv channel diversity and function.

3. The findings provide valuable insights into the structural determinants underlying the functional diversity of Hv molecules, shedding light on their roles in various physiological processes and potentially informing future research and therapeutic strategies targeting Hv channels.

### Acknowledgements

Kerscher'sche Stiftung to GC

- Chaves G, Ayuyan AG, Cherny VV, Morgan D, Franzen A, Fieber L, Nausch L, Derst C, Mahorivska I, Jardin C, DeCoursey TE, Musset B. Unexpected expansion of the voltage-gated proton channel family. FEBS J. 2023 Feb;290(4):1008-1026. doi: 10.1111/febs.16617. Epub 2022 Sep 20. PMID: 36062330; PMCID: PMC10911540.
- Chaves G, Jardin C, Franzen A, Mahorivska I, Musset B, Derst C. Proton channels in molluscs: A new bivalvianspecific minimal HV 4 channel. FEBS J. 2023 Jul;290(13):3436-3447. doi: 10.1111/febs.16751. Epub 2023 Mar 3. PMID: 36788452.
- 3. Chaves G, Jardin C, Derst C, Musset B. Voltage-Gated Proton Channels in the Tree of Life. Biomolecules. 2023 Jun 24;13(7):1035. doi: 10.3390/biom13071035. PMID: 37509071; PMCID: PMC10377628.

# Sodium Valproate Modulates Cortical Morphology in Juvenile Myoclonic Epilepsy

<u>Bernardo Crespo Pimentel</u><sup>1,2,3,4</sup>, Giorgi Kuchukhidze<sup>1,4</sup>, Fenglai Xiao<sup>2,3</sup>, Lorenzo Caciagli<sup>2,3,5</sup>, Julia Höfler<sup>1,4</sup>, Lucas Rainer<sup>1,6</sup>, Christian Vollmar<sup>8</sup>, John Duncan<sup>2,3</sup>, Eugen Trinka<sup>1,4,9,10</sup>, Matthias Koepp<sup>2,3</sup>, Britta Wandschneider<sup>2,3</sup>

<sup>1</sup>Department of Neurology, Neurointensive Care and Neurorehabilitation, Christian Doppler University Hospital, Paracelsus Medical University, Salzburg, Austria;

<sup>2</sup>Department of Clinical & Experimental Epilepsy, UCL Queen Square Institute of Neurology, London, UK;

<sup>3</sup>Chalfont Centre for Epilepsy, Chalfont St. Peter, UK;

<sup>4</sup>Neuroscience Institute, Christian-Doppler University Hospital, Centre for Cognitive Neuroscience Salzburg, Austria; <sup>5</sup>Department of Neurology, Inselspital, Sleep-Wake-Epilepsy-Center, Bern University Hospital, University of Bern, Bern, Switzerland;

<sup>6</sup>Department of Child and Adolescent Psychiatry, Christian Doppler University Hospital, Paracelsus Medical University, Salzburg, Austria;

<sup>7</sup>Department of Psychology, University of Salzburg, Salzburg, Austria;

<sup>8</sup>Department of Neurology, Epilepsy Center, University Hospital of the Ludwig-Maximilians-University of Munich, Munich, Germany.;

<sup>9</sup>Department of Public Health, Health Services Research and Health Technology Assessment, UMIT- University of Health Sciences, Medical Informatics and Technology, Hall in Tirol, Austria;

<sup>10</sup>Karl Landsteiner Institute for Neurorehabilitation and Space Neurology, Salzburg, Austria;

Contact: bcpimentel93@gmail.com

### Objective

Juvenile myoclonic epilepsy (JME) is associated with cortical thinning of the motor areas (1). The relative contribution of anti-seizure medication to cortical thickness is unknown. We aimed to investigate how valproate influences cortical morphology of JME.

### Methods

In this cross-sectional study, individuals with JME with and without valproate, temporal lobe epilepsy (TLE) with valproate and controls were selected through propensity score matching. Participants underwent T1-weighted brain imaging and vertex-wise calculation of cortical thickness.

### Conclusions

Valproate was associated with JME-specific and dose-dependent thinning of the motor regions. This suggests that valproate may be a key modulator of cortical morphology in JME, on the mechanisms behind valproate's high efficacy in this syndrome.

### References

 Whelan CD, Altmann A, Botía JA, Jahanshad N, Hibar DP, Absil J, Alhusaini S, Alvim MKM, Auvinen P, Bartolini E, Bergo FPG, Bernardes T, Blackmon K, Braga B, Caligiuri ME, Calvo A, Carr SJ, Chen J, Chen S, Cherubini A, David P, Domin M, Foley S, França W, Haaker G, Isaev D, Keller SS, Kotikalapudi R, Kowalczyk MA, Kuzniecky R, Langner S, Lenge M, Leyden KM, Liu M, Loi RQ, Martin P, Mascalchi M, Morita ME, Pariente JC, Rodríguez-Cruces R, Rummel C, Saavalainen T, Semmelroch MK, Severino M, Thomas RH, Tondelli M, Tortora D, Vaudano AE, Vivash L, von Podewils F, Wagner J, Weber B, Yao Y, Yasuda CL, Zhang G, Bargalló N, Bender B, Bernasconi N, Bernasconi A, Bernhardt BC, Blümcke I, Carlson C, Cavalleri GL, Cendes F, Concha L, Delanty N, Depondt C, Devinsky O, Doherty CP, Focke NK, Gambardella A, Guerrini R, Hamandi K, Jackson GD, Kälviäinen R, Kochunov P, Kwan P, Labate A, McDonald CR, Meletti S, O'Brien TJ, Ourselin S, Richardson MP, Striano P, Thesen T, Wiest R, Zhang J, Vezzani A, Ryten M, Thompson PM, Sisodiya SM. Structural brain abnormalities in the common epilepsies assessed in a worldwide ENIGMA study. Brain. 2018 Feb 1;141(2):391-408. doi: 10.1093/brain/awx341. PMID: 29365066;

# Small-Molecule Library Screen Identifies Tanespimycin as a YAP-modulator in SH-SY5Y Cells

Daniel Devlitšarov<sup>1</sup>, Mirjana Malnar<sup>1</sup>, Nicole Schider<sup>1</sup>, Jan Pruszak<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cell Biology, Paracelsus Medical University, Salzburg, Austria; Contact: daniel.devlitsarov@pmu.ac.at

### Objective

Neuroblastoma (NB), the most common extra-cranial solid tumor in children, displays a heterogeneous clinical spectrum due to its diverse cellular composition. Because of the phenotypic variability presented by this disease, timely diagnosis, accurate prognosis and choice of treatment remain challenging. Despite recent advances in treatment approaches, current therapies often fall short of achieving complete tumor eradication, highlighting the urgent need for novel therapeutic approaches that target high-risk therapy-resistant NB, and selectively modulate tumor behavior. A preliminary study conducted by Ferlemann et al. provided the first comprehensive flow cytometric characterization of surface molecule expression in NB cell lines, and employed an unbiased clustering algorithm (SPADE) to resolve cellular heterogeneity and identify distinct neuroblastoma subsets based on combinatorial expression of CD15, CD24, CD44, CD57, CD184, and TrkA [1].Building on that study, we screened a small-molecule library consisting of protein-protein interaction inhibitors, using the same methodology as the preliminary investigation. The goal of the screen was to identify drug candidates capable of modulating NB surface marker expression profiles.

### Methods

SH-SY5Y NB cells were pre-cultured in DMEM/F12 supplemented with 10% FBS and 1% Pen/Strep until reaching 85% confluence. Cells were harvested with TrypLE and used to seed 6-well plates. After 24 hours, medium was exchanged for DMEM/F12 supplemented with 10% KnockOut Serum Replacement. Tanespimycin was added to a 1 $\mu$ M final concentration. DMSO served as the vehicle control, at a final concentration of 0.01%. Medium was changed after 24 hours and fresh treatment compounds were added. After 48 hours of treatment, cells were harvested, labeled with fluorescent antibodies and analyzed by flow cytometry.

### Results

Among the potential therapeutic candidates, we identified tanespimycin (17-AAG), as a modulator of the YAPpositive phenotype in SH-SY5Y cells. Tanespimycin significantly altered the composition of specific NB subpopulations in-vitro, as evidenced by downregulation of CD44 and CD49d (markers associated with a YAPpositive phenotype [2]). Upregulation of CD184 and CD24 was also observed. The selective regulation of YAPpositive subpopulations by tanespimycin expands our understanding of this drug's mechanism of action in NB cells.

### Conclusions

This study builds upon the groundwork laid by the preliminary investigation [1], advances our insights into NB cellular heterogeneity, and corroborates the potential of tanespimycin as a selective modulator of surface marker expression and the YAP pathway. These findings highlight the significance of tanespimycin as a promising addition to our therapeutic arsenal for combating this challenging pediatric malignancy.

- Ferlemann FC, Menon V, Condurat AL, Rößler J, Pruszak J. Surface marker profiling of SH-SY5Y cells enables small molecule screens identifying BMP4 as a modulator of neuroblastoma differentiation. Scientific Reports. 2017 Oct 19;7(1).
- 2. Hindley CJ, Condurat AL, Menon V, Thomas R, Azmitia LM, Davis JA, Pruszak J. The Hippo pathway member YAP enhances human neural crest cell fate and migration. Scientific Reports. 2016 Mar;6(1).

# Effect of Probiotic Supplementation on the Intestinal Microbiome of Preterm Infants

<u>Ronald Ebbert</u><sup>1</sup>, Sabrina Hofmann<sup>1</sup>, Adrian Rosenberg-Granzner<sup>1</sup>, Krisztina Somogyi<sup>1</sup>, Niels Rochow<sup>2</sup>, Katja Knab<sup>2</sup>, Christoph Fusch<sup>2</sup>

<sup>1</sup>TH Nürnberg Georg Simon Ohm, Fakultät AC;

<sup>2</sup>Department/Institute of Neonatology, Children and Adolescents, Paracelsus Medical University, Nuremberg, Germany; Contact: ronald.ebbert@th-nuernberg.de

## Objective

In the current study, 30 preterm infants (mean gestational age 205 d) received the probiotic ProPrems (1) containing Bifidobacterium longum subsp. infantis, Streptococcus thermophilus and Bifidobacterium animalis subsp. lactis. 14 Patients without supplementation (mean gest. age 237 d) served as control.

## Methods

Longitudinal stool samples were collected: 28 samples before, 118 during and 69 after supplementation and 44 of the control group. Bacteria were identified by Nanopore sequencing of the complete 16S gene. Sequence analysis was performed by a workflow based on QIIME 2.

## Results

PCoA based on Bray-Curtis distances, PERMANOVA and ANCOM analysis show significantly higher abundances of the ProPrems species "During" supplementation compared to "Before". Unexpectedly, both S. thermophilus and B. animalis are strongly reduced in "After" samples compared to "During".

### Conclusions

All three probiotic strains are found during supplementation, but apparently only B. longum subsp. infantis successfully colonizes the preterm gut.

# References

1. Garland SM, Tobin JM, Pirotta M, Tabrizi SN, Opie G, Donath S, Tang ML, Morley CJ, Hickey L, Ung L, Jacobs SE; ProPrems Study Group. The ProPrems trial: investigating the effects of probiotics on late onset sepsis in very preterm infants. BMC Infect Dis. 2011 Aug 4;11:210. doi: 10.1186/1471-2334-11-210. PMID: 21816056; PMCID: PMC3199779.

# Sunflower Syndrome – Creating Awareness & International Medication Survey

### Kirsten Eschermann<sup>1</sup>

<sup>1</sup>Research Institute for Rehabilitation, Transition and Palliation; PMU Salzburg; Austria; Contact: kirsten.eschermann@pmu.ac.at

## Objective

Sunflower syndrome (SFS) is a rare epileptic disorder characterized by pronounced stereotyped seizures. People with SFS look at a light source, usually the sun, and wave a hand in front of their eyes. In connection with these hand-waving episodes (HWE) eyelid myoclonus can occur. In addition to HWE, other forms of seizures such as forms such as absences and generalized tonic-clonic seizures. A genetic cause is suspected, but has not yet been clarified.Objectives of our study:1. AwarenessSFS is insufficiently recognized and is not described in detail in the literature. HWE is often incorrectly classified as a tic disorder or obsessive-compulsive disorder. This lack of awareness and understanding, both among healthcare professionals and the general public, has led to many patients developing a false self-perception and therefore poor self-esteem. With our survey, we would like to raise awareness of the condition among both the treating physicians and the patients' social environment.2. Effectiveness of medicationUnfortunately, SFS often proves to be resistant to treatment to the currently available anticonvulsant treatments. Non-pharmacological stimulus avoidance measures such as wearing hats and sunglasses can be beneficial, but these strategies alone do not lead to seizure freedom. Our platform PATRE (PATient based phenotyping and evaluation of therapy in Rare Epilepsies; www.patre.info) provides a standardized procedure for researching therapies for rare epilepsies based on information provided by patients and their parents.

### Methods

The PATRE project for the SFS is a cooperation between the University of Leuven, the University of Genoa, Harvard University and the PMU Salzburg. For our electronic questionnaires, we use our PATRE server with REDCap for secure data collection. In an initial survey, we asked patients and their parents about their subjective assessment of the effect of medication on SFS. The effect on seizures and behavior was measured on a scale of -50 to +50.

### Results

So far, 14 patients have taken part in the study. Medications for which at least two patients had provided information were evaluated. Treatment effects on seizures with valproic acid and levetiracetam have been described in the literature. For this reason, the poor performance of levetiracetam in our group is particularly surprising. According to our survey, ethosuximide and lamotrigine appear to achieve good anticonvulsant results. The positive anticonvulsant effect of valproic acid was confirmed. Lamotrigine and valproic acid also had a good effect on behavior, as did clobazam.

### Conclusions

These results are only based on data from a small number of patients and are therefore only of limited value. We are continuing the survey and would be very pleased if you would contact us at sunflower@patre.info if you care for patients with Sunflower syndrome.

### References

1. Nascimento FA, Thiele EA. Teaching Video NeuroImage: Shedding Light on Sunflower Syndrome. Neurology. 2021 Aug 17;97(7):e749. doi: 10.1212/WNL.000000000011776. Epub 2021 Mar 15. PMID: 33722995.

# Prenatal Psychosocial Risk: Association with Stress, Social Support and Resilience

<u>Figen Isik Esenay</u><sup>1,2,3</sup>, Alina Oesterreich<sup>1,3</sup>, Barbara Strubegger<sup>4</sup>, Elisabeth Bürgler<sup>5</sup>, Ulrike Metzger<sup>6</sup>, Gabriela Markova<sup>1,3,7</sup>, Beate Priewasser<sup>1,3</sup>

<sup>1</sup>Institute for Early Life Care, Paracelsus Medical University, Salzburg, Austria;

<sup>2</sup>Department of Pediatric Nursing, Faculty of Nursing, Ankara University, Ankara, Turkey;

<sup>3</sup>Department of Pediatrics, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria;

<sup>4</sup>Institute of Psychology, Paris Lodron University Salzburg, Salzburg, Austria;

<sup>5</sup>University Clinic for Gynecology and Obstetrics, University Hospital Salzburg, Salzburg, Austria;

<sup>6</sup>University Clinic for Pediatric and Adolescent Surgery, University Hospital Salzburg, Salzburg, Austria;

<sup>7</sup>Faculty of Psychology, University of Vienna, Vienna, Austria;

Contact: figen-isik.esenay@pmu.ac.at

# Objective

Psychosocial stress in parents is an important risk factor for child development. Already prenatal maternal stress can affect fetal brain development (1). In addition, postpartum psychosocial stress can negatively affect caregiving behavior toward the infant, which can, in turn, lead to impairments in child development. Despite these serious and long-term effects, the risk factor of psychosocial stress in parents remains widely neglected in primary care and no screening instruments are available that could assess psychosocial stress quickly and resource-effectively. The aim of the study is to validate a short screening, consisting of seven evidence-based questions, can accurately assess psychosocial stress among expectant mothers.

## Methods

The study sample is part of a research project "Validierung von Screening-Fragen zur Erfassung von Psychosozialen Risikofaktoren" at the Early Life Care Institute, PMU. We recruited 80 pregnant women in the 37th-39th week of pregnancy during birth registration at the University Hospital Salzburg. Participating women were on average 33.4 (SD = 1.59) years old and in the 38th (SD = 4.22) gestation week. At time of recruitment, participants filled out online questionnaires to assess their psychosocial risks, perceived stress, social support and resilience.

# Results

Correlation analysis showed that higher psychosocial risk was associated with higher perceived stress ( $\beta$ =.673; t=7.117; p<.001), lowered perceived social support ( $\beta$ =-.247; t = -2.915, p=.005) and higher resilience ( $\beta$ =.223; t = 2.578, p=.012). Perceived stress, social support and resilience together explained 54.6% of the variance in psychosocial risk (R2 = .546; F=31.253; p<.001).

### Conclusions

Our findings indicate that a very short screening instrument may be suitable to assess psychosocial risk in expectant mothers. The use of universal screening in primary care practice can remove the stigma of screening selected families and reduce the likelihood of overlooking at-risk families.

### Acknowledgements

The project is financially supported by Gesundheit Österreich GmbH (as part of the EU program "NextGenerationEU" and commissioned by the BMSGPK), the Ferry Porsche Foundation, and the Austrian Society for Pediatrics and Adolescent Medicine.

### References

1. Buss et al. (2012). Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumesand affective problems. Proceedings of the National Academy of Sciences of the United States of America, 109(20), E1312-1319. https://doi.org/10.1073/pnas.1201295109

# FROM LAPTOP TO TOP-LAB: A COMBINED IN SILICO IN VITRO APPROACH TO IDENTIFY NOVEL MICROTUBULE STABILIZERS AS POTENTIAL DRUG CANDIDATES

Florian Fischer<sup>1</sup>, Daniela Schuster<sup>1</sup>, Veronika Temml<sup>1</sup>

<sup>1</sup>Institute of Pharmacy, Department of Pharmaceutical and Medicinal Chemistry; Contact: florian.fischer@pmu.ac.at

# Objective

Microtubules are integral components of eukaryotic cell structures, providing essential support and facilitating various cellular processes. Composed of  $\alpha$ - and  $\beta$ -tubulin subunits, they form dynamic filaments crucial for structural integrity, intracellular transport, and cell division. Through dynamic polymerization and depolymerisation of these tube-like aggregates, these proteins represent an important element in the life cycle of eukaryotic cells. Disruption of microtubule dynamics can induce cell death, particularly in rapidly dividing cells like cancer or immune cells, by promoting apoptosis. One strategy involves inhibiting continuous depolymerization using microtubule stabilizers from the taxane family, such as paclitaxel and docetaxel. However, resistance to these potent chemotherapeutics can develop in cancer cells. Consequently, there is considerable scientific interest in identifying new compound classes and scaffolds with potential antitumor activity through microtubule stabilization, offering promising avenues for cancer therapeutics.[1, 2]

### Methods

Our approach combines innovative in silico techniques such as pharmacophore modeling with in vitro studies to discover new enhancers for the  $\beta$ -tubulin subunit. High-quality pharmacophore models were generated using LigandScout version 4.4.5 [3]. Pharmacophore modeling identifies key physicochemical features of protein-ligand interactions, facilitating the screening of molecules with similar interaction patterns. In this study, a combination of structure- and ligand-based modeling was employed. Structure-based models utilized 3D crystal structures, while ligand-based pharmacophore models were developed by aligning active molecules from the literature. The optimized and theoretically validated pharmacophore models were used to screen the SPECS database, which contains > 200,000 commercially available compounds (april2023). Selected Hits identified by our models were further investigated in vitro using a fluorescence-based tubulin polymerization assay kit (Cytoskeleton, BK011P).

### Results

A comprehensive literature search was conducted to compile microtubule stabilizers targeting the taxane binding site of the beta-tubulin subunit, resulting in a curated set of 56 active compounds. Additionally, a decoy set of 1907 compounds was generated for comparison. Using LigandScout 4.4.5, databases were converted into multi-conformational screening datasets, facilitating the training of pharmacophore models. A dataset consisting of 56 biologically tested microtubule stabilizers (MSTs) and 1907 decoy compounds was transformed into a 3D dataset. One structure-based (SB) and three ligand-based (LB) pharmacophore models were utilized to identify active substances from the SPECS database. Among 3616 candidates, 22 were screened in vitro at a concentration of 30  $\mu$ M, with 8 exhibiting significantly higher polymerization rates compared to the blank. This integrated in silico and in vitro approach led to the discovery of novel microtubule stabilizers, which will undergo further investigation using cancer cell models

### Conclusions

Our highly optimized structure-based and ligand-based pharmacophore models have the capability to predict substances that also exhibit the desired activity in an in vitro assay with a success rate of 36%. To demonstrate the potency of the top-ranked biologically tested compounds, further concentration-dependent screenings are now required.

- 1. Cao YN, Zheng LL, Wang D, Liang XX, Gao F, Zhou XL. Recent advances in microtubule-stabilizing agents. Eur J Med Chem. 2018 Jan 01;143:806-28.
- 2. Lei L, Wang XJ, Tang SC. Novel taxanes in development: Hopes or hypes? Crit Rev Oncol Hematol. 2022 Aug;176:103727.
- 3. Wolber G, Langer T. LigandScout: 3-D pharmacophores derived from protein-bound ligands and their use as virtual screening filters. J Chem Inf Model. 2005;45(1):160-9.

# Safe Forests: The psychological wellbeing of harvester and forwarder drivers – a cross-sectional study

Vera Foisner<sup>1</sup>, Christoph Haas<sup>2</sup>, Christoph Huber<sup>2</sup>, Arnulf Hartl<sup>1</sup>

<sup>1</sup>Institute of Ecomedicine, Paracelsus Medical University Salzburg, Austria; <sup>2</sup>Austrian Research Centre For Forests, Upper Austria, Austria; Contact: vera.foisner@pmu.ac.at

## Objective

The use of fully mechanized wood harvesting systems allows forwarder and harvester machines to operate in steep terrain, thereby reducing harvesting costs and rising work safety for the operators. But working in such challenging conditions poses hazards and stressors for the drivers, such as high levels of mental stress and strain. We focused on the subjective wellbeing, stress, and strain the drivers are facing during their work and compared them to the average population. Aiming to provide a better understanding of their psychological wellbeing and providing better work-time-organisation.

## Methods

Online-Questionnaire: SF-12 [1] to measure physical and mental health, RESTQ-work 27 [2] to record recovery and strain at work, OrgFit [3] for risk assessment of mental stress and work-related strain, and environmental factors, such as steepness of the terrain. Setting: recruitment by calling companies, link on webpage. Participants: mainly male healthy forestry workers in Austria, using a) harvester and b) forwarder machines with an age between 18-55 years. Analyses: parametric (unpaired t-test ( $\alpha$ = 0.05, 95% CI)) and non-parametric (Spearman correlation) tests.

### Results

Population: 42 forestry workers (one women) with an average age of 32.2 years (SD 9.26), 88.62 kg (SD 16.44), and 27.11 BMI (SD 4.28) participated. Comparison to norm samples: participants (M= 53.94; SD = 3.39) reported a significantly higher (t (2845)= 3.39; df= 2845; p < 0.001; d= 0.69) physical health (SF-12) compared to the norm sample (M= 49.03; SD= 9.35), comprising of a representative German general population (n= 2524). According to Cohen (1992), this is a medium to high difference.Other factors: Environmental factors have no effect on the driver's wellbeing. The higher the assessed difficulty of the last operation, the steeper the terrain (r= 0.6757; p< 0.001) (Fig. 1). And the better their social and organisational climate (OrgFit), the higher their recovery from stress and strain (RESTQ-work 27) (r= 0.6484, p< 0.001) (Fig. 2). According to Cohen (1992), both are strong correlations.

# Conclusions

being or working in nature benefits human health: shown by higher subjective physical health and recovery from stressSocial interactions of importance for psychological wellbeing: most participants work in small, family-like companies; they showed better social and organisational climate, which was also associated with higher recoveryfurther analysis with higher number of participants is of need

- 1. Drixler et al., 2020
- 2. Jiménez & Kallus, 2016
- 3. Jiménez et al, 2022

# A comparative study of three different bioactive glass scaffolds tailored for cartilage tissue engineering

<u>Clemens Gögele<sup>1</sup></u>, Silvana Müller<sup>2</sup>, Sven Wiltzsch<sup>2</sup>, Kerstin Schäfer-Eckart<sup>3</sup>, Bernd Minnich<sup>4</sup>, Astrid Obermayer<sup>4</sup>, Armin Lenhart<sup>2</sup>, Gundula Schulze-Tanzil<sup>1</sup>

<sup>1</sup>Institut für Anatomie und Zellbiologie, Paracelsus Medizinische Universität, Salzburg und Nürnberg, Nürnberg, Deutschland; <sup>2</sup>Fakultät für Werkstoffwissenschaften, Technische Hochschule Nürnberg Georg Simon Ohm, Nürnberg, Deutschland; <sup>3</sup>Klinik für Onkologie/Hämatologie, Klinikum Nürnberg Nord, Nürnberg, Deutschland; <sup>4</sup>Department für Umwelt und Biodiversität, Paris Lodron Universität Salzburg, Salzburg, Österreich; Contact: clemens.goegele@pmu.ac.at

## Objective

The hyaline cartilage has a reduced regenerative capacity. In order to prevent the progression of osteoarthritis, cartilage defects should be treated. Cartilage tissue engineering using bioactive glass (BG) is a promising approach. Until now, conventional BG (45S5, 60S5, BG1393) has been used for bone regeneration due to the formation of hydroxyapatite. The aim of this in vitro study is to compare two BGs with a novel ionic composition, specifically developed for cartilage tissue with the conventional BG1393.

### Methods

The novel BG CAR12N (1) variants were activated by leaching with HCI. The highly porous and 100% bioglass scaffolds (CAR12N, CAR12 + Ca2+/Mg2+, BG1393) with the same topology) were seeded with either primary porcine articular chondrocytes (pACs) or human mesenchymal stem cells (hMSCs) and cultured dynamically for 21 days. Subsequently, the sulfated glycosaminoglycan (sGAG) content per cell was determined quantitatively, the vitality assay was performed qualitatively and quantitatively, and scanning electron micrographs (SEM) were taken. Cartilage-specific protein expressions (collagen type II, proteoglycans, SOX9 and collagen type I) were visualized by immunocytochemistry.

### Results

Macro- and microscopic images showed that the leaching did not lead to any structural changes in the scaffold and that the porosity was above 75%. During leaching, the pH value changed in both HCl and water. The viability assay showed that both pACs and hMSCs can survive on all three scaffold variants. The sGAG expression of pACs increased in trend on all glass variants. After 21 days, expression is highest on the BG1393 scaffolds colonized with pACs. A significant increase was shown in the BG1393 glass at this time. The sGAG expression of the hMSCs is significantly reduced compared to the pACs. The SEM showed deep insights into the cell-material interaction as well as the surface morphology of the scaffolds. The expression of collagen type II, cartilage-specific proteoglycans and the transcription factor SOX9 was detected in pACs and hMSCs colonized CAR12N scaffolds.

### Conclusions

Highly porous bioactive glass scaffolds with a comparable morphology could be produced and cultured with mesenchymal cells for several weeks. Although the BG1393 scaffolds showed the densest cell growth, it was shown that the expression of cartilage markers and therefore the onset of successful chondrogenese, was highest on pACs CAR12N scaffolds but was also seen in hMSCs. This is currently being confirmed by gene expression analysis.

### Acknowledgements

The work is financially supported by the OPED GmbH.

### References

 Goegele C, Wiltzsch S, Lenhart A, Civilleri A, Weiger TM, Schaefer-Eckart K, et al. Highly porous novel chondroinstructive bioactive glass scaffolds tailored for cartilage tissue engineering. Mater Sci Eng C Mater Biol Appl. 2021;130:112421. PMID: 34702508

# Chondrogenesis of hMSCs in 3D printed bioglass/hydrogel scaffolds for cartilage regeneration

<u>Clemens Gögele<sup>1</sup></u>, Nils Fleischmann<sup>1</sup>, Silvana Müller<sup>2</sup>, Tanja Liesenberg<sup>1</sup>, Gina Pizzadili<sup>1</sup>, Sven Wiltzsch<sup>2</sup>, Armin Lenhart<sup>2</sup>, Christian Werner<sup>1</sup>, Kerstin Schäfer-Eckart<sup>3</sup>, Gundula Schulze-Tanzil<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cell Biology, Paracelsus Medical University, Nuremberg and Salzburg, 90419 Nuremberg, Germany; <sup>2</sup>Faculty of Materials Science, Technische Hochschule Nürnberg Georg Simon Ohm, 90489 Nuremberg, Germany; <sup>3</sup>Bone marrow Transplantation Unit, Medizinische Klinik 5, Klinikum Nürnberg, Paracelsus Medical University, 90419 Nuremberg, Germany; Contact: clemens.goegele@pmu.ac.at

### Objective

The self-healing capacity of cartilage tissue is low after injury. For this reason, there is still an urgent need to develop a new, cost-effective, implantable and chondroinductive matrix with mesenchymal stem cells (hMSCs) that induces cartilage formation. In contrast to conventional bioglasses, which are designed for bone regeneration our glass (CAR12N) (1) has a faster degradation rate and no hydroxyapatite deposition. This basic research project focuses on cartilage formation (chondrogenesis) in 3D printed bioglass/hydrogel scaffolds for the precise and personalized regeneration of cartilage defects.

### Methods

The grid-like scaffolds (7 x 7 x 1.5 mm), consisting of a hydrogel (1.3% alginate with methyl cellulose without/with CAR12N glass spheres (25-40 µm)) and hMSCs, are produced using a "layer-to-layer" process with the 3D printer and have a porosity of 70%. After scaffold printing, they are cross-linked and then cultivated statically in different culture media. In order to determine the influence of the ions on the MSCs, the scaffolds are cultured either: in isolation medium, in chondrogenic differentiation medium. After long-term cultivation of up to 21 days, extensive analyses (morphological examinations, vitality assay, metabolic activity assay, DNA and sulfated glycosaminoglycan (sGAG) content, chondrogenic matrix components at protein and gene level were carried out in order to be able to adequately assess the progress of chondrogenesis.

#### Results

Extensive morphological studies have shown that the hMSC-colonized scaffolds remain dimensionally stable even after 21 days in the different media. In general, significantly more hMSCs survived over time in the isolation medium than in the chondrogenic differentiation medium. The bioactive glass spheres influence cell viability. However, the glass spheres do apparently influence proliferation and sGAG synthesis as these were always higher in the presence of the spheres compared to the control groups. Increased relative gene expression of cartilage-specific matrix components, such as aggrecan and collagen type II, was detected in hydrogel scaffolds cultured in chondrogenic differentiation medium compared to the isolation medium.

### Conclusions

A printed 3D hydrogel model made of bioink alginate/methylcellulose was established for the investigation of MSC chondrogenesis. The functionalization with bioactive glass spheres enhances the effect on vitality, proliferation and sGAG synthesis of primary hMSCs. In the future, the quantity ratio and the printing pressure parameters are to be coordinated even more precisely in order to achieve faster chondrogenic differentiation.

#### Acknowledgements

The work is financially supported by the Förderverein der Paracelsus Medizinischen Privatuniversität.

#### References

 Goegele C, Wiltzsch S, Lenhart A, Civilleri A, Weiger TM, Schaefer-Eckart K, et al. Highly porous novel chondroinstructive bioactive glass scaffolds tailored for cartilage tissue engineering. Mater Sci Eng C Mater Biol Appl. 2021;130:112421. PMID: 34702508

# Hiatal hernia after transthoracic esophagectomy for cancer: the key role of omentectomy

Luca Giulini, Irina Avramovska<sup>1</sup>, Lucas Thumfart<sup>1</sup>, Felix Hüttner<sup>1</sup>, Wolfgang Hitzl<sup>2</sup>, Markus Diener<sup>1</sup>, Attila Dubecz<sup>1,3</sup>

<sup>1</sup>Department of General, Visceral and Thoracic Surgery, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Biostatistics and Publication od Clinical Trial Studies, Research and Innovation Management (RIM), Paracelsus Medical University, Nuremberg, Germany;

<sup>3</sup>Department of Surgery, Helios Clinic Erfurt, Academic Hospital of the University of Jena, Erfurt, Germany; Contact: Luca.Giulini@klinikum-nuernberg.de

### Objective

Hiatal Hernia after esophagectomy for cancer represents a potentially life-threatening long-term complication of which pathogenesis is unclear. However, it is well-known that its incidence is higher after minimally-invasive procedures. Aim of this single-center controlled cohort-study was to compare the incidence of hiatal hernia after open vs minimally-invasive esophagectomy (MIE), and to identify specific risk factors associated with its occurrence. Furthermore, the objective was to validate our hypothesis that the remaining omentum after MIE has a key role in the development of a post-esophagectomy hiatal hernia.

### Methods

Our prospectively maintained database was retrospectively queried for patients who underwent transthoracic esophagectomy (Ivor Lewis or McKeown) for cancer over a 15 years period. Groups were defined according to the procedure (open = omentectomy vs. minimally-invasive = non-omentectomy). Minimally-invasive procedures were performed both laparoscopic and robotically- assisted. In case of a conversion to laparotomy the patient was included in the open group. The 2 groups were compared according to demographics, operative, histologic and postoperative parameters. Risk factors for hiatal hernia after esophagectomy were analyzed.

### Results

A total of 897 patients were included. A hiatal hernia was registered in 1/490 (0.2%) in the open and in 21/407 (5.16%) of the minimally-invasive treated patients [Odds ratio = 26.6 (95% CI: 3.56-198.7), p = 0.0001]. Otherwise, significant differences (two-tailed p<0.05) between the two groups were noted only in 8 out of further 31 investigated variables, as a parameter of homogeneity between the groups. Within the minimally-invasive group, patients with ASA Score of 2 and 3 had significantly lower risk to develop a hiatal hernia when compared to ASA 1 subjects (Odds ratio 0.18, 95% CI 0.06 – 0.55, p=0.002 and 0.11, 95% CI 0.03 – 0.39, p=0.0007 respectively). No further risk factors were identified. All patients underwent an omentectomy in the open, and none in the minimally-invasive group.

### Conclusions

In our cohort, hiatal hernia was significantly lower in patients who underwent open esophagectomy compared to MIE. Omentectomy might play a key role in the prevention of post-esophagectomy hiatal hernia and should thus be considered also during minimally-invasive procedures. However, further multicentric and randomized studies are needed.

- 1. Kuvendjiska J et al. Postoperative Hiatal Hernia after Ivor Lewis Esophagectomy-A Growing Problem in the Age of Minimally Invasive Surgery. J Clin Med. 2023;12(17):5724. Published 2023 Sep 1. PMID: 37685791
- 2. Murad H et al. Postoperative hiatal herniation after open vs. minimally invasive esophagectomy; a systematic review and meta-analysis. Int J Surg. 2021;93:106046. PMID: 34411750
- Fetzner UK et al. A commentary on "Postoperative hiatal herniation after open vs. minimally invasive esophagectomy; a systematic review and meta-analysis" (Int J Surg 2021;93:106046). Paraconduit hiatal hernia: A relevant complication after radical esophagectomy with reconstruction via gastric pull-up. Suggestions for prevention. Int J Surg. 2022;102:106659. PMID: 35589050
- 4. Hertault H et al. Incidence and Risk Factors for Diaphragmatic Herniation Following Esophagectomy for Cancer. Ann Surg. 2021;274(5):758-765. PMID: 34334646

# IMPACT OF ANASTOMOTIC LEAK VS PNEUMONIA ON FAILURE TO RESCUE AFTER TRANSTHORACIC ESOPHAGECTOMY FOR CANCER

Luca Giulini (MD, FACS)<sup>1</sup>, Melissa Kemeter (MD)<sup>1</sup>, Lucas Thumfart (MD)<sup>1</sup>, Felix J. Hüttner (MD)<sup>1</sup>, Patrick Heger (MD)<sup>1</sup>, <u>Filitsa Farmaki (MD)</u>, Oliver Koch (MD, FEBS)<sup>2, 6</sup>, Michael Grechenig (MD)<sup>2</sup>, Klaus Emmanuel (MD, FACS)<sup>2, 6</sup>, Wolfgang Hitzl (PhD)<sup>3, 4, 5</sup>, Markus K. Diener (MD, MBA)<sup>1, 6</sup>, Attila Dubecz (MD, FACS)<sup>6, 7</sup>

1. Department of Surgery, Paracelsus Medical University Nuremberg, Nuremberg, Germany. 2. Department of General, Visceral and Thoracic Surgery, Paracelsus Medical University Salzburg, Salzburg, Austria. 3. Biostatistics and Publication of Clinical Trial Studies/Machine Learning, Research and Innovation Management (RIM), Paracelsus Medical University Salzburg, Salzburg, Austria. 4. Department of Ophthalmology and Optometry, Paracelsus Medical University Salzburg, Salzburg, Austria. 5. Research Program Experimental Ophthalmology and Glaucoma Research, Paracelsus Medical University Salzburg, Salzburg, Salzburg, Austria. 6. Paracelsus Medical University, Nuremberg (Germany) and Salzburg (Austria). 7. Department of Surgery, Helios Clinic Erfurt, Academic Hospital of the University of Jena, Erfurt, Germany Contact: filitsafarmaki@gmail.com

# Objectives

Data about failure-to-rescue (FTR) after oesophagectomy for cancer and its association with patient and procedure-related risk factors are sparse. Aim of this study was to analyze such aspects, particularly focusing on the impact of pneumonia and anastomotic leak on FTR.

## Methods

Patients who underwent an Ivor Lewis oesophagectomy for cancer between 2008 and 2022 in two tertiary European centers were prospectively identified. Patients were classified and compared according to the type of operation (open, hybrid-laparoscopic, hybrid-robotic, minimally-invasive or robotic minimally-invasive). FTR was defined as in-hospital death following a major complication. Risk factors for FTR were identified with a univariable model. Mortality after pneumonia and anastomotic leak were calculated and compared across the groups.

# Results

Sevenhundred-and-eight patients were included. There were 355 (50.1%) open, 204 (28.8%) hybrid-laparoscopic, 121 (17.1%) hybrid-robotic, 15 (2.1%) standard minimally-invasive and 11 (1.6%) robotic minimally-invasive procedures. Overall morbidity was 60%, and failure-to-rescue rate 4.5%. Anastomotic leak, pneumonia, postoperative bleeding, sepsis, pulmonary embolism, arrhythmia and need for blood transfusion were the risk factors significantly associated with in-hospital mortality (p<0.05). There was no particular type of operation significantly associated with mortality (p=0.42). Pneumonia and leak associated FTR rates did not significantly differ among the groups (p=0.99).

# Conclusions

Anastomotic leak and pneumonia are equally dangerous complications after oesophagectomy for cancer. Hybrid or minimally-invasive methods do not seem to negatively affect the FTR rates. However, implementation of minimally-invasive oesophagectomy should be considered carefully and only in high-volume centers, whereas efforts should be made to improve both tailored-approach and postoperative care.

# Mixed Methods Evaluation of Early Home-Based Interventions for Children at Risk

Elke Greif<sup>1,2</sup>, Antonia Dinzinger<sup>1,2</sup>, Beate Priewasser<sup>1,2</sup>

<sup>1</sup>Institute for Early Life Care, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Pediatrics, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria; Contact: elke.greif@pmu.ac.at

### Objective

In early childhood, the brain is especially malleable, making early experiences crucial for later development (1). For example, the lack of appropriate treatment for visual disorders in a very limited time period in infancy can affect the entire life course, compromising the development of the visual system as well as motor, cognitive and emotional development (2), highlighting the need for early intervention. Several organizations in Vienna, funded by the Vienna Social Fund, provide early childhood interventions for children with a variety of disabilities or developmental risks and their families. These home-based interventions, delivered by trained facilitators, include not only targeted exercises with the children, but also support and guidance for parents in dealing with their child's impairment. The facilitators assist parents in handling bureaucratic barriers and societal reactions to the impairment, which parents of disabled children have identified as major obstacles (3). The aim of this study is to assess the impact of these interventions on the family system and its sustainability as well as to determine the extent to which the families' needs are met.

### Methods

To review theoretical assumptions and clarify the research questions, we are currently conducting a qualitative pre-study. This explorative study includes expert interviews with early intervention facilitators as well as a focus group with eight parents of children receiving the intervention. Based on the results, we design a quantitative longitudinal study and a qualitative case study to be conducted from 2024 to 2029. The quantitative study employs a panel design, with each family being assessed three times: at the start of the intervention (T1), at the end of the intervention (T2), and one year later (T3). At each time point, the parents evaluate their parenting sense of competence, their quality of life, the child's abilities and participation, and the family's network with other supporting institutions. In addition, we will survey the parents' expectations at T1, the fulfillment of expectations and effectiveness of the intervention at T2, and long-term effects at T3. The qualitative case study will compare five cases to reconstruct the course of family histories and individual family support. We estimate 200 families to participate in the evaluation each year between 2024 and 2028.

### Results

We anticipate the family system to undergo sustainable improvements including enhanced quality of life and sense of competence among parents, as well as increased levels of participation among children.

### Conclusions

With regard to other studies indicating the positive impact of early childhood interventions (4), this study aims to ascertain the value and sustainability of these interventions from the parents' perspective and to contribute to optimizing care for children at risk in Austria.

- 1. Shonkoff JP, Phillips DA. From neurons to neighborhoods: The science of early child development. Washington, D.C.: National Academy Press; 2000
- 2. Käsmann-Kellner B, Seitz B. Ausgewählte Aspekte der Kinderophthalmologie für Nichtkinderophthalmologen. Der Ophthalmologe. 2012 Jun;109(6):603–22. doi:10.1007/s00347-011-2495-6
- 3. Liljeberg H, Magdanz E, Reuse S. Eltern von Kindern mit Beeinträchtigungen Unterstützungsbedarfe und Hinweise auf Inklusionshürden. Vol. FB613, SSOAR. Berlin: Bundesministerium für Arbeit und Soziales; 2022
- 4. Resch B, Hasenbacher V, Kurz R. Die Bedeutung und Effizienz interdisziplinärer Frühförderung für Kinder mit Beeinträchtigung oder drohender Behinderung eine Literaturrecherche. Pädiatrie & Pädologie. 2020 Jul 13;55(4):186–96.

# First insights into SESAM: "SonoElastography: Sarcopenia Associated Muscle Change"

<u>Basel Habboub</u><sup>1</sup>, Guntje Kneiseler<sup>2</sup>, Robert Speer<sup>1</sup>, Marlene Bock<sup>1</sup>, Ursula Berger<sup>3,4</sup>, Katrin Singler<sup>1,5</sup>, Markus Gosch<sup>1</sup>

<sup>1</sup>Department of Geriatrics, University Hospital, Paracelsus Medical University Nuremberg, Prof. Ernst-Nathan-Str.1, 90419 Nuremberg, Germany; <sup>2</sup>Department of Gastroenterology, Hepatology and Endocrinology, University Hospital, Paracelsus Medical University Nuremberg, Prof. Ernst-Nathan-Str.1, 90419, Nuremberg, Germany; <sup>3</sup>Institute for Medical Information Processing, Biometry and Epidemiology, Faculty of Medicine, LMU Munich, Munich, Germany; <sup>4</sup>Pettenkofer School of Public Health, Munich, Germany; <sup>5</sup>Institute for Biomedicine of Ageing, Friedrich-Alexander-University, Kobergerstr. 60, 90408 Erlangen-Nuremberg, Germany; Contact: Basel-h92@hotmail.com

## Objective

Sarcopenia poses a growing global health concern and is associated with an increase in negative health outcomes. Shear wave elastography (SWE) and B-mode ultrasound are two promising techniques for detection of muscle mass loss.

### Methods

This diagnostic cross-sectional study with two control groups has 66 mobile participants assigned equally into three groups: sarcopenic older adults (65+), non-sarcopenic older adults (65+) and younger adults ( $\leq$ 30). The rectus femoris (RF) muscle of the dominant limb is examined using standardized protocols. The main outcome — RF muscle stiffness in resting position — is measured using SWE. Secondary outcomes include muscle stiffness, cross-sectional area and pennation angle. Muscle stiffness is compared in resting position between the three groups. Diagnostic performance of the outcomes and their correlation with appendicular skeletal muscle mass (ASM) measured using bioelectrical impedance analysis are examined.

### Results

The current sample (n = 34, ongoing inclusion) shows a RF stiffness in resting position in sarcopenic individuals of mean 24.7 kPa (SD 11.4) compared to 19.5 kPa (7.2) in non-sarcopenic older adults and 19.8 (5) among younger adults. The Spearman correlation coefficient between muscle stiffness in resting position and ASM is 0.35, compared to -0.17 in the passively stretched position. B-Mode measurements correlations with ASM are: muscle thickness 0.68, cross-sectional area 0.73 and pennation angle 0.34.

### Conclusions

SESAM preliminary results show slightly higher RF stiffness in sarcopenic adults, which seems an interesting feature in sarcopenia detection. Also, B-Mode measurements seem to correlate closely with muscle mass.

- Bastijns S, De Cock AM, Vandewoude M, Perkisas S. Usability and Pitfalls of Shear-Wave Elastography for Evaluation of Muscle Quality and Its Potential in Assessing Sarcopenia: A Review. Ultrasound in medicine & biology. Nov 2020;46(11):2891-2907. doi:10.1016/j.ultrasmedbio.2020.06.023
- Alfuraih AM, O'Connor P, Hensor E, Tan AL, Emery P, Wakefield RJ. The effect of unit, depth, and probe load on the reliability of muscle shear wave elastography: Variables affecting reliability of SWE. J Clin Ultrasound. 2018/02/01/ 2018;46(2):108–115. doi:10.1002/jcu.22534
- Alfuraih AM, Tan AL, O'Connor P, Emery P, Wakefield RJ. The effect of ageing on shear wave elastography muscle stiffness in adults. Aging Clinical and Experimental Research. 2019/12/01 2019;31(12):1755-1763. doi:10.1007/s40520-019-01139-0
- 4. Akagi R, Yamashita Y, Ueyasu Y. Age-Related Differences in Muscle Shear Moduli in the Lower Extremity. Ultrasound in medicine & biology. Nov 2015;41(11):2906-12. doi:10.1016/j.ultrasmedbio.2015.07.011
- Janczyk EM, Champigny N, Michel E, et al. Sonoelastography to Assess Muscular Stiffness Among Older Adults and its Use for the Diagnosis of Sarcopenia: A Systematic Review. Ultraschall Med. 2020/11/13/ 2020;doi:10.1055/a-1293-8057

# Parents Media Guide: How parental Smartphone use influences family life

Marion Hantinger<sup>1</sup>, Plattner, B.<sup>2</sup>, Priewasser, B.<sup>1</sup>, Lang, H.<sup>1</sup>, Brisch, Brisch<sup>1</sup>

<sup>1</sup>Institute for Early Life Care, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Division of Child and Adolescent Psychiatry, CDK, Salzburg, Austria; Contact: marion.hantinger@stud.pmu.ac.at

## Objective

Scientific studies report that the mere presence of a smartphone reduces the quality and enjoyment of the interaction between conversation partners, e.g., when a phone is used / is present during meal (1,2). It appears that the ability of adults to interact closely with their babies and young children is influenced by smartphone use. In addition, mothers who report that they use their smartphone more often, are less able to calm their child down after a stressful situation (3). Constant interruptions also influence the relationship between mother and child. For example, mothers - depending on their spontaneous smartphone use while eating - show less verbal and non-verbal contact with their children (4).

## Methods

The present empirical work investigates the influence of parental smartphone-use on the quality of family life e.g., infants' behavioral and physiological stress responses due to parental smartphone use and sums up several disadvantages of parental smartphone use next to their children. In the prospective follow-up interview study, parents are asked whether there has been a change in their behavior as a result of participation in the Smart Baby study. The aim of the work is to develop recommendations for parents and professionals working with parents or parents-to-be.

### Results

The results of the Smart Baby Study show a correlation between the interaction interruptions and the heart rate of the children, shown as a statistically significant increase in heart rate (HR) and thus a physiological stress response in the children. These findings will be discussed in the light of current knowledge based on further related empirical data. Together with the findings of the interviews, valuable statements can be made about appropriate smartphone use of parents.

### Conclusions

For creating the new parental smartphone guidebook "Parents Media Guide", we concentrate on the main influence factors of parental smartphone use, based on empirical data. Each chapter is devoted to a different topic, from the possible negative effects of smartphone use on family life to the testimonials and finally the summary of the findings. The results of the Smart Baby Study will be included in the evaluation.

- 1. Dwyer, R. J., Kushlev, K., & Dunn, E. W. (2018). Smartphone use undermines enjoyment of face-to- face social interactions. Journal of Experimental Social Psychology, 78, 233-239. doi:https://doi.org/10.1016/j.jesp.2017.10.007
- 2. Misra, S., Cheng, L., Genevie, J., & Yuan, M. (2016). The iPhone Effect. Environment and Behavior, 48(2), 275-298. doi:10.1177/0013916514539755.
- 3. Myruski, S., Gulyayeva, O., Birk, S., Pérez-Edgar, K., Buss, K. A., & Dennis-Tiwary, T. A. (2018). Digital disruption? Maternal mobile device use is related to infant social-emotional functioning. Developmental Science, 21(4), e12610.
- 4. Radesky, J., Miller, A. L., Rosenblum, K. L., Appugliese, D., Kaciroti, N., & Lumeng, J. C. (2015). Maternal mobile device use during a structured parent-child interaction task. Academic Pediatrics, 15(2), 238–244.

# Role of galanin receptor 2/3 in inflammation associated with peritonitis

Sara Huber<sup>1</sup>, Stefanie Gaisbauer<sup>1</sup>, Antonia Gantschnigg<sup>2</sup>, Christina Brandstötter<sup>1,3</sup>, <u>Theresa</u> <u>Haslauer<sup>1</sup></u>, Barbara Kofler<sup>1</sup>

<sup>1</sup>Research Program for Receptor Biochemistry and Tumor Metabolism, Department of Pediatrics, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Surgery, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Department of Biosciences and Medical Biology, Paris-Lodron University of Salzburg, Salzburg, Austria; Contact: th.haslauer@salk.at

## Objective

Peritonitis poses a serious medical challenge due to abdominal wall inflammation, often triggered by infectious agents. If not medicated timely, it can lead to life-threatening conditions. Therefore, understanding its immunological mechanisms is crucial for innovative therapies. The regulatory peptide galanin and its receptors (GAL1-3-R) are expressed on various immune cells and play pivotal roles in diverse immunoregulatory mechanisms and inflammation (1) upon mediation of the cross-communication between the nervous- and immune system. Thus, we aimed to explore the impact of GAL2/3-R knockout (KO) in a preclinical model of peritonitis.

### Methods

The impact of GAL2/3-R knockout (KO) was assessed in a preclinical peritonitis model, using aged 4% thioglycollate medium (TGM) injected intraperitoneal in GAL2/3-R KO and corresponding WT mice. Comparison to vehicle controls (0.9% saline) was included to discern treatment-specific effects. After 6 hours, peritoneal fluid, spleen and blood cells were collected. Phenotyping of myeloid cell populations was performed via flow cytometry. Chemokine levels in the peritoneum were assessed via multiplex immunoassays.

#### Results

Comparison of TGM-treated versus vehicle mice exhibited successful induction of acute peritonitis. Notably, in GAL2/3-R KO mice the induction of chemokines, especially RANTES (regulated on activation, normal T cell expressed and secreted), IP-10 (Interferon gamma-induced protein 10), and MIG (Monokine induced by Gamma-Interferon), upon TGM treatment was less prominent compared to WT mice. Simultaneously, increased levels of peritoneal macrophages were detected by flow cytometry. Immunophenotyping of blood and spleen revealed reduced neutrophil levels in both compartments upon induction of peritonitis in GAL2/3-R KO compared to WT mice, accompanied by elevated levels of dendritic cells in the blood.

### Conclusions

GAL2/3-R KO mice lacked the characteristic induction of neutrophils by the innate immune system but showed induction of cells involved in early initiation of the adaptive immune responses. The observed alterations in chemokine levels and immune cell populations specifically in GAL2/3-R KO mice suggest a regulatory role for GAL2/3-R in modulating the immune response in peritonitis. Consequently, antagonism of GAL2/3-R could potentially dampen the sometimes overshooting immune response observed in peritonitis, thus representing a promising avenue for therapeutic interventions.

### Acknowledgements

The project was supported by the Austrian Science Fund (P32403-B).

#### References

1. Koller A, Brunner SM, Bianchini R, Ramspacher A, Emberger M, Sternberg F, Schlager S, Kofler B. Galanin is a potent modulator of cytokine and chemokine expression in human macrophages. Sci Rep. 2019 May 10;9(1):7237. PMID: 31076613.

# Therapeutic Drug Monitoring in critically ill patients with urosepsis

Wolfgang Hitzl<sup>2,3,4</sup>, Rainer Höhl<sup>5</sup>, Sascha Pahernik<sup>6</sup>, Clemens Hüttenbrink<sup>6</sup>, Fritz Sörgel<sup>7</sup>, Jens Kubitz<sup>1</sup>

<sup>1</sup>Department of anesthesiology, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Research and Innovation Management (RIM), Biostatistics and Publication of Clinical Trial Studies, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Department of Ophthalmology and Optometry, Paracelsus Medical University, Salzburg, Austria; <sup>4</sup>Research Program Experimental Ophthalmology and Glaucoma Research, Paracelsus Medical University, Salzburg, Austria; <sup>5</sup>Institute of Clinical Hygiene, Medical Microbiology and Infectiology, Paracelsus Medical University, Nuremberg General Hospital, Germany; <sup>6</sup>Department of Urology, Paracelsus Medical University, Nuremberg General Hospital, Germany; <sup>7</sup>Institute for Biomedical and Pharmaceutical Research, Nuremberg-Heroldsberg, Germany; Contact: anne.kamphausen@klinikum-nuernberg.de

# Objective

Sepsis remains one of the leading causes of death worldwide which is defined as a dysregulated response to microbial infection. There is a strong recommendation for early and effective antimicrobial treatment. However, as antimicrobial treatment may "kill the germ but also the patient", a therapeutic drug monitoring (TDM) may be used to optimize and individualize antimicrobial therapy.

### Methods

A single-center retrospective analysis of patients with urosepsis and therapeutic drug monitoring was performed. The influence of different factors such as SOFA-Score, origin of urosepsis, pathogen in urine and blood samples, and the use of renal replacement on the target concentration of optimal and minimal trough levels was investigated. Using logistical regression, the impact of various parameters on the minimum and optimal plasma-level of antibiotics in TDM was analyzed.

### Results

Between 2017 and 2019, 222 patients were treated for urosepsis at our institution, of which 84 patients needed intensive care treatment. In 20 cases, a therapeutic drug monitoring was performed Plasma level of antimicrobial therapy in patients with urosepsis caused by E. coli and Proteus mirabilis, the most common pathogen in blood and urine culture was significantly more frequent in the target concentrations of the optimal trough level of antimicrobial treatment. In addition, there was a significant correlation between the SOFA-Score and target concentrations: the higher the score, the higher the probability that the concentrations were within the target range.

### Conclusions

TDM supports early and effective antimicrobial treatment in urosepsis. Patients with a low SOFA score and pathogens others than E.coli or Proteus mirabilis may profit from TDM with more plasma levels being within target range. A more pronounced organ dysfunction (high SOFA-Score) correlates positively with the achievement of the desired trough level. TDM should be considered in every patient with urosepsis.

- 1. Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). JAMA. 2016;315(8):801-810. doi:10.1001/jama.2016.0287
- 2. Phe K, Heil EL, Tam VH. Optimizing Pharmacokinetics-Pharmacodynamics of Antimicrobial Management in Patients with Sepsis: A Review. J Infect Dis. 2020;222(Suppl 2):S132-S141. doi:10.1093/infdis/jiaa118
- Brunkhorst FM, Weigand MA, Pletz M, et al. S3 Sepsis-Praevention-Diagnose-Therapie-Nachsorge. Published online February 2020. Accessed March 10, 2022. https://www.awmf.org/uploads/tx\_szleitlinien/079-001k\_S3\_Sepsis-Praevention-Diagnose-Therapie-Nachsorge\_2020-02.pdf
- 4. Roberts JA, Norris R, Paterson DL, Martin JH. Therapeutic drug monitoring of antimicrobials. Br J Clin Pharmacol. 2012;73(1):27-36. doi:10.1111/j.1365-2125.2011.04080.x

## First results from a multimodal day-care psychosomatic post-COVID treatment approach

<u>Hanna Hofmann</u><sup>1</sup>, Angela Fricke<sup>1</sup>, Paul Köbler<sup>1</sup>, Christine Hertle<sup>1</sup>, Eva Krauß-Köstler<sup>1</sup>, Markus Müller<sup>1</sup>, Barbara Stein<sup>1</sup>, Peter Radermacher<sup>2</sup>, Christiane Waller<sup>1</sup>

<sup>1</sup>Department of Psychosomatic Medicine and Psychotherapy, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Anesthesiological Pathophysiology and Process Engineering, University Hospital, Ulm, Germany; Contact: Hanna.Hofmann@klinikum-nuernberg.de

## Objective

The WHO defines post-COVID syndrom as a range of different symptoms that persist or develop at least three months after a COVID infection, persist for a minimum of two months and cannot be explained otherwise (1). Patients with post-COVID syndrome are in urgent need of an adequate treatment option (2). However, published studies with an interdisciplinary, multimodal approach for the therapy of post-COVID syndrome are still limited (3). We established a half-day multimodal treatment programme at the Department of Psychosomatic Medicine and Psychotherapy in a large acute care hospital for post-COVID patients. Women and men with post-COVID have been treated there and scientifically monitored to evaluate the effects of the psychosmatic treatment.

### Methods

This observational study between January 2022 and March 2023 conducted as documentation the Patient Health Questionnaire (PHQ-D consisting of PHQ-15 (somatization), PHQ-9 (depressive symptoms), PHQ-Stress and GAD (anxious symptoms)), the ICD-10 Symptom Rating (ISR) and specialized psychological measurement like the Fatigue Scale (FS) and the Health Status Questionnaire (SF-12 consisting of the Physical Composite Score (PCS) and the Mental Composite Score (MCS)) at admission and discharge. Therapy lasted 4 weeks and consisted of a multimodal psychosomatic treatment as well as neuropsychological training and individualised physical training.

### Results

A total of 65 patients, 48 women and 17 men, consented to post-COVID treatment after an appeal via the public media. At admission, PHQ and FS showed a high level of somatoform complaints (PHQ-15: M = 16.00, SD = 5.57) and fatigue symptoms (FS: M = 27.06, SD = 4.42) whereas depressive (PHQ-9: M = 13.98, SD = 5.27) and anxious symptoms (GAD: M = 9.55, SD = 5.62) were moderately pronounced. SF-12 revealed values below the 75th percentile (PCS: M = 29.16, SD = 6.65; MCS: M = 31.16, SD = 12.81). Ten patients ("dropout group") discontinued treatment after an average of one week. Symptom burden in these patients was sigificantly higher in all measured scales compared to the continuing treatment group (effect sizes ranging from d = 0.51 to d = 0.60). Compared to patients from our standard clinical settings, including patients with somatic comorbidity, post-COVID patients had a comparably high or even higher psychological symptom burden (e.g. PHQ-15: p < .001, d = 0.79; PHQ-9: p < .05, d = 0.39). Compared to admission, symptomatology at discharge was significantly improved (e.g. PHQ-15: p < .001, d = 0.43; GAD: p < .001, d = 0.38; FS: p < .01, d = 0.32; SF-12, MCS: p < .001, d = 0.39).

# Conclusions

Despite the short treatment time compared to standard psychosomatic therapy, all patients were able to significantly benefit from the treatment. Further data will show whether increasing the duration of treatment can further strengthen the positive effects of multimodal psychosomatic therapy.

- 1. Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. The Lancet Infectious Diseases. 2022 Apr;22(4):e102–7.
- Koczulla AR, Ankermann T, Behrends U, Berlit P, Berner R, Böing S, et al. AWMF S1-Leitlinie Long/ Post-COVID (Stand 17.08.22). 2022; Available from: https://register.awmf.org/assets/ guidelines/020-027I\_S1\_Post\_COVID\_Long\_COVID\_2022-08.pdf
- 3. Kupferschmitt A, Langheim E, Tüter H, Etzrodt F, Loew TH, Köllner V. First results from post-COVID inpatient rehabilitation. Front Rehabil Sci. 2023 Jan 23;3:1093871.

# The ubiquitin-proteasome system component Zbtb preferentially interacts with pathogenic variants of SLC26A4 (pendrin)

<u>Florian Huber</u><sup>1</sup>, Emanuele Bernardinelli<sup>1</sup>, Raffaella Liuni<sup>1</sup>, Houssein Nasser<sup>1</sup>, Martin Jakab<sup>2</sup>, Sebastian Roesch<sup>3</sup>, Silvia Dossena<sup>1,4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Center of Physiology, Pathophysiology and Biophysics, Institute of Physiology and Pathophysiology Salzburg, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Department of Otorhinolaryngology, Paracelsus Medical University, Salzburg, Austria; <sup>4</sup>FIZ - Regenerative Medicine & Novel Therapies (RM & NT); Contact: f.huber@pmu.ac.at

## Objective

The SLC26A4 gene codes for the electroneutral CI-/I-/HCO3- exchanger pendrin (PDS). The amino acid (AA) substitutions p.L236P and p.R409H are found in two of the most common pathogenic PDS protein variants leading to Pendred syndrome and non-syndromic autosomal recessive deafness B4 (DFNB4) within the Caucasian population (1). A yeast two-hybrid screening identified the zinc finger and BTB domain-containing protein (Zbtb), which is part of the ubiquitin-proteasome system (UPS) (2), as a potential PDS-interacting partner. Based on this, we want to elucidate the degradation pathway of PDS.

## Methods

Mice cochlea sections were immunostained with anti-PDS and anti-Zbtb antibodies and imaged by confocal microscopy. HEK cells were co-transfected with wild type (WT), L236P, or R409H PDS with or without Zbtb, or transfected with WT PDS or C-terminal truncated PDS variants. Cells were incubated with or without a proteasome inhibitor. Western blots or co-immunoprecipitation (Co-IP) experiments were done.

### Results

PDS and Zbtb co-localize within the stria vascularis of the mouse cochlea. There is a direct molecular interaction between Zbtb, WT PDS and PDS L236P & R409H. WT PDS and PDS L236P & R409H are ubiquitinated. Zbtb preferentially interacts and ubiquitinates PDS L236P & R409H compared to the WT. The truncated PDS variants 738X, 674X, and 584X show significantly decreased expression levels compared to the WT, while PDS 530X levels were not decreased.

# Conclusions

The co-localization of PDS and Zbtb in the inner ear, their direct molecular interaction, as well as the increased ubiquitination levels of PDS variants in the presence of Zbtb indicate that the UPS might be responsible for the degradation of WT PDS and the pathogenic PDS variants. The lack of degradation of the PDS 530X indicates that the interaction site between PDS and the UPS is located between the AA positions 530 and 584. Targeting this sequence might represent a new pharmacological strategy for Pendred syndrome/DFNB4.

### Acknowledgements

This research was funded by projects 2022-lif-004-DOSSENA and FIZ RM&NT Talent Pool SR 042023 to Silvia Dossena. The present address of S. Roesch is: Department of Otorhinolaryngology, University Hospital Regensburg, Regensburg, Germany.

- 1. Smith RJH, Iwasa Y, Schaefer AM. Pendred Syndrome / Nonsyndromic Enlarged Vestibular Aqueduct. In: Adam MP, Everman DB, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, et al., editors. GeneReviews(®). Seattle (WA): University of Washington, Seattle; 1993.
- 2. Furukawa M, He YJ, Borchers C, Xiong Y. Targeting of protein ubiquitination by BTB-Cullin 3-Roc1 ubiquitin ligases. Nat Cell Biol. 2003 Nov;5(11):1001-7. doi: 10.1038/ncb1056. Epub 2003 Oct 5. PMID: 14528312.

# Comparison of image quality in CT neck imaging between different protocols with linearly reduced radiation doses and contrast agent amounts

## **Christian Hummel**

Institute of Radiology and Nuclear Medicine, Paracelsus Medical University, Nuremberg, Germany; Contact: christian.hummel@klinikum-nuernberg.de

# Objective

One of the primary aim in CT imaging is to reduce radiation dose while maintaining diagnostic image quality, according to the ALARA principle (as low as reasonably achievable). Another aim, especially in elderly patients and in patients with impaired renal function is to reduce the amount of administered contrast agent/media (CA/CM). Therefore, continuous improvement is necessary especially in case of neck imaging, where organs sensitive to radiation are located such as the thyroid gland and salivatory gland.

## Methods

Patients randomly distributed to six groups consisting of at least 34 clients each group undergone a CT scan of the neck without preselection. Body weight adapted doses at 120 kV to a final lowest dose of 70 kV in steps of ten with corresponding diminished dose of contrast agent have been applicated. Dependent on the concentration (mg/ml) of the CA the amount was calculated in corresponding steps of 10 %. ROI-evaluation of different target organs was performed with measured correcting values of Hounsfield units at five different levels of surrounding air. Furthermore, a semiquantitive evaluation of the quality of obtained scans with correlated to a Likert scale was obtained. The corresponding parameters of each kV-level are correlated, and the MANOVA-analysis was the statistical method we used.

### Results

The hypothesis is that there is no difference in diagnostic value in varying amounts of CA at different kV-levels. Diagnostical value of the scans are evaluated with a semiquantitive Likert scale. The results show a decrease in the image quality with diminishing kV-level. In levels of 100, 90 and 80 kV there is almost no difference. The level with 70 kV differs slightly with a lower score of averaging around 2,5. Further results show differing variance in organs as the thyroid and the sternocleid muscle by almost no variance of HU in the other organs/structures. Last part indicates no lack of diagnostic information. Furthermore we reduce the cost of CA at almost halving it.

# Conclusions

Preliminary results show only slightly varying variance in kV-levels of 100, 90 and 80 with reduced image quality at 70 kV. In correlation with reduced consumption of CA at reasonably rates the aim to reduce amount the volume of it at reduced radiation doses are achieved. The diagnostic value by fostering radiosensitive organs according to the results of the Likerts scale show reduced imaging quality at low levels at 70 kV.

### References

1. Eller A, Wuest W, Kramer M, May M, Schmid A, Uder M, Lell MM. Carotid CTA: radiation exposure and image quality with the use of attenuation-based, automated kilovolt selection. AJNR Am J Neuroradiol. 2014 Feb;35(2):237-41. doi: 10.3174/ajnr.A3659. Epub 2013 Aug 1. PMID: 23907241; PMCID: PMC7965763.

# Within and between effects of mothers' and fathers' attachment, mentalizing and caregiving

# <u>Selina Ismair</u><sup>1,2</sup>, Antonia Dinzinger<sup>1,2</sup>, Gabriela Markova<sup>1,2,3</sup>, Jonas Schropp<sup>4</sup>, Karl Heinz Brisch<sup>1,5</sup>, Beate Priewasser<sup>1,2</sup>

<sup>1</sup>Institute for Early Life Care, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Pediatrics, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>University of Vienna, Austria; <sup>4</sup>Inselspital, Bern University Hospital, University of Bern, Switzerland; <sup>5</sup>Ludwig-Maximilian University Munich, Germany; Contact: selina.ismair@pmu.ac.at

# Objective

Parental sensitivity towards their children's needs is one of the most important factors influencing a child's social, emotional and cognitive development. Recent studies indicate that secure attachment representations (1) as well as the ability to reflect on one's own and others' mental states has a positive effect on sensitive caregiving in mothers (2) and fathers (3). However, it remains unclear how these traits and skills of one parent influence the other parents' skills and interaction patterns. Thus, the goal of the present study was to investigate the interplay between maternal and paternal attachment representations and reflective functioning (RF) on parental sensitivity, using a family system perspective.

# Methods

We observed N = 40 first-time families in a longitudinal, multi-method study. We used the Adult Attachment Interview to assess parental attachment and general RF during pregnancy, and the Parent Development Interview to measure mothers and fathers parental RF. When children were 6 months (Mage = 6.5 month), we observed interactions between each parent and their child and used the Emotional Availability Scales to assess parental sensitivity.

# Results

A structural equation model showed good fit (X2 = 15.34, p = .571, CFI = 1; RMSEA < 0.001). The model reveals individual associations between attachment representations and general RF in mothers ( $\beta$  = 0.68, p < .001) and fathers ( $\beta$  = 0.71, p < .001), mothers and fathers general RF ( $\beta$  = 0.46, p < .001), as well as a mediating effect of fathers parental RF on the association between mothers' parental RF and paternal sensitivity ( $\beta$  = 0.14; b [95% CI] = 0.12 [0.02; 0.26]). Fathers' parental RF does not appear to influence maternal sensitivity directly or indirectly (all p > .05).

# Conclusions

These results suggest that parents might influence each other in their reflective functioning and caregiving behavior, supporting family system theory. The findings align with recent evidence showing interdependencies between mothers and fathers (4). This study extends beyond previous research by focusing on attachment representations and reflective abilities as an important basis for sensitive interactions. This research contributes to a more holistic understanding of the modern family by investigating the effects of both mothers and fathers on each other and their children in a longitudinal study employing multiple non-self-report methods.

- 1. van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2019). Bridges across the intergenerational transmission of attachment gap. Current Opinion in Psychology, 25, 31–36
- Stacks, A. M., Muzik, M., Wong, K., Beeghly, M., Huth-Bocks, A., Irwin, J. L., & Rosenblum, K. L. (2014). Maternal reflective functioning among mothers with childhood maltreatment histories: links to sensitive parenting and infant attachment security. In Attachment & human development 16(5), 515–533.
- Dinzinger, A., Ismair, S., Brisch, K. H., Sperl, W., Deneault, A. A., Nolte, T., ... & Priewasser, B. (2023). Mentalizing in first-time fathers: reflective functioning as a mediator between attachment representation and sensitivity. Attachment & Human Development, 25(5), 544-565.
- 4. Gao, M. M., Du, H., Davies, P. T., & Cummings, E. M. (2019). Marital Conflict Behaviors and Parenting: Dyadic Links Over Time. Family Relations, 68(1), 135–149.

# Deckung komplexer pararektaler Beckendefekte: Rolle der freien myokutanen Musculus vastus lateralis Lappenplastik

Denis Ehrl<sup>1</sup>, Moritz Billner<sup>1</sup>, David Breidung<sup>1</sup>, Andre Barth<sup>1</sup>, Konrad Karcz<sup>1</sup>

<sup>1</sup>Department of Plastic, Reconstructive and Hand Surgery; Contact: konrad.karcz@klinikum-nuernberg.de

## Objective

Hintergrund: Aufgrund eines fortgeschrittenen Rektumkarzinoms kann eine Eviszeration mit Rektumamputation notwendig werden. Dabei können vor allem bei entsprechenden Voroperationen und neoadjuvanter Radiatio, ausgedehnte sowie tiefgreifende Weichteildefekte entstehen, die eine freie mikrovaskuläre Gewebetransplantation nötig werden lassen. In dieser Fallserie zeigen wir die erfolgreiche Kombination der muskulokutanen M. vastus lateralis-Lappenplastik (MVL) mit direktem Anschluss an die A. glutealis superior (1).

### Methods

Material und Methoden: Wir untersuchten über einen Zeitraum von 47 Monaten, 11 retrospektiv Fälle von Patienten, bei denen dorsale Beckendefekte nach Eviszeration und Rektumamputation bestanden, die mit lokalen oder regionalen Mitteln nicht zu verschließen waren. Bei all diesen Patienten erfolgte bei ausgedehnten Defekten mit tiefen pararektal Wundhöhlen eine Defektdeckung durch eine freie myokutaner MVL-Lappenplastik mit direkte Gefäßanschluss an die superioren Glutealgefäße.

### Results

Ergebnisse: Die Defektgröße betrug im Mittel 290,0 cm<sup>2</sup> (SD: 131,2; Range: 200 – 600 cm<sup>2</sup>). Die Defekttiefe betrug im Mittel 10,5 cm. Die Defekte machten somit MVL-Lappenplastiken mit einer mittleren Größe von 336,3 cm<sup>2</sup> notwendig. Es waren aufgrund von Nachblutungen drei operative Revisionen nötig. Es traten weder arterielle noch venöse Thrombosen sowie kein Lappenverlust auf. Einzig wurde eine Lappenspitzennekrose beobachtet, die sekundär per Direktnaht korrigiert werden konnte. Die Auswertung des Case-Mix ergab einen Wert von im Mittelwert 24,251 (SD: 21,699; Range: 7,036 – 65,748) Punkten, was die Komplexität der Fälle unterstreicht.

# Conclusions

Schlussfolgerungen: Unsere Ergebnisse zeigen, dass eine freie mirkovaskuläre MVL-Lappenplastik eine praktikable Therapieoption für pararektale Defekte darstellt, die nicht durch lokoregionale Methoden verschlossen werden können. Die A. glutea superior ist dafür ein sicheres und suffizientes Anschlussgefäß. In Kombination lassen sich auch ausgedehnte Defekte erfolgreich verschließen.

### References

1. Heppert, V., Kessler, T., Malze, K. et al. Vastus-lateralis-Lappen – Ein ideales Verfahren zur definitiven Sanierung von Infekthöhlen an der Hüfte. Unfallchirurg 103, 938–944 (2000). https://doi.org/10.1007/s001130050649

# Influencing factors on SARS-CoV-2-specific IgG antibody development following vaccination

Tanja Karl<sup>1,2</sup>, Anja Schuster<sup>1</sup>, Lea M. Stangassinger<sup>1</sup>, Gertie J. Oostingh<sup>1</sup>, Janne Cadamuro<sup>3</sup>

<sup>1</sup>Biomedical Sciences, Salzburg University of Applied Sciences, Puch/Salzburg, Austria; <sup>2</sup>Medical Sciences, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Department of Laboratory Medicine, Paracelsus Medical University, Salzburg, Austria; Contact: tanja.karl@fh-salzburg.ac.at

# Objective

Achieving immunization of the entire population by utilizing vaccine formulations targeting the same molecular structure was one of the goals during the severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) pandemic. This presupposes that a uniform vaccine will generate a comparable antibody response in all individuals, resulting in similar antibody levels. Influencing factors that affect our immune system are likely to have a substantial impact on the response towards an immunization. (1) Therefore, the presented study focused on the immune response following SARS-CoV-2 vaccination by considering various influencing factors, including the type of vaccine and its combinations, time, sex assigned at birth, body mass index (BMI), age, chronic diseases, and long-term treatment.

# Methods

A single venous blood sample (serum, 3 ml) was taken from subjects (aged 18 years and older, n = 1,106) from the county of Salzburg (Austria) between January and August 2022. All participants were vaccinated and had not knowingly acquired coronavirus disease 2019 (COVID-19). A questionnaire was distributed to the participants to assess their health state, the kind and date of vaccination(-s), anthropometric data, pre-existing illnesses, and chronic disorders. SARS-CoV-2 spike-protein-specific IgG was then detected utilizing the VITROS® Anti-SARS-CoV-2 IgG detection technique in an automated chemiluminescence immunoassay. The results are displayed in terms of binding antibody units (BAU) per ml. Statistical analysis was performed using the IBM SPSS Statistics (version 27) and GraphPad Prism (version 9.2.0) software.

### Results

Within this population study, the 3 times vaccinated subjects had a higher median antibody titer (835.5 BAU/ml) compared to the 2 times vaccinated participants (361.5 BAU/ml). Immunization with various vaccines results in significant variations in the titer for the 3 times vaccinated group (Spikevax 1,855 BAU/ml, Comirnaty 831.5 BAU/ml, Vaxzevria 304.5 BAU/ml). The highest antibody median value was obtained with 2 doses Comirnaty with 1 Spikevax dosage (2,115 BAU/ml) followed by 3 doses Spikevax and 1 dose Comirnaty with 2 doses Spikevax (1,780 BAU/ml). A strong negative connection was discovered between the amount of IgG antibodies and the weeks after the previous vaccination in the 3 times vaccinated group (r = -0.65; p-value  $\leq 0.0001$ ). Concerning the ongoing medication as an influencing factor, participants taking medication had a significantly lower median antibody titer (716 BAU/ml) than those who did not take any medication (921 BAU/ml). Sex assigned at birth (p-value 0.126), BMI (r: - 0.008; p-value 0.815), age (r: -0.052; p-value 0.11), and chronic illnesses (p-value 0.106) had no significant impact on the antibody titer.

# Conclusions

The results of this study showed, that the number, the type of vaccine and its combinations as well as the time to the last immunizing event and the intake of ongoing medication have an impact on the amount of SARS-CoV-2 specific IgG antibodies. This data is essential for future vaccine development, particularly in the context of personalized treatment.

### References

1. Zimmermann P, Curtis N. Factors That Influence the Immune Response to Vaccination. Clin Microbiol Rev. 2019 Mar 13;32(2):e00084-18. doi: 10.1128/CMR.00084-18. PMID: 30867162; PMCID: PMC6431125.

# Collection efficiency of mononuclear cells in offline extracorporeal photopheresis: can processing time be shortened?

<u>Orkan Kartal</u><sup>1</sup>, Nadja Lindlbauer<sup>1</sup>, Sandra Laner-Plamberger<sup>1</sup>, Eva Rohde<sup>1</sup>, Fabian Föttinger<sup>1</sup>, Laura Ombres<sup>1</sup>, Georg Zimmermann<sup>2,3</sup>, Cornelia Mrazek<sup>4</sup>, Wanda Lauth<sup>2,3</sup>, Christoph Grabmer<sup>1</sup>

<sup>1</sup>Department of Transfusion Medicine, Paracelsus Medical University Hospital Salzburg, Salzburg, Austria; <sup>2</sup>Team Biostatistics and Big Medical Data, IDA Lab, Paracelsus Medical University Salzburg, Salzburg, Austria; <sup>3</sup>Research and Innovation Management, Paracelsus Medical University Salzburg, Salzburg, Austria; <sup>4</sup>Department of Laboratory Medicine, Paracelsus Medical University Hospital Salzburg, Salzburg, Austria; Contact: o.kartal@salk.at

# Objective

Extracorporeal photopheresis (ECP) is a well-established but lengthy and burdensome cell-based therapy for various diseases such as cutaneous T-cell lymphoma, graft-versus-host disease and organ rejection after transplantation. The number of mononuclear cells (MNCs) that needs to be collected to obtain a clinical response to ECP is still under debate. The purpose of this retrospective study was to determine the number of lymphocytes, monocytes and neutrophils in mononuclear cell products (MCP) by flow cytometry and the collection efficiency in the offline ECP setting.

# Methods

We collected data from 10 different patients undergoing 162 ECP procedures using the Spectra Optia device for MNC collection. White blood cell (WBC) count of MCP was determined using a hematology analyzer. MNCs were analyzed for CD45 and CD14 expression by flow cytometry to exactly determine the collected lymphocyte and monocyte fractions.

# Results

Collected MCP showed high cell yields with 55.3 x 106/kg MNCs and 41.1 x 106/kg lymphocytes. MCP were characterized by high MNC (81.3 %) and low neutrophils (18.7 %) percentage. Mean collection efficiency for WBCs and for MNCs was 23.9 % and 62.0 %, respectively. The MNC fraction showed a moderate to high correlation between peripheral blood cell count of patients and MCP count.

# Conclusions

This study is one of a few reports showing the monocyte-to-lymphocyte relation in MCP for ECP determined by flow cytometry. In comparison to historical data from inline ECP, the offline ECP processing one total blood volume results in considerably higher cell yields. For this reason, and to reduce the burden on patients, we propose that the offline ECP processing time can be substantially reduced.

- 1. Piccirillo N, Putzulu R, Massini G, Di Giovanni A, Giammarco S, Metafuni E, et al. Inline and offline extracorporeal photopheresis: device performance, cell yields and clinical response. J Clin Apher 2021; 36: 118-126. doi: 10.1002/jca.21851., PMID: 33058243
- Worel N, Lehner E, Fuhrer H, Kalhs P, Rabitsch W, Mitterbauer M, et al. Extracorporeal photopheresis as second-line therapy for patients with acute graft-versus-host disease: does the number of cells treated matter? Transfusion 2018; 58: 1045-1053. doi: 10.1111/trf.14506., PMID: 29446444

# Characterization of patients with classical myeloproliferative neoplasms: A retrospective, single-center analysis.

<u>Dominik Kiem</u><sup>1</sup>, Lisa Pleyer<sup>1, 2</sup>, Kim Föhrenbach<sup>2</sup>, Michael Leisch<sup>1</sup>, Gabriel Rinnerthaler<sup>1</sup>, Alexander Egle<sup>1, 2</sup>, Richard Greil<sup>1, 2</sup>, Thomas Melchardt<sup>1, 2</sup>

<sup>1</sup>Department of Internal Medicine III with Hematology, Medical Oncology, Hemostaseology, Rheumatology and Infectiology, Oncologic Center, Salzburg Cancer Research Institute-Laboratory for Immunological and Molecular Cancer Research (SCRI-LIMCR), Paracelsus Medical University, 5020 Salzburg, Austria; <sup>2</sup>Austrian Group Medical Tumor Therapy, 5020 Salzburg, Austria; Contact: d.kiem@salk.at

Austrian Group Medical Tumor Therapy, 5020 Salzburg, Austria; Contact: d.

### Objective

Classic Philadelphia-chromosome negative myeloproliferative neoplasms (MPN) are a group of clonal malignant hematologic disorders. They include polycythemia vera (PV), essential thrombocythemia (ET) and myelofibrosis (MF). Patients with MPN have an increased risk for thromboembolic events. PV or ET can evolve to MF over time, and transformation into blast phase, i. e. acute leukemia, is possible. We aim to analyze the course of disease in all patients treated at our center over the last 20 years. Here, we provide a first characterization of the cohort, with the goal of further data collection within the Austrian Myeloid Registry of the Austrian Group Medical Tumor Therapy (AMR-AGMT).

# Methods

Data were collected retrospectively. We conducted a search in the clinical database of the Department of Internal Medicine III and collected data of all MPN patients diagnosed and/or treated at our center since 2003. Statistical analysis is descriptive. For this analysis, patients with unclassifiable MPN were excluded.

# Results

Overall, 882 patients were included into the database, of which 124 were excluded for unclassifiable MPN, thereby leaving 758 patients. The largest subgroups were PV (n=348) and ET (n=338), followed by MF (n=150). 34 patients with PV (9,7%) and 32 patients with ET (8,6%) transformed into MF, respectively. Therefore, 84 patients were classified as primary MF (56%). Overall, 12 patients (1,70%) transformed into blast crisis. A JAK2V617F mutation was detected in most patients with PV (97,1%). It was also the most common mutation in ET (66,1%) and MF (65,3%). Mutations in CALR and MPL were found in 21,8% and 6% of patients with ET, and in 22,7% and 4,4% with MF. Of the entire cohort, only 4,5% patients had no classic driver mutation. For patients with a diagnosis of PV or ET (without transformation), at least one thrombotic event was documented in 36,5% and 31,5% of patients, respectively, whereas 21% and 33% of patients with primary and secondary MF had a thrombotic event. A cytoreductive therapy with hydroxyurea was performed in 76% of patients with PV and in 61% with ET, whereas anagrelide was used in 10,9% of ET. Overall, interferon alpha was chosen for 109 patients, mainly PV (n=50) and ET (n=57), whereas ruxolitinib was given to 132 patients. Of the patients with PV, 15% received interferon alpha and 16% ruxolitinib. 49 patients with primary MF and 43 patients with secondary MF received ruxolitinib, meaning that 61,3% of patients with MF received the JAK2 inhibitor.

# Conclusions

In our cohort, PV and ET are the most prevalent, with fewer patients with MF. The distribution of driver mutations is comparable to other studies. Potentially disease-modifying treatment with interferon alpha and ruxolitinib was chosen in a significant proportion of patients. The cohort seems suitable for more in depth analysis within the AMR-AGMT.

### References

 Mahmud M, Vasireddy S, Gowin K, Amaraneni A. Myeloproliferative Neoplasms: Contemporary Review and Molecular Landscape. Int J Mol Sci. 2023 Dec 12;24(24):17383. doi: 10.3390/ijms242417383. PMID: 38139212; PMCID: PMC10744078.

# A personalised and customisable serious gaming system for children with movement disorders

### Lorenz Kiwull<sup>1</sup>

<sup>1</sup>Research Institute for Rehabilitation, Transition and Palliation, PMU Salzburg, Austria; Contact: lorenz.kiwull@pmu.ac.at

### Objective

Special body-controlled computer games known as "serious games" are very well suited to motivating children with neurological movement disorders to engage in therapeutic exercises. Between 2016 and 2018, I conducted a study with 20 patients to gain initial experience in the use of serious games and obtain patient feedback. The result of the study at that time was that although the exercises were very motivating, they were not challenging enough and did not focus enough on the patients' goals: 50% of the children did not work directly on their rehabilitation goals at that time1. Our aim was to develop a system based on this feedback that allows the body-controlled games to be tailored precisely to the therapists' requirements and the patients' goals.

### Methods

Initially, the therapy system was developed iteratively in close consultation with patients and therapists. To adapt the system to the patients, we developed a "Body Selector" that allows therapists to easily enter goals by clicking on body parts and selecting the desired functional results. In addition, we have developed small gamified assessments so that the system can adapt to the patient's conditions, for example to the individual range of motion, and thus remain challenging. For long-term motivation, we developed a virtual world in which the patient can move independently as an "avatar" to the next therapeutic exercise, collect coins and solve tasks. The system was then certified as a medical device and standardized, anonymous feedback from patients and therapists was obtained from digital questionnaires in 6 different partner clinics after over 1000 therapy sessions. Standardized questions were rated on a scale from 0 to 100.

#### Results

Patient feedback showed that over 70% patients very much enjoyed the therapeutic games and they were neither too easy nor too difficult for over 65% to train. The Serious Gaming System was mostly used by the therapists for mixed therapy goals (63%), otherwise for coordination (17%), range of motion (8%) and balance (6%). More than 85% of therapists rated the therapy as effective or very effective for the therapy goal of the respective session.

### Conclusions

The Serious Gaming System was mostly used by the therapists for mixed therapy goals (63%), otherwise for coordination (17%), range of motion (8%) and balance (6%). More than 85% of therapists rated the therapy as effective or very effective for the therapy goal of the respective session. The data suggest that the Serious Gaming System now provides the right level of challenge and effort and is largely effective for the therapy goals set by the therapist.

#### References

1. Kiwull et al; Assessment of Therapeutically Guided Serious Gaming from the Point of View of the Treated Children and Adolescents—A Pilot Study; Neuropediatrics 2018; 49(S 02): S1-S6; DOI: 10.1055/s-0038-1676025

### Relationship between Education and Steatotic liver disease

<u>Florian Koutny</u><sup>1,2</sup>, Vanessa Frey<sup>3,4</sup>, Christian Datz<sup>5</sup>, Sophie Gensluckner<sup>6</sup>, Andreas Maieron<sup>2</sup>, Andrea Mega<sup>7</sup>, Patrick Langthaler<sup>3,8,9</sup>, Maria Flamm<sup>10</sup>, Daniel Weghuber<sup>1</sup>, Bernhard Iglseder<sup>11</sup>, Eugen Trinka<sup>3,13</sup>, Bernhard Paulweber<sup>6</sup>, Elmar Aigner<sup>6</sup>, Bernhard Wernly<sup>5,10,14</sup>

<sup>1</sup>Department of Human Medicine, PhD Medical Science, Paracelsus Medical University, Strubergasse 21, 5020 Salzburg, Austria; <sup>2</sup>Department of Internal Medicine 2, Gastroenterology and Hepatology and Rheumatology, Karl Landsteiner University of Health Sciences, University Hospital of St. Pölten, St. Pölten, Austria; <sup>3</sup>Department of Neurology, Christian Doppler University Hospital, Paracelsus Medical University and Centre for Cognitive Neuroscience, Ignaz-Harrer-Straße 79, 5020, Salzburg Austria.; <sup>4</sup>Neuroscience Institute, Christian Doppler University Hospital, Paracelsus Medical University and Centre for Cognitive Neuroscience, Ignaz-Harrer-Straße 79 5020, Salzburg Austria.; <sup>5</sup>Department of Internal Medicine, General Hospital Oberndorf, Teaching Hospital of the Paracelsus Medical University, Paracelsusstraße 37, 5110 Salzburg, Austria; <sup>6</sup>First Department of Medicine, University Clinic Salzburg, Paracelsus Medical University Strubergassse 21, 5020 Salzburg, Austria.; 7Gastroenterology Department, Bolzano Regional Hospital, Lorenz-Böhler-Straße 5, 39100 Bolzano, Italy; <sup>8</sup>Department of Artificial Intelligence and Human Interfaces, Paris Lodron University Kapitelgasse 4/6, 5020 Salzburg, Austria.; 9Team Biostatistics and Big Medical Data, IDA Lab Salzburg, Paracelsus Medical University, Strubergasse 16. A-5020 Salzburg, Austria: <sup>10</sup>Institute of General Practice, Family Medicine and Preventive Medicine, Paracelsus Medical University of Salzburg, Strubergasse 21, 5020, Salzburg, Austria; <sup>11</sup>Department of Geriatric Medicine, Christian-Doppler-Klinik, Paracelsus Medical University, Ignaz-Harrer-Straße 79,5020 Salzburg Austria; <sup>12</sup>Center for Public Health and Healthcare Research Paracelsus Medical University of Salzburg, Strubergasse 21, 5020, Salzburg, Austria; <sup>13</sup>Department of Public Health, Health Services Research and Health Technology Assessment, UMIT - University for Health Sciences, Medical Informatics and Technology, Eduard-Wallnöfer- Zentrum 1, 6060 Hall in Tirol, Austria; <sup>14</sup>Department of Internal Medicine, Saint John of God Hospital, Teaching Hospital of the Paracelsus Medical Private University, Salzburg, Austria.; Contact: florian.koutny@stpoelten.lknoe.at

#### Objective

A lower level of education has been shown to be associated with a higher risk of developing various health conditions like myocardial infarct or stroke (1). This is due to unhealthy and sedentary lifestyle and limited access to healthcare (2). But, in case of non-alcoholic fatty liver disease (NAFLD) this association remains unclear. Therefore, the aim of this study was to prove if there is an independent relationship between NAFLD and educational level, defined by the International Standard Classification of Education (ISCED).

#### Methods

This cross-sectional study included 8,315 participants from the Paracelsus 10,000 study. Subjects were randomly selected from local population. The primary finding of the study was to address the etiological question of NAFLD in relation to education through relative risk analysis. The primary endpoint was NAFLD defined by the Fatty liver Index (FLI). A score  $\geq$  60 was considered elevated (1). Education level was defined by: (i) those with a low education level (less than a high-school education), (ii) those with an intermediate education level (from high school to less than university or college degree) and (iii) those with a high education level (university or college degree) (3). Statistical analyses, included descriptive statistics, logistic regression models and risk analysis. Outcomes were adjusted for the covariables of age and sex in model 1 and for income, employment, and marital status in model 2. Included were those with an assessment for FLI, ISCED score and supplied data regarding age, sex, income, employment, and marital status. Those with known viral hepatitis or excessive alcohol consumption (>40g in females and >60g in males) were excluded.

#### Conclusions

The studies findings indicate that a lower education level increases the risk of NAFLD independent of confounding factors. Therefore, these findings highlight the potential impact of educational attainment on NAFLD risk and emphasize the need for targeted interventions in vulnerable populations.

#### Acknowledgements

We would like to extend our heartfelt gratitude to the initiators, medical personnel, participants, and the dedicated team involved in the Paracelsus 10,000 Study. Their invaluable contributions and support have been instrumental in the success of this research. Thank you all for your unwavering commitment and effort.

- 1. Mantovani A, Scorletti E, Mosca A, Alisi A, Byrne CD, Targher G. Complications, morbidity and mortality of nonalcoholic fatty liver disease. Metabolism. 10 2020;111S:154170. doi:10.1016/j.metabol.2020.154170
- 2. Adler NE, Newman K. Socioeconomic disparities in health: pathways and policies. Health Aff (Millwood). 2002;21(2):60-76. doi:10.1377/hlthaff.21.2.60
- 3. Silke S. The International Standard Classification of Education 2011. Comparative Social Research. 30. 365-379. 10.1108/S0195-6310(2013)0000030017.

# Caring for a relative with dementia and the lived experiences of boundaries in caregiving – a qualitative study in a rural area of Salzburg

Simon Krutter<sup>1</sup>, Andreas Büscher<sup>2</sup>, Margitta B. Beil-Hildebrand<sup>1</sup>

<sup>1</sup>Institute of Nursing Science and Practice, Paracelsus Medical University, Salzburg; <sup>2</sup>University of Applied Sciences Osnabrück, Osnabrück; Contact: simon.krutter@pmu.ac.at

### Objective

Family caregivers of persons with dementia provide an important contribution to the care of communitydwelling older people, especially in rural areas. Besides positive aspects, caring for a relative with dementia is often a demanding process (1). If caregiver burden leads to feelings of caregiver overload, the boundaries of dementia caregiving are reached and the institutionalization phase of the relative with dementia sets in. Objective of the study: How do families caring for relatives with dementia perceive the boundaries in family caregiving and what are their lived experiences when making decision to institutionalize the relative with dementia?

### Methods

A qualitative approach was used to explore the life-world of family caregivers within a case study design and a phenomenological framework. For data collection 21 problem-centred interviews with family caregivers in Salzburg county were conducted. The interviews were analysed according to the hermeneutical sociology of knowledge approach.

### Results

The results illustrate specific life-world aspects of family caregiving that exemplify the boundaries in the course of caregiving. Core characteristics described are "a build-up of events in the lived experience", "dramatic events", or "cascading recruitment of formal services", whereby typical boundaries experienced by family caregivers are described.

# Conclusions

The results indicate that the experience of boundaries is not an isolated episode, but a core aspect in the course of caring for a relative with dementia. The crossing of boundaries and dangerous courses of caring for a relative with dementia could be prevented with a counselling concept that takes a lifeworld approach.

### Acknowledgements

Supported by the dissertation funding program "Discite!" of Paracelsus Medical University, Salzburg (project number: D-17/02/005-KRU).

### References

1. Chiao, C. Y., Wu, H. S. & Hsiao, C. Y. (2015). Caregiver burden for informal caregivers of patients with dementia: A systematic review. Int Nurs Rev, 62(3), 340-350. doi:10.1111/inr.12194

# Interatrial block improves prediction of new-onset atrial fibrillation after cardiac surgery

<u>Spela Leiler<sup>1,2</sup></u>, Andre Bauer<sup>3</sup>, Rok Bernik<sup>2</sup>, Patrick Sluet<sup>2</sup>, Valentin Guenzler<sup>2</sup>, Matthias Angerer, Theodor Fischlein<sup>1,2</sup>, Jurij Matija Kalisnik<sup>1,4,5</sup>

<sup>1</sup>Department of Cardiac Surgery, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Paracelsus Medical University, Nuremberg, Germany; <sup>3</sup>Department of Computer Sciences, University of Chicago, United States; <sup>4</sup>Department of Cardiothoracic and Vascular Surgery, University of Graz affiliated Clinic KABEG, Klagenfurt, Austria; <sup>5</sup>Medical Faculty, University of Ljubljana, Slovenia;

Contact: spela.leiler@gmail.com

# Objective

This study seeks to examine the relationship between interatrial conduction block and postoperative atrial fibrillation, a condition that can lead to acute cardiopulmonary instability and is linked to heart failure, stroke, and increased mortality after cardiac surgery (1, 2).

# Methods

Perioperative 12-channel electrocardiograms from 3405 patients undergoing myocardial revascularization, valve surgery, ascending aorta surgery, or combinations thereof, were considered. Clinical and electrographic parameters were compared between patients with and without atrial fibrillation, and significant variables were analysed using univariate and stepwise multivariate logistic regression. Statistical significance was defined as p<0.05.

# Results

Among 2108 analysed patients, 764(36.2%) developed atrial fibrillation. Preoperative interatrial block was a strong independent predictor (3.18[2.55,3.96],p<0.001 and improved atrial fibrillation prediction, enhancing the area under the ROC curve from 71.8% to 75.6%. Delong's test confirmed the model with interatrial block had superior discrimination performance (p=0.013). Other independent predictors of atrial fibrillation included advanced age (1.05[1.03,1.07],p<0.001), female gender (1.86[1.45,2.38],p<0.001), history of cardiogenic shock (1.44[0.99,2.09],p=0.057), reduced left ventricular ejection fraction <40% (1.57[1.06,2.33],p=0.024),  $\beta$ -blockers (1.17[0.95,1.46],p=0.145), CHA2DS2-VASc and EuroScore II (0.87[0.79,0.97],p=0.01 and 1.04[0.99,1.11],p=0.138), preexisting left bundle branch block (1.59[0.92,2.74],p=0.097), cardiopulmonary bypass time (1.00[1.00,1.00],p=0.049), bicaval cannulation (1.45[0.88,2.41],p=0.035), cardiac surgery-associated acute kidney injury (3.19[2.45,4.15],p<0.001), and postoperative atrioventricular block (1.20[0.96,1.51],p=0.105), particularly Mobitz I (6.73[1.98,31.51],p=0.005).

# Conclusions

Incorporating ECG-derived parameters, especially interatrial block, can enhance the predictive accuracy for postoperative atrial fibrillation. Additional research is needed to clarify the relationship between conduction abnormalities and postoperative atrial fibrillation, which will aid in the targeted delivery of prophylactic therapies and interventions for patients at high risk.

- 1. Gaudino M, Di Franco A, Rong LQ, Piccini J, Mack M.Postoperative atrial fibrillation: from mechanisms to treatment. Eur Heart J.2023;44(12):1020-1039.doi:10.1093/eurheartj/ehad019
- Tse G, Wong CW, Gong M, Wong WT, Bazoukis G, Wong SH, Li G, Wu WKK, Tse LA, Lampropoulos K, Xia Y, Liu T, Baranchuk A; International Health Informatics Study (IHIS) Network. Predictive value of inter-atrial block for new onset or recurrent atrial fibrillation: A systematic review and meta-analysis. Int J Cardiol. 2018 Jan 1;250:152-156. doi: 10.1016/j.ijcard.2017.09.176. Epub 2017 Oct 3. PMID: 29017777.

# Novel genetic variants lead to sensorineural hearing loss and Enlarged Vestibular Aqueduct (EVA)

<u>Raffaella Liuni</u><sup>1</sup>, Emanuele Bernardinelli<sup>1</sup>, Florian Huber<sup>1</sup>, Sebastian Roesch<sup>2</sup>, Giorgia Girotto<sup>3</sup>, Silvia Dossena<sup>1,4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Otolaryngology, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Institute for Maternal and Child Health-I.R.C.C.S. "Burlo Garofolo", University of Trieste, Trieste, Italy; <sup>4</sup>FIZ - Regenerative Medicine & Novel Therapies (RM & NT), Paracelsus Medical University, Salzburg, Austria; Contact: raffaella.liuni@pmu.ac.at

# Objective

Biallelic pathogenic variants in the SLC26A4 gene, coding for the anion exchanger pendrin, are responsible for Pendred syndrome and nonsyndromic recessive hearing loss DFNB4. Both these clinical entities are associated with an enlarged vestibular aqueduct (EVA), which is the most common malformation of the inner ear (1). We have recruited the first Austrian cohort of 33 patients with hearing loss and EVA to define the prevalence and type of pathogenic sequence alterations in SLC26A4 and discover novel EVA-associated genes.

### Methods

The coding region and intron-exon boundaries of known EVA genes were amplified by end-point PCR and Sanger sequenced. The possible presence of the Caucasian EVA (CEVA) haplotype was determined with the rhAmp® SNP Assays (IDT). Copy number variation (CNV) in the SLC26A4 and STRC genes was assessed using the TaqMan<sup>™</sup> Copy Number Assays and QuantStudio 3D Digital PCR. For patients who remain undiagnosed, whole exome sequencing (WES) was performed. The pathogenicity of novel SLC26A4 and TJP2 variants was evaluated by functional and molecular assays of the corresponding proteins.

#### Results

Biallelic pathogenic variants in SLC26A4 were detected in 5/33 patients (15%). Based on the positive perchlorate discharge test, one of these 5 patients (1/33, 3%) had Pendred syndrome. Monoallelic variants in SLC26A4 were detected in 5/33 patients (15%). Of these, two were benign based on functional and molecular tests. The CEVA haplotype was found in 6/33 (18%) of patients, 3 of whom carried monoallelic pathogenic SLC26A4 variants, 2 carried biallelic pathogenic SLC26A4 variants, and one carried a monoallelic non-pathogenic SLC26A4 variant. Furthermore, biallelic pathogenic variants in FOXI1 (1/33, 3%), monoallelic X-linked pathogenic variants in POU3F4 (2/33, 6%), and biallelic pathogenic variants in GJB2 (2/33, 6%) have been found. No CNV of SLC26A4 and STRC genes was found. WES of patients with bilateral EVA and negative for known causative genes (15/33, 45%) detected mutations in 6 EVA-unrelated genes (SCD5, REST, EDNRB, TJP2, TMC1, and CDH23) in 5/33 patients (15%). Cell-based assays showed that the novel gene variant in TJP2 leads to an aberrantly localized protein product, supporting its pathogenicity.

### Conclusions

Sequence alterations in SLC26A4 and/or the CEVA haplotype, FOXI1, POU3F4, and GJB2 genes are responsible for hearing loss and EVA in 42% (14/33) of patients of this cohort. WES led to the identification of 6 genes previously not associated with EVA and allowed for the diagnosis of an additional 15% (5/33) of patients. The genetic causes remain unidentified in 42% (14/33) of patients. Functional and molecular studies are needed to assign or exclude the pathogenicity of novel variants and establish a causal link with disease.

### Acknowledgements

The present address of S Roesch is: ENT, University Hospital Regensburg, Germany.

### References

1. Roesch S, Rasp G, Sarikas A, Dossena S. Genetic Determinants of Non-Syndromic Enlarged Vestibular Aqueduct: A Review. Audiology research. 2021;11(3):423-42. PMID: 34562878

# Oxidative stress-related cellular aging causes dysfunction of the Kv3.1/KCNC1 channel reverted by melatonin

<u>Raffaella Liuni</u><sup>1</sup>, Sara Spinelli<sup>2</sup>, Alessia Remigante<sup>2</sup>, Gianluca Mantegna<sup>2</sup>, Giuseppe Legname<sup>3</sup>, Angela Marino<sup>2</sup>, Rossana Morabito<sup>2</sup>, Silvia Dossena<sup>1,4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina, Italy; <sup>3</sup>Laboratory of Prion Biology, Department of Neurosciences, Scuola Internazionale Superiore di Studi Avanzati (SISSA), Trieste, Italy; <sup>4</sup>FIZ -Regenerative Medicine & Novel Therapies (RM & NT), Paracelsus Medical University, Salzburg, Austria; Contact: raffaella.liuni@pmu.ac.at

## Objective

Kv3.1/KCNC1 is a voltage-gated channel that is abundantly expressed in fast-spiking principal neurons and GABAergic inhibitory interneurons throughout the ascending auditory pathway and in various brain regions. Inactivating mutations in the KCNC1 gene lead to various forms of epilepsy, and Kv3.1 expression within the auditory pathway declines with age, which might contribute to age-related hearing loss (1). As oxidative stress plays a fundamental role in the pathogenesis of both epilepsy and age-related hearing loss, we hypothesized that an oxidative insult might affect the function of this channel.

## Methods

To obtain a cellular model of oxidative stress-related aging, HEK 293 Phoenix cells were exposed to 100 mM D-galactose (D-Gal) for 3-72 hours and melatonin (100 uM for 48 hours) was used as an antioxidant. As a control, 100 mM D-Mannitol (D-Man) was used to obtain equal osmolarity of the cell culture medium. Total intracellular ROS levels were evaluated with 2',7'-dichlorofluorescein diacetate. Potassium currents were measured by patch-clamp in whole-cell configuration. Total RNA was extracted from cultured cells and Kv/KCN transcripts were quantified by RT-qPCR. Western blot analysis was performed to test the abundance of Kv3.1a protein at the plasma membrane, calreticulin, GAPDH, tubulin, phosphorylated Src, and phosphorylated Cav-1.

### **Results**

In this model of oxidative stress-related aging, intracellular ROS were dysregulated and the current of Kv3.1 was significantly reduced. The Kv3.1 reduction of function was not determined by a reduction in its transcript levels. Instead, exposure of cells to D-Gal resulted in a reduction of Kv3.1 protein abundance in the plasma membrane-enriched fraction. Melatonin reverted all these effects. The increased Src and Cav-1 phosphorylation levels, as well as metabolic reprogramming and endoplasmic reticulum stress denoted by elevated GAPDH and reduced calreticulin expression respectively, might have all contributed to a decreased Kv3.1 trafficking to the plasma membrane.

# Conclusions

These data identify Kv3.1 as a novel target of oxidative stress and suggest that Kv3.1 dysfunction caused by oxidative stress might contribute to age-related hearing loss and increased prevalence of epilepsy during aging. The pharmacological use of the antioxidant melatonin might be protective in these settings.

### Acknowledgements

The research was funded by projects 2022-lif-004-DOSSENA and FIZ RM&NT Talent Pool SR 042023 to Silvia Dossena.

# References

1. Peixoto Pinheiro B, Vona B, Löwenheim H, Rüttiger L, Knipper M, Adel Y. Age-related hearing loss pertaining to potassium ion channels in the cochlea and auditory pathway. Pflugers Archiv : European journal of physiology. 2021;473(5):823-40. PMID: 33336302

# Fructose appears to be able to influence the immune system

# Sebastian Lobmann<sup>1,2</sup>, Boris Musset<sup>1</sup>

<sup>1</sup>Center of Physiology, Pathophysiology and Biophysics, Klinikum Nuremberg Medical School, Paracelsus Medical University Nuremberg, 90419 Nuremberg, Germany;

<sup>2</sup>Department of gastroenterology, hepatology and endocrinology, General Hospital Nuremberg, Paracelsus Medical Private University Nuremberg, 90419 Nuremberg, Germany;

Contact: Sebastian.Lobmann@klinikum-nuernberg.de

## Objective

The respiratory burst of phagocytes is an important part of the innate immune defense. Oxygen radicals are formed to eliminate pathogens. At the Institute<sup>1</sup> we investigate the influence of different hexoses in different concentrations on the respiratory burst of human phagocytes.

## Methods

The desired cell fraction was isolated directly at the institute from fresh whole blood from volunteer, clarified donors. This was carried out in hexose-free solutions. After the isolation, the cells were transferred to solutions with the appropriate concentrations of hexoses. We trigger the respiratory burst using PMA. We determine the activity of this respiratory burst by detecting  $H_2O_2$  using the Amplex Red method (1). Two to three different hexoses were examined simultaneously in one essay. Glucose was set as control to establish a reference and comparability with other measurements.

### Results

The different hexoses have very individual characteristic changes on the  $H_2O_2$  production during the respiratory burst. As an example, eosinophils with fructose and glucose are presented here. Eosinophils have a reduced maximum  $H_2O_2$  production with fructose. In the first 25 min of the measurements there were no differences in rate of production at different D - fructose concentrations. Later (>25 min) results clearly deviating at different concentrations with fructose. Our previous evaluation by determining Vmax is insufficient here as the differences at the end could not be represented, therefore the first derivative was determined additionally at the time t=90 min.

# Conclusions

It appears that eosinophils metabolize fructose with a slight delay. However, this does not occur in the first 25 min, when the highest production rate of  $H_2O_2$  is found. It can be observed that an initial plateau begins at all fructose concentrations between 20 and 25 min before the increase continues at the high fructose concentrations. The storage capacity for glycogen of the cells can be calculated from the values measured in the measurements without hexose.

### References

1. Karakuzu O, Cruz MR, Liu Y, Garsin DA. Amplex Red Assay for Measuring Hydrogen Peroxide Production from Caenorhabditis elegans. Bio Protoc. 2019 Nov 5;9(21):e3409.

# Influence of the isolation method on the measurement of the respiratory burst

## Sebastian Lobmann<sup>1,2</sup>, Boris Musset<sup>1</sup>

<sup>1</sup>Center of Physiology, Pathophysiology and Biophysics, Klinikum Nuremberg Medical School, Paracelsus Medical University Nuremberg, 90419 Nuremberg, Germany;

<sup>2</sup>Department of gastroenterology, hepatology and endocrinology, General Hospital Nuremberg, Paracelsus Medical Private University Nuremberg, 90419 Nuremberg, Germany;

Contact: Sebastian.Lobmann@klinikum-nuernberg.de

### Objective

At the Institute1 we are investigating various factors influencing the respiratory burst of human phagocytes. We currently have two methods at our disposal for isolating the desired cell fraction. A ready-to-use isolation kit (MACSxpress® (1) to extract the target cell fraction from the unprocessed whole blood with antibodies in a magnetic field and a method with density gradient centrifugation and subsequent separation of the cell fractions using antibodies in a magnetized column.

### Methods

We trigger the respiratory burst in the desired cell fraction using PMA. We determine the activity of the respiratory burst by detecting  $H_2O_2$  using the Amplex Red method (2) in a fluorescence essay. The two methods were compared by repeating the identical experiments in which only the method of isolation was varied.

### Results

For neutrophils, there were no differences between the two methods, outside the variance. There was a clear difference between the two methods for eosinophils. Even by changing the manufacturer's instructions, only a slight increase in  $H_2O_2$  production was achieved, but no comparable results were obtained.

### Conclusions

Both methods appear to provide equivalent results for the isolation of neutrophils. However, it should be noted that in density gradient isolation, the neutrophils are loaded with antibodies. With the MACSxpress® method, the neutrophils should be isolated untouched according to the manufacturer. Despite adaptation of the MACSxpress® instructions, there are still clear differences in the eosinophils. Whether this difference is due to other buffers / antibodies in the MACSxpress® kit or to the eosinophils themselves has not yet been determined, as the manufacturer does not wish to disclose the components of the isolation kit. The measurements by Patch-clamp support the measurement results we obtain with the density gradient isolation.

- 1. https://www.miltenyibiotec.com/DE-en/products/macsxpress-separator-and-starting-kits.html#130-101-319
- 2. Karakuzu O, Cruz MR, Liu Y, Garsin DA. Amplex Red Assay for Measuring Hydrogen Peroxide Production from Caenorhabditis elegans. Bio Protoc. 2019 Nov 5;9(21):e3409.

# Telemedicine for the management of behavioral disorders in nursing home residents with dementia

Francesca Lubian<sup>1, 2</sup>, Albert March<sup>1</sup>, Nicola Vanacore<sup>3</sup>, Markus Gosch<sup>2</sup>

<sup>1</sup>South Tyrol Health Authority, Bolzano Department of Geriatrics, Italy; <sup>2</sup>Department of Geriatrics, Paracelsus Medical University, Nuremberg, Germany; <sup>3</sup>National Institute of Health, Rome, Italy; Contact: francesca.lubian@stud.pmu.ac.at

## Objective

The management of behavioral disorders in Nursing Homes (NHs) residents with dementia (BPSD – Behavioral and Psychological Symptoms of Dementia) is particularly difficult and often leads to unnecessary hospitalizations and incorrect or excessive use of physical and pharmacological restraint (1). Telemedicine can become the means to deliver quality care in individuals with BPDS in NHs, as other studies have shown (2, 3). However, an evaluation of specific health outcomes is needed. The aim of the study is to verify the effectiveness of teleconsultations for the management of BPSD in residents with dementia and to assess NH staff's satisfaction with the new method.

## Methods

This prospective cohort study involves monthly teleconsultations between the Geriatrician of the Memory Clinic of Bolzano and the General Practitioners (GPs) and Head Nurses (HNs) in charge of two NH in the Province of Bolzano: one where the Geriatrician visited in person monthly before the pandemic (Egna/Montagna) and one where consultations had never been conducted before (Renon). Two trainings were prepared for the staff involved, one in the technical area for the use of the POHEMA telemedicine platform, and one in the clinical area for the recognition of BPSD and delirium. At each teleconsultation, the characteristics of residents affected by dementia and BPSD were noted, and three outcome indicators were registered: the number of referrals to emergency rooms (ERs) or memory clinics for problems related to behavioral disorders, the number of physical restraints, and the number of therapeutic adjustments, according to STOPP/START criteria. The data were compared with data for the year 2019 from the NH software. Every 6 months NH staff completed an ad-hoc composed satisfaction questionnaire. For statistical comparisons, the Fisher exact test was used.

# Conclusions

One year of frequent and constant telemedicine contacts between the Bolzano Memory Clinic and the two NHs have helped to avoid sending any patients to ERs or Memory Clinics for reasons related to behavioral disorders. The data is comparable to that obtained when the Geriatrician traveled to the facility in person. In contrast, the use of physical restraints was not shown to be related to telemedicine (higher number of restrained subjects in 2023 than in 2019). The findings could be explained by an increase in physical restraint during the Covid-19 pandemic, as supported by literature. The number of therapeutic adjustments demonstrates that the presence of the Geriatrician (in person or remotely) contributes to better management of residents with dementia. The results of the operator satisfaction questionnaire encourages the use of teleconsultations in other NHs and memory clinics in the area.

- 1. Valk-Draad MP, Bohenet-Joschko S. Nursing-Homes sensitive hospitalizations and the relevance of telemedicine: a scoping review. Int J Environ Res Public Health 2022;19:12944
- Nkodo JA, Gana W, Debacq C, Aidoud A, Poupin P, Camus V, Fougere B. The role of telemedicine in the management of the behavoiural and psychological symptoms of dementia: a systematic review. Am J Geriatr Psychiatry 2022;30:1135-50
- 3. Groom LL, McCarthy M, Stimpfel AW, Brody AA. Telemedicine and telehealth in Nursing Homes; an integrative review. J Am Med Dir Assoc 2021;22:1784-1801
## Impact of sugars on reactive oxygen species production during the respiratory burst of phagocytes

<u>Iryna Mahorivska</u><sup>1</sup>, Eva Schrader<sup>1</sup>, Gustavo Chaves<sup>1</sup>, Stefan Stein<sup>2</sup>, Matthias Schweizer<sup>3</sup>, Boris Musset<sup>1</sup>

<sup>1</sup>Institute of Physiology, Pathophysiology and Biophysics, PMU, Nürnberg, Germany; <sup>2</sup>Flow Cytometry Unit, Institute for Tumor Biology and Experimental Therapy, Georg-Speyer-Haus, Frankfurt, Germany; <sup>3</sup>Federal Institute for Vaccines and Biomedicines, Paul-Ehrlich-Institut, Langen, Germany; Contact: Iryna.Mahorivska@klinikum-nuernberg.de

#### Objective

Reactive oxygen species (ROS) are produced by NADPH oxidase during the respiratory burst of phagocytes (1). This is crucial for defending against bacteria as well as for signal pathways. However, high production of ROS causes oxidative stress and tissue damage. ROS generation is substrate dependent and sugars are preferably used. However, little is known about the impact of common monosaccharides on ROS production.

#### Methods

We compared the impact of glucose, fructose, galactose and 2-deoxyglucose on H2O2 production using the fluorometric amplex red method in three different phagocytic cells: a genetically modified leukemia cell line PLB-985 X-CGD SgpsIBW (BW), neutrophils and monocytes, freshly isolated from human peripheral venous blood. Furthermore, we investigated the gene expression of glucose transporters using real-time PCR.

#### Results

H2O2 production by leukemia cell line BW, neutrophils and monocytes after incubation with different concentrations of several sugar substrates: glucose, fructose, galactose and 2-deoxyglucose was investigated. Glucose is a substrate for NADPH oxidase and triggers ROS generation. The minimal concentration of external glucose effecting H2O2 production in BW cells and neutrophils is 100µM and the maximal – 3mM. For monocytes the minimal concentration of glucose is below 10µM and maximal above 30mM. The difference in the concentrations of glucose effecting H2O2 production inbetween cells suggest difference in glucose transporting. Fructose and galactose are not a substrate for H2O2 production. 2-deoxyglucose competes with glucose and inhibits glucose-6-phosphate dehydrogenase in neutrophils, therefore H2O2 production decreases. For BW cells and monocytes 2-deoxyglucose without presence of external glucose has no effect. Values are means  $\pm$  SD. n = 3Differentiated and not differentiated BW cells were harvested on the stated day with subsequent RNA isolation and gene expression of GLUT1 was determined using real-time PCR, GAPDH was used as housekeeping gene. n = 2, mean  $\pm$  SD. No change in GLUT1 mRNA expression was seen on day 1 and 6 during DMSO-induced differentiation compared to undifferentiated cells (d0). Values are mean  $\pm$  SD, n = 2.

#### Conclusions

Our results show that sugars play a substantial role in ROS generation in phagocytes and thus might affect oxidative stress. ROS production seems to be highly dependent on glucose but not on other monosaccharides. Glucose is transported via GLUT1 transporter into BW cells. Gene expression of GLUT1 does not change during differentiation process of BW cells into phagocyte-like cells.

#### Acknowledgements

Supported by the Kerscher'sche Stfitung

#### References

1. Nguyen GT, Green ER, Mecsas J. Neutrophils to the ROScue: Mechanisms of NADPH Oxidase Activation and Bacterial Resistance. Front Cell Infect Microbiol; 2017, PMID: 28890882

#### Embroidered silk fibroin scaffolds for ACL tissue engineering

#### Yasir Majeed<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cell Biology;

<sup>2</sup>Workgroup BioEngineering, Institute of Polymers Materials, Leibniz-Institut für Polymerforschung Dresden e.V., Germany; Contact: yasir.majeed@pmu.ac.at

#### Objective

Anterior cruciate ligament (ACL) rupture requires reconstruction due to poor healing capacity of the ACL. The resulting joint instability increases the risk for the onset of osteoarthritis (OA). Silk has been used since centuries for medical applications due to high cytocompatibility and biomechanical strength. Bombyx (B.) mori (silk worm) derived silk consists of two components, fibroin and sericin. The latter is usually removed to reduce immunogenicity and improve biocompatibility as well as processability. Embroidering allows adaption of biomechanical properties by selecting biomimetic embroidery patterns. This project aims to identify an embroidered silk fibroin-based scaffold variant facilitating ACL tissue engineering and ligamentogenesis.

#### Methods

For removal of sericin, two different cleaning procedures were applied to the silk threads: a standard ("raw silk") and a novel ("purified silk") technique patented (DE102021118652A1) by the start-up company providing the threads. Silk fibroin threads of both cleaning degrees were either used to embroider pure silk scaffolds or were combined with PLA-co-caprolactone (P(LA-CL)) fibers (7:0). A previously established and approved zig zag embroidery pattern (1.8 stitch length, 15° stitch angle, 0.2 duplication shift) was conducted with a lower and upper thread and three embroidered plies were stacked together by means of a silk locking thread. Scaffolds had the dimension of the rabbit ACL (10 x 4 x 2-3 mm) and were disinfected with 70% ethanol. Hence, four groups (1: "raw silk", 2: "raw silk/P(LA-CL)", 3: "purified silk", 4: "purified silk/P(LA-CL)") were seeded with lapine cruciate ligament fibroblasts (LCL-Fb, 1.5-2 mio cells per scaffold, dynamical cultivation). Cytotoxicity assays were conducted with extracts (48 hours extraction in growth medium) of each scaffold variant using MTS assay and LCL and L929 fibroblasts (Fbs). Cell adherence and vitality was monitored with fluorescein diacetate/propidiumiodide and immunostaining of cytoskeletal and extracellular matrix proteins after 7 and 14 days.

#### Results

The fibroin single fiber diameters of un/treated silk was nearly similar. LCL-Fb adherence on scaffolds differed, being low on the scaffolds made of conventionally treated silk, higher on raw silk/P(LA-CL). Highest cell adherence was found on the purified silk scaffolds and purified silk/P(LA-CL). Interestingly, the P(LA-CL) integration into the scaffold increased also the cytocompatibility of raw silk fibers. Cell survival on the scaffolds was high for the whole observation period (14 days). The focal adhesion component paxillin was abundantly expressed in cells on purified silk scaffolds underlining intimate cell material interaction and also typical ECM could be detected.

#### Conclusions

Purified silk fibroin threads are highly cytocompatible and represent a versatile material, suitable to prepare embroidered LCL-Fb carriers without the necessity of further functionalization.

#### References

1. Hahn J, Gögele C, Schulze-Tanzil G. Could an Anterior Cruciate Ligament Be Tissue-Engineered from Silk? Cells. 2023,12(19):2350. PMID: 37830564

### Transcriptomic analysis unveils the regulatory network of Sturge-Weber syndrome-associated $G\alpha q$ protein during neural differentiation

Rakesh Mani<sup>1</sup>, Anna Maria Ernst<sup>1</sup>, Mirjana Malnar<sup>1</sup>, Christine Deisl<sup>1</sup>, Jan Pruszak<sup>1</sup>

<sup>1</sup>Institute of Anatomy and Cell Biology, Paracelsus Medical University, 5020 Salzburg, Austria.; Contact: rakesh.mani@pmu.ac.at

#### Objective

Sturge-Weber syndrome (SWS) is a rare neurovascular disorder attributed to a spontaneous mutation in the GNAQ gene, responsible for encoding the guanine nucleotide-binding protein G(q) subunit alpha ( $G\alpha q$ ). This syndrome presents distinct clinical features, including a facial port-wine birthmark (PWB), glaucoma, and various neurological symptoms such as intellectual disability and seizures. Despite the prevalence of neurological manifestations, there is a significant knowledge gap regarding the molecular intricacies underlying SWS within neural cells. This study aims to uncover the molecular mechanisms regulated by  $G\alpha q$  in neural cells.

#### Methods

Pluripotent stem cells (PSC) were subjected to neuronal differentiation using a dual SMAD inhibition protocol established in our lab. Gαq knockouts were generated using the CRISPR-mediated non-homologous endjoining method. Knockout samples were screened via western blot analysis, and validation was carried out using Sanger sequencing. Transcriptional differences between control and Gαq knockout neural stem cells (NSC) were assessed through RNA sequencing. Gene ontology analysis was performed using the online tool PANTHER, while pathway enrichment analysis was carried out using Integrated Molecular Pathway Level Analysis (IMPaLA).

#### Results

Confirmation of neuronal differentiation using wild-type PSC was shown by elevated levels of neural markers NESTIN and MAP2. The expression patterns of the Gaq gene showed an increase during the proliferative stage of neural stem cells, indicating its potential role in neural proliferation. RNA sequencing analysis comparing knockout and control cells revealed increased differential expression of genes in Gaq knockout neural stem cells compared to Gaq knockout embryonic stem cells, aligning with previous qPCR results. Fifteen genes, predominantly non-coding RNA, were identified as commonly differentially expressed between embryonic stem cells and neural stem cells showed differential regulation of 94 genes in Gaq knockout NSC. Pathway analysis using the IMPaLA tool highlighted neural differentiation as the top pathway represented by 8 of the 94 genes, with a very high p-value. Notably, Gaq knockout in NSC modulated ASCL1, a master regulator of neurogenesis.

#### Conclusions

Sturge-Weber syndrome associated Gaq protein modulates genes related to neural differentiation, potentially contributing to the neurological symptoms observed in this condition.

#### Acknowledgements

We greatly acknowledge Sturge-Weber foundation for the grant to initiate this project. We thank Leonie Enzinger, Shamsudeen Suleiman, Chiara Thommes, and other members of the Institute of Anatomy and Cell Biology, PMU for their technical support to the project.

- 1. Uhlén M et al,. Tissue-based map of the human proteome. Science. 2015; 347(6220).
- Shirley MD, Tang H, Gallione CJ, Baugher JD, Frelin LP, Cohen B, North PE, Marchuk DA, Comi AM, Pevsner J. Sturge-Weber syndrome and port-wine stains caused by somatic mutation in GNAQ. N Engl J Med. 2013;368(21):1971-1979.

#### EZH1 - a new therapeutic target for biliary tract cancer?

<u>Christian Mayr</u><sup>1,2</sup>, Daniela Mayr<sup>3</sup>, Petra Huber-Cantonati<sup>4</sup>, Anna Wipplinger<sup>1</sup>, Markus Ritter<sup>1</sup>, Daniela Neureiter<sup>5</sup>, Tobias Kiesslich<sup>1,2</sup>

<sup>1</sup>Center for Physiology, Pathophysiology and Biophysics, Institute of Physiology and Pathophysiology, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Internal Medicine I, University Clinics Salzburg, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Research Program for Experimental Ophthalmology, Department of Ophthalmology and Optometry, University Hospital of the Paracelsus Medical University, Salzburg, Austria; <sup>4</sup>Institute of Pharmacy, Department of Pharmaceutical Biology and Clinical Pharmacy, Paracelsus Medical University, Salzburg, Austria; <sup>5</sup>Institute of Pathology, University Clinics Salzburg, Paracelsus Medical University, Salzburg, Austria; Contact: christian.mayr@pmu.ac.at

#### Objective

Biliary tract cancer (BTC) is a tumor with limited therapeutic options and dismal outcome. Hence, the identification and investigation of new therapeutic approaches is of utmost importance. The polycomb repressive complex 2 (PRC2) is a multi-protein complex that regulates gene expression via trimethylation of lysine 27 at histone 3 (H3K27me3). The canonical PRC2 consists of the histone methyltransferase EZH2 and the core subunits EED and SUZ12. Aberrant PRC2 activity is associated with epigenetic silencing of tumor suppressor genes and tumor progression. Recently, a non-canonical PRC2 has been described, harboring EZH1 instead of EZH2 as the histone methyltransferase. Interestingly, inhibition of EZH2 did not result in sufficient re-expression of PRC2-target genes, which might be explainable by compensatory function of EZH1. Consequently, combined EZH1/EZH2 inhibition has been identified as an attractive therapeutic strategy, which resulted in the development of dual EZH1/EZH2 inhibitors. Currently, there are no data available regarding EZH1 and BTC. Hence, in this study, we aim for an initial investigation of the effect of dual EZH1/EZH2 inhibition in BTC. Additionally, we want to clarify, whether EZH1 is expressed in BTC patient samples and whether EZH1 expression can be associated with clinical BTC progression and aggressiveness.

#### Methods

The in vitro experiments were performed using a comprehensive BTC cell line model (n = 9 BTC cell lines and n = 1 non-tumor cell line). mRNA levels of EZH1 were measured using rtPCR.Protein levels of EZH1/EZH2 and H3K27me1/2/3 were analyzed using Western Blot. Cytotoxic effects of Valemetostat were evaluated via a resazurin assay.

#### Results

We found that EZH1 and EZH2 are heterogeneously expressed in BTC cells. Similarly, H3K27me1/2/3 were expressed in a cell line-dependent manner. Interestingly, we found no correlation between expression levels. Next, we tested whether the dual EZH1/2 inhibitor Valemetostat shows cytotoxicity in BTC cells. Therefore, we treated the cells with different concentrations of Valemetostat for 72 hours. However, no specific cytotoxic effect was found. Therefore, we established a new protocol that allowed long-term incubation of BTC cells with Valemetostat for up to 14 days. Using this new protocol, we observed a significant decline in cell viability in selected cell lines following treatment for 10 to 14 days.

#### Conclusions

We here demonstrate that EZH1 is expressed in BTC cells. However, we found no correlation between EZH1 expression and protein levels of EZH2 and H3K27me1/2/3. Furthermore, we found that long-term incubation with Valemetostat results in cell line-dependent decline of cell viability. In a next step, we aim to test whether treatment of BTC cells with Valemetostat results in reduced H3K27me1/2/3 levels and re-expression of EZH1/EZH2 target genes - in comparison to the effect of sole EZH2 inhibition. We will complement these data by double EZH1/EZH2 knockdown experiments. Lastly, we will analyze EZH1 expression in BTC patient samples and investigate, whether EZH1 expression can be associated with clinical outcome.

# PREVALENCE OF METABOLIC DYSFUNCTION-ASSOCIATED FATTY LIVER DISEASE (MAFLD) AND EFFECTS OF GLUCOSE-LOWERING MEDICATIONS IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

<u>Andrea Mega</u><sup>1</sup>, Elmar Aigner<sup>2</sup>, Florian Koutny<sup>3</sup>, Sophie Gensluckner<sup>2</sup>, Fabio Vittadello<sup>4</sup>, Marco Dauriz<sup>5</sup>

<sup>1</sup>Department of Gastroenterology, Regional Hospital, Bolzano, Italy; <sup>2</sup>First Department of Internal Medicine, University clinic of Salzburg, Paracelsus Medical University Salzburg, Salzburg, Austria; <sup>3</sup>Department of Internal Medicine 2, Gastroenterology and Hepatology and rheumatology, Karl Landsteiner University of Health Sciences, University Hospital of St. Poelten, Iower Austria; <sup>4</sup>Explora-Research and Statistical Analysis, Padua, Italy; <sup>5</sup>Department of Internal Medicine, Medicine, Regional Hospital, Bolzano, Italy; Contact: andrea.mega@sabes.it

#### Objective

The aim of this study is to investigate the effects of antidiabetics on liver steatosis and fibrosis in a big court of patients with type 2 diabetes (T2D) and non alcoholic fatty liver diseae (NAFLD) and to evaluate the correlatons between liver stiffness and other clinical and biochemical parameters in T2D patients

#### Methods

101 consecutively patients affected by T2D and concomitant liver steatosis were enrolled from november 2022 to november 2023. All patients underwent biohumoral samplings and anthropometric values, upper abdominal ultrasound (for steatosis) and fibroscan at T0. NFS score, APRI score, FIB-4 and NAFLD score for liver fibrosis were calculated. All the enrolled patients started treatment with antidiabetic therapy with one or more of the new oral antidiabetics (PPAR-gammaAr; SGLUT-2 In; GLP1-ra) at T0. All the above data were then collected at 6 months and 12 months after the start of the T2DM treatment

#### Results

37.9% of the patients presented metabolic syndrome hepatic steatosis at T0. A significant reduction in GGT and ALT was documented at T6. In addition, a significant reduction in triglyceride levels, glycated Hb and blood glucose was documented at the end of the follow up (T12). With regard to fibrosis, APRI score and NFS score showed a significant reduction at T6 and T12, whereas no significant changes in liver stiffness (LS) were observed. No drugs correlated with changes in APRI score or NFS score, or with LS, probably due to the small sample size studied

#### Conclusions

Diabetes, metabolic syndrome and NAFLD/NASH are often related (1). Treatment of diabetes improves blood sugar but also liver metabolism (2). The improvement of liver metabolism and the decrease in liver inflammation leads to rearrangement of free fatty acids with their decrease and improvement of the general metabolism as well as the systemic inflammatory state (1, 2). A longer follow-up is needed to define the effects of T2D treatment on liver fibrosis

- Binet Q, Loumaye A, Preumont V, Thissen JP, Hermans MP, Lanthier N. Non- invasive screening, staging and management of metabolic dysfunction-associated fatty liver disease (MAFLD) in type 2 diabetes mellitus patients: what do we know so far ? Acta Gastroenterol Belg. 2022 Apr-Jun;85(2):346-357. doi: 10.51821/85.2.9775. PMID: 35709779
- Binet Q, Loumaye A, Hermans MP, Lanthier N. A Cross-sectional Real-life Study of the Prevalence, Severity, and Determinants of Metabolic Dysfunction- associated Fatty Liver Disease in Type 2 Diabetes Patients. J Clin Transl Hepatol. 2023 Nov 28;11(6):1377-1386. doi: 10.14218/JCTH.2023.00117. Epub 2023 Jun 30. PMID: 37719967; PMCID: PMC10500296

#### Risk of Interprosthetic Femur Fracture Is Associated with Implant Spacing - A Biomechanical Study

Mischa Mühling<sup>1, 2</sup>, Sabrina Sandriesser<sup>1, 2</sup>, Claudio Glowalla<sup>3</sup>, Sven Herrmann<sup>1,2</sup>, Peter Augat<sup>1, 2</sup>, Sven Hungerer<sup>1, 2, 3</sup>

<sup>1</sup>Institute for Biomechanics, BG Unfallklinik Murnau, Germany; <sup>2</sup>Institute for Biomechanics, Paracelsus Medical University Salzburg, Austria; <sup>3</sup>Department of Arthroplasty, BG Unfallklinik Murnau, Murnau, Germany; Contact: mischa.muehling@bgu-murnau.de

#### Objective

Ipsilateral revision surgeries of total hip or knee arthroplasties due to periprosthetic fractures or implant loosening are becoming more frequent in aging populations. Implants in revision arthroplasty usually require long anchoring stems. Depending on the residual distance between two adjacent knee and hip implants, we assume that the risk of interprosthetic fractures increases with a reduction in the interprosthetic distance. The aim of the current study was to investigate the maximum strain within the femoral shaft between two ipsilateral implants tips.

#### Methods

A simplified physical model consisting of synthetic bone tubes and metallic implant cylinders was constructed and the surface strains were measured using digital image correlation. The strain distribution on the femoral shaft was analyzed in 3-point- and 4-point-bending scenarios. The physical model was transferred to a finite element model to parametrically investigate the effects of the interprosthetic distance and the cortical thickness on maximum strain. Strain patterns for all parametric combinations were compared to the reference strain pattern of the bone without implants.

#### Results

The presence of an implant reduced principal strain values but resulted in distinct strain peaks at the locations of the implant tips. A reduced interprosthetic distance and thinner cortices resulted in strain peaks of up to 180 % compared to the reference. At low cortical thicknesses, the strain peaks increased exponentially with a decrease in the interprosthetic distance. An increasing cortical thickness reduced the peak strains at the implant tips.

#### Conclusions

A minimum interprosthetic distance of 10 mm seems to be crucial to avoid the accumulation of strain peaks caused by ipsilateral implant tips. Interprosthetic fracture management is more important in patients with reduced bone quality.

#### Acknowledgements

The authors would like to thank Martin Winkler and Johanna Finkel for their preliminary work in this project.

- Kosters, C.; den Toom, D.; Metzlaff, S.; Daniilidis, K.; Barz, L.; Rosslenbroich, S. Peri- and Interprosthetic Femoral 1. Frac-tures-Current Concepts and New Developments for Internal Fixation. J. Clin. Med. 2022, 11, 1371. https://doi.org/10.3390/jcm11051371.
- Loucas, M.; Loucas, R.; Akhavan, N.S.; Fries, P.; Dietrich, M. Interprosthetic Femoral Fractures Surgical Treatment in 2. 21514593211013790. Geriatric Patients. Geriatr. Orthop. Surg. Rehabil. 2021. 12. https://doi.org/10.1177/21514593211013790.
- Lipof, J.S.; Amitai, A.D.; Judd, K.T.; Gorczyca, J.T. Radiographic Risk Factors For Interprosthetic Femur Fractures. Iowa Orthop. J. 2017, 37, 35–39. Soenen, M.; Baracchi, M.; De Corte, R.; Labey, L.; Innocenti, B. Stemmed TKA in a femur with a total hip arthroplasty: 3.
- 4. distance J. Arthroplast. ls there safe between the stem tips? 2013. 28, 1437–1445. а https://doi.org/10.1016/j.arth.2013.01.010.
- 5. Stoffel, K.; Sommer, C.; Kalampoki, V.; Blumenthal, A.; Joeris, A. The influence of the operation technique and implant used in the treatment of periprosthetic hip and interprosthetic femur fractures: A systematic literature review of 1571 cases. Arch. Orthop. Trauma Surg. 2016, 136, 553–561. https://doi.org/10.1007/s00402-016-2407-y. Mühling, M.; Sandriesser, S.; Glowalla, C.; Herrmann, S.; Augat, P.; Hungerer, S. Risk of Interprosthetic Femur Fracture Is Associated with Implant Spacing—A Biomechanical Study. J. Clin. Med. 2023, 12, 3095.
- https://doi.org/10.3390/jcm12093095

## Trauma-focused art therapy in patients with miscarriage or stillbirth: Psychometric treatment effects of the randomized controlled MALT! study

<u>Markus Müller</u><sup>1</sup>, Barbara Stein<sup>1</sup>, Andrea Dorenkamp<sup>1</sup>, Bernadette Enders<sup>1</sup>, Luis Gerber<sup>2</sup>, Michael Gröger<sup>3</sup>, Alexandra Hass<sup>1</sup>, Olaf Prusik-Lutz<sup>1</sup>, Peter Radermacher<sup>3</sup>, Sofie Roecken<sup>1</sup>, Nicolas Rohleder<sup>2</sup>, Sophia Urbanczyk<sup>1</sup>, Fabian Zink<sup>3</sup>, Christiane Waller<sup>1</sup>

<sup>1</sup>Department of Psychosomatic Medicine and Psychotherapy, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Chair of Health Psychology, Friedrich-Alexander-University, Erlangen, Germany;

<sup>3</sup>Institute for Anesthesiological Pathophysiology and Process Engineering, Ulm University, Germany;

Contact: markus.mueller@klinikum-nuernberg.de

#### Objective

Women who experienced a miscarriage or stillbirth are at elevated risk of suffering from symptoms of posttraumatic stress, anxiety or depression.[1] While watchful waiting is often recommended, art therapy offers the opportunity to process and integrate the experience. The aim of the MALT! study was to examine whether trauma symptoms decrease to a greater extent after an art therapy intervention compared to a watchful waiting condition.

#### Methods

In a randomized controlled trial, women were assigned to either an eight-week outpatient one-on-one traumafocused art therapy (AT)[2] or a watchful waiting (WW) group within six months after miscarriage/stillbirth. Participants were examined at the beginning (T1) and after three months (T2). At T1, a control group (CG) of women without miscarriage/stillbirth was recruited. Primary outcome measure was change in post-traumatic symptoms, using the Impact of Event Scale - Revised (IES-R). Secondary outcome measures included depression, using the Patient Health Questionnaire 9 (PHQ-9). Statistical analyses were carried out crosssectionally using ANCOVAs and longitudinally using mixed models with repeated measures (intent-to-treat).

#### Results

For the study, 41 women after miscarriage/stillbirth were randomized, 20 in the AT group (age M = 33.5; SD = 5.7), and 21 the WW group (age M = 34.7; SD = 4.6,. The CG consisted of 28 women (age M = 27.3; SD = 1.4). At T1, women after miscarriage/stillbirth showed significantly higher scores than the CG in the PDS, IES-R and PHQ (all p < .001) with high effect sizes (all Cohen's d > 1.00). From T1 to T2, we found significant changes in IES-R (main effects for TIME), but no significant interactions between the AT and the WW group. Post-hoc analyses showed greater reductions in the symptoms of intrusion for the AT group (d = -1.0) than for the WW group (d = -0.5). In exploratory analyses, we found that for women with depression at T1, intrusions were only reduced by AT (d = -0.9) and not by WW (d = +0.1).

#### Conclusions

Women after miscarriage or stillbirth were more psychologically and biologically stressed than women without an event. MALT! art therapy lead to a significant reduction in intrusions, especially in participants with depression. Results indicate that art therapy plays an important role in the processing of traumatic events such as stillbirth and miscarriage and is feasible as a method for the prevention of PTSD symptoms.

#### Acknowledgements

This work was funded by STAEDTLER Stiftung, Nuremberg, Germany

- 1. Quenby S, Gallos ID, Dhillon-Smith RK, et al. Miscarriage matters: the epidemiological, physical, psychological, and economic costs of early pregnancy loss. Lancet. 2021;397(10285):1658-1667.
- 2. Roecken S, Prusik-Lutz O, Müller MM. MALT! Kunsttherapie als präventives Angebot nach einer Fehlgeburt. Psychotherapie im Dialog. 2024;24(01).

#### The voltage-gated proton channel discovers its family and function

#### Boris Musset<sup>1</sup>

<sup>1</sup>CPPB, Institute of Physiology, Pathophysiology and Biophysics, Paracelsus Medical University, Nuremberg, Germany; Contact: Boris.Musset@pmu.ac.at

#### Objective

In each species investigated, there is only a single gene representing voltage-gated proton channels. This dogma is reversed by our discovery of a whole new ion channel family.

#### Methods

We used modern bioinformatics, patch-clamp, and MD simulations to show the structure and function of proton channels.

#### Results

At least 4 new voltage-gated proton channels have been discovered.

#### Conclusions

HVs have numerous connections to human health (Immune system, cancer, reproduction, energy homeostasis ...)The full range of implications of HVs tissues/cells are up to date unknownNot every cell or species express HVsCommonly used standard model systems as: Drosophila melanogaster, Saccharomyces cerevisiae, Caenorhabditis elegans, Dictyostelium discoideum, show no indication of HVs.HVs evolved before 1.8 billion years during the appearance of Eukaroytes.Metabolically active cells and/or gametes use HVs to regulate membrane potential and/or pH homeostasis.After the Perm-Trias mass extinction (2000 ppm CO2 atmosphere//95 % of marine species died) Molluscs recovered well, potentially due to better adaptationProton channels might function as a scaffold for additional protein tasks.

#### References

1. Chaves, G., C. Jardin, C. Derst, and B. Musset. 2023. Voltage-Gated Proton Channels in the Tree of Life. Biomolecules. 13:1035.

#### PATHOGENIC PENDRIN (SLC26A4) VARIANTS ARE FUNCTIONALLY RESCUED BY CARFILZOMIB

Houssein Nasser<sup>1</sup>, Emanuele Bernardinelli<sup>1</sup>, Florian Huber<sup>1</sup>, Martin Jakab<sup>2</sup>, Sebastian Roesch<sup>3</sup>. Silvia Dossena<sup>1,4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, Paracelsus Medical University, Salzburg;

<sup>2</sup>Institute of Physiology and Pathophysiology, Paracelsus Medical University, Salzburg;

<sup>3</sup>Department of Otorhinolaryngology, Paracelsus Medical University, Salzburg; <sup>4</sup>FIZ - Regenerative Medicine & Novel Therapies (RM & NT), Paracelsus Medical University, Salzburg;

Contact: houssein.nasser@pmu.ac.at

#### Objective

Approximately 466 million people globally, or 5% of the world's population, experience disabling hearing loss, which significantly impacts adults and notably about 34 million children (1). Our research focuses on Pendrin. a vital protein for ion management in the inner ear, thyroid, and kidneys. Genetic variants of Pendrin, frequently associated with Pendred syndrome and DFNB4, manifest in hearing and vestibular dysfunctions. These variants are often characterized by reduced functionality and diminished expression levels, hinting at potential degradation processes. This observation has prompted us to hypothesize that such degradation might occur through the ubiquitin-proteasome system (UPS), a hypothesis we aim to investigate further in our study, with the additional aim of understanding how to recover this decrease in functionality and expression.

#### Methods

To investigate pendrin ubiquitination, HEK 293 Phoenix cells were transfected for 48 hours with plasmid vectors encoding FLAG-tagged wild-type pendrin and the pathogenic variants PDS L236P and PDS R409H. Post-transfection, co-immunoprecipitation was performed using an anti-FLAG antibody, followed by Western blot analysis with anti-ubiquitin and anti-FLAG antibodies. We assessed the functionality of pendrin variants by measuring the iodide influx in transfected HEK 293 Phoenix cells. Additionally, we evaluated pendrin expression levels in transfected HeLa cells using EYFP emission intensity, which we normalized against DAPI emission intensity.

#### Results

The results highlight ubiguitination and a significant reduction in the expression levels of pathogenic variants of pendrin compared to the wild-type form, confirming their degradation mediated by the ubiquitin-proteasome system (UPS). Furthermore, it has been demonstrated that the use of carfilzomib, a proteasome inhibitor, effectively restored both functionality and compromised expression.

#### Conclusions

Our study demonstrates the involvement of the ubiquitin-proteasome system (UPS) in the degradation of pathogenic pendrin variants associated with sensorineural deafness. Additionally, we highlight the potential therapeutic benefit of utilizing carfilzomib, a proteasome inhibitor, in restoring pendrin expression and transport function. These findings offer novel insights into the treatment of hearing loss linked to pathogenic pendrin variants and propose the UPS as a promising target for future therapeutic interventions.

#### Acknowledgements

This research was funded in part by projects 2022-lif-004-DOSSENA and FIZ RM&NT Talent Pool SR 042023 to Silvia Dossena. The present address of S. Roesch is: ENT Department, University Hospital Regensburg, Regensburg, Germany.

#### References

1. C. Schmucker, P. Kapp, E. Motschall, J. Loehler & J. J. Meerpohl. Prevalence of hearing loss and use of hearing aids among children and adolescents in Germany: a systematic review (2019) doi: 10.1186/s12889-019-7602-7. PMID: 31533687

# WGS-based characterization of putative hypervirulent Klebsiella pneumoniae identified in a tertiary care hospital in Germany

<u>Bernd Neumann</u><sup>1</sup>, Felix Aurnhammer<sup>1</sup>, Lisa Marr<sup>1</sup>, Thomas Kohl<sup>2</sup>, Stefan Niemann<sup>2</sup>, Anca Rath<sup>3</sup>, Bärbel Kieninger<sup>3</sup>, Wulf Schneider-Brachert<sup>3</sup>, Jörg Steinmann<sup>1</sup>

<sup>1</sup>Institute of Clinical Microbiology, Infectious Diseases and Infection Control, Paracelsus Medical University, Nuremberg General Hospital, Nuremberg, Germany;

<sup>2</sup>Molecular and Experimental Mycobacteriology, Research Center Borstel, Leibniz Lung Center, Borstel, Germany; German Centre for Infection Research (DZIF), Partner Site Hamburg-Lübeck-Borstel-Riems, Germany;

<sup>3</sup>Department of Infection Control and Infectious Diseases, University Hospital Regensburg, Regensburg, Germany; Contact: bernd.neumann@klinikum-nuernberg.de

#### Objective

Hypervirulent Klebsiella pneumoniae strains (hvKp), in contrast to classical K. pneumoniae strains, can cause invasive community-acquired infections in healthy patients of all ages. In this study, K. pneumoniae isolates from routine microbiological diagnostics were tested via string-test for hypermucoviscous phenotype and PCR for virulence genes to screen for hvKp, which were further analyzed by whole-genome sequencing (WGS). The prevalence of "hidden" hypervirulent hvKp, including non-invasive strains and their population structure should be analyzed in a tertiary-care hospital in Southern Germany(1).

#### Methods

For the study period 1.5 years, all K. pneumoniae isolates were string-tested. A multiplex PCR for hvKp genes was applied to all string-test positive isolates. PCR-positive isolates were subjected to WGS to determine genotyping and phylogeny.

#### Results

A total of 10.9% (331/3044) isolates with hmKp phenotype were detected by string-test. The patients' age rangend from 0 to 95 years, with a mean of 69 years. In total, 13.3% (44/331) isolates were tested positive by PCR for genes associated with hvKp. cgMLST revealed that 41.5% of sequenced isolates belonged to international hvKp clonal lineages ST23/K1 with high virulence scores and close phylogenetic relationships. In contrast, 22.6% of isolates belonged to the ST86/K2 with lower virulence scores. Isolates of liver abscesses (7.5%) belonged to ST23, ST25 and ST268, but without phylogenetic relatedness to other isolates.

#### Conclusions

We identified hypervirulent K. pneumoniae within the study period with an overall prevalence of 1.4%. No transmissions were identified. Isolates were assigned to the international hvKp lineage ST23/K1 and lineage ST86, bearing the potential to spread further in the community. The known association of multidrug-resistance ST86, was not observed in this study. Presented cases of liver abscesses seem to represent individual occurrences, so far. In conclusion, the screening of hmKp phenotypes in routine diagnostics seems to be a suitable surveillance method. Further, the population size and structure of "hidden" putative hvKp were more complex than expected.

#### References

1. Neumann et al 2023 (https://www.nature.com/articles/s41598-023-46221-w)

# Impact of Mesenchymal Stem Cells Derived Laminin-Binding Extracellular Vesicles on Schwann Cells and in Peripheral Nerve Regeneration Processes

#### Mai Quyen Nguyen<sup>1,2</sup>

<sup>1</sup>1Ludwig Boltzmann Institute for Experimental and Clinical Traumatology, Donauschingenstraße 13, 1020 Vienna, Austria; <sup>2</sup>Austrian Cluster for Tissue Regeneration;

Contact: mai.quyen.nguyen@trauma.lbg.ac.at

#### Objective

Injuries to peripheral nerves is a common health problem, often resulting in sensory and motor dysfunction in respective parts of the body. One of the most important facilitators of the regeneration process are Schwann cells (SCs), which basement membrane is chiefly comprised of laminin. Extracellular vesicles (EVs) are considered playing an important role in the intercellular communication and transfer of biological information. Especially, Mesenchymal stem cell-derived EVs (MSC-EVs) have been identified as a promising novel therapeutic option due to their potential function as target-directed drug delivery vehicles. However, the precise delivery of EVs upon administration is still a huge challenge due to the accumulation of EVs at off-target sites. To remedy this problem, this study focuses on the production of laminin-binding EVs derived from MSCs by modification of the large extracellular loop (LEL) of the overexpressed EV surface marker protein CD81, from the tetraspanin protein family, to increase their binding affinity to laminin.

#### Methods

Specific CD81-LEL sequences are cloned into lentiviral vectors encoding the expression cassette for full-length CD81 proteins fused with eGFP or Akaluciferase under the control of human cytomegalovirus promoter. Stable cell lines are obtained upon transformation of Wharton's Jelly MSCs (WJ-MSCs). EVs derived by WJ-MSCs are further isolated and characterized by NTA. Their uptake by SCs is evaluated by detecting the emitted fluorescence with FACS.

#### Results

Our results obtained from NTA demonstrate the successful cloning and production of laminin-binding EVs. The positive tracking of fluorescence with FACS verifies their binding affinity on rodent laminin and their increased uptake and pro-regenerative effect on SCs.

#### Conclusions

This study represents a substantial advance for further investigation on EVs regarding their influence on peripheral nerve regeneration processes.

#### References

 Vogt S, Bobbili MR, Stadlmayr G, Stadlbauer K, Kjems J, Rüker F, Grillari J, Wozniak-Knopp G. An engineered CD81based combinatorial library for selecting recombinant binders to cell surface proteins: Laminin binding CD81 enhances cellular uptake of extracellular vesicles. J Extracell Vesicles. 2021 Sep;10(11):e12139. doi: 10.1002/jev2.12139. PMID: 34514736; PMCID: PMC8435527.

# Concentrations of antiseizure medications during pregnancy and breastfeeding in women with epilepsy: evaluation of acute side effects and systematic follow-up of development of breasted infant. A international multicenter study.

<u>Bruna Nucera<sup>1,2</sup></u>, Barbara Mostacci<sup>3</sup>, Valentina Franco<sup>4,5</sup>, Gudrun Kalss<sup>6</sup>, Fabrizio Rinaldi<sup>1,7</sup>, Fedele Dono<sup>8</sup>, Luca Panebianco<sup>2,9</sup>, Georg Zimmermann<sup>10,11</sup>, Torbjörn Tomson<sup>12</sup>, Eugen Trinka<sup>6</sup>

<sup>1</sup>Department of Neurology, Hospital of Merano-Meran (SABES-ASDAA), Merano-Meran, Italy; <sup>2</sup>Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Istituto di Ricerca e Cura a Carattere Scientifico (IRCCS) Istituto delle Scienze Neurologiche di Bologna, Reference Center for Rare and Complex Epilepsies, Full Member of the ERN EpiCARE, Bologna, Italy; <sup>4</sup>Department of Internal Medicine and Therapeutics, Clinical and Experimental Pharmacology Unit, University of Pavia, Pavia, Italy; <sup>5</sup>IRCCS Mondino Foundation, Pavia, Italy; <sup>6</sup>Department of Neurology, Christian Doppler University Hospital, Paracelsus Medical University and Centre for Cognitive Neuroscience, Member of the ERN EpiCARE, Salzburg, Austria; <sup>7</sup>Paracelsus Medical University, Nuernberg, Germany; <sup>8</sup>Department of Neuroscience, Imaging and Clinical Science, "G. D'Annunzio" University of Chieti-Pescara, Chieti, Italy; <sup>9</sup>Department of Radiological Functions, Hospital of Merano (SABES-ASDAA), Merano-Meran, Italy; <sup>10</sup>Team Biostatistics and Big Medical Data, Intelligent Data Analytics (IDA) Lab Salzburg, Paracelsus Medical University, Salzburg, Austria; <sup>11</sup>Research and Innovation Management, Paracelsus Medical University Salzburg, Salzburg, Austria; <sup>12</sup>Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. Contact: bruna.nucera@stud.pmu.ac.at

#### Objective

Despite that recent position papers and guidelines recommend women with epilepsy (WWE) to breastfeed and highlight the benefits of breastfeeding, for both mother and child, only 42% of WWE breastfeed at three months. Several factors are involved, but the main one is the misconception that drugs taken by the mother are transferred to the baby through breastmilk and could cause acute or long-term adverse events. Although different studies have quantified antiseizure medications (ASMs) in breastmilk of WWE, there are limited data on the safety of specific ASMs during lactation, based on clinical experience, case reports and case series or observational studies. In addition, data on the possible effect of drugs on a child's neuro-cognitive development are also very scarce. Related to the premise, the objective of the current study is to characterize the second and third-generations ASMs concentrations in breastmilk and nursing-infant's plasma and assess adverse effect on the child's exposure to maternal drugs.

#### Methods

Women of childbearing age with a confirmed diagnosis of epilepsy (according to the ILAE 2014 practical definition), any type of seizure (focal or generalized), age > 18 years, on mono-or polytherapy with ASMs will be selected and enrolled. Breast milk samples, maternal and infant serum/plasma for quantification of the ASMs concentration will be collected until 4-8 weeks postpartum. Short and long-term adverse events on infant will be evaluated. Short-term breastfed infant outcomes are both clinical (impaired suckling, rash, sluggish, hypotonic, sedated, tremor, unmotivated sobbing) and laboratory tests (elevated platelet counts, alterations on electrolytes or hepatic profile enzymes). The long-term effect includes neurodevelopmental outcome of the infant at 12-24 months will be assessed using clinical scales.

#### **Results**

This is an international, multicenter, prospective study, which will be developed involving four epilepsy centers in Austria and Italy. The number of cases required to calculate the statistical power of the study is difficult to realize because it depends on the number of WWE wishing to become pregnant. However, considering that the reported rate of use of ASMs in pregnant women (3-4 cases per 1000) and the 2022 birth rate in the centers involved (total of 16.210 pregnancies), we consider enrolling approximately 80 women and children. The study received the approval of the South Tyrol Ethics Committee in November 2023 and was discussed in December 2023 in Salzburg Ethics Committee. It also received funding from the South Tyrol Health Authority.

#### Conclusions

The expected results of this study may be essential to guide epileptologists in the management of WWE taking ASMs during breastfeeding and to provide them with information on the safety of breastfeeding.

#### References

1

Veiby G, Bjørk M, Bernt A. Engelsen, Nils Erik Gilhus, Epilepsy and recommendations for breastfeeding, Seizure 28 (2015) 57–65, http://dx.doi.org/10.1016/j.seizure.2015.02.013

## Digital assessment of real-world walking activity in pulmonary hypertension: A prospective bicenter clinical trial

<u>Simone Panzer</u><sup>1</sup>, Nina Lutz<sup>4</sup>, Friederike Klenner<sup>2</sup>, Axel Hueber<sup>3</sup>, Jochen Klenk<sup>5</sup>, Alessandro Ghiani<sup>4</sup>, Claus Neurohr<sup>4</sup>, Simon Jäger<sup>1,4</sup>

<sup>1</sup>Institute of Clinical Pharmakology, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>ASV PH Ambulance, Medical Clinic 3, Klinikum Nuremberg, Nuremberg, Germany; <sup>3</sup>Rheumatology, Klinikum Nuremberg, Nuremberg, Germany; <sup>4</sup>RBK Lung Center, Robert Bosch Hospital, Stuttgart, Germany; <sup>5</sup>Robert Bosch Hospital, Stuttgart, Germany; Contact: simone.panzer@pmu.ac.at

#### Objective

Pulmonary hypertension (PH) has a prevalence of around 1% in the global population and comprises five main clinical subtypes that reflect a different pathophysiology. Shortness of breath and fatigue are the most important symptoms, and the significant loss of mobility makes it difficult to perform tasks of daily living (1,2). The clinical endpoints currently used in PH studies (e.g. six minute walk test) reflect impairments in daily mobility only to a limited extent. A digital recording of activity in the home environment over a longer period of time is therefore an attractive way to quantify the actual loss of mobility in more accurately. The aim of this study is to assess whether digital mobility outcomes (DMO) are suitable for monitoring the response to treatment, predicting the course of the disease or whether they can be helpful in the risk stratification of patients with PH.

#### Methods

This prospective observational study in pre-capillary PH is conducted at Klinikum Nuremberg and Robert Bosch Hospital in Stuttgart. At each time of medically indicated right heart catheterization, real life walking activity for seven consecutive days will be recorded with the Axivity AX6 sensor and characterized through eight newly validated DMOs: walking speed, step count, number of walking episodes, walking episode duration, cadence, number of body turns, stride duration, stride length (3). Primary endpoint will be the correlation of walking speed with hemodynamic data from RHC in a baseline and follow-up examination after 3 months. Along patient-reported outcomes, further clinical and laboratory data will be analysed in depth to characterize their relationship to real-world walking activity. Sample size calculation showed that 70 patients have to be included over a recruitment period of two years. Extensive analysis of the sensor's raw data will involve two different algorithms to interpret walking parameters.

#### Results

The study was successfully initiated at the two study centers in April 2024, with the first patient enrolled on April 19th.

#### Conclusions

This study will generate comprehensive real-world data on mobility of PH patients and the correlation and usefulness of DMO in research and clinical practice. In addition, a structured database for patients with PH in Stuttgart and Nuremberg will be implemented to facilitate further research, also comprising real-world effectiveness of PH-specific medications.

- Galiè N, Humbert M, Vachiery JL, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). Eur Respir J. Oct 2015;46(4):903-75. doi:10.1183/13993003.01032-2015
- 2. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. Eur Respir J. Jan 2019;53(1). doi:10.1183/13993003.01913-2018
- 3. Polhemus A, Ortiz LD, Brittain G, et al. Walking on common ground: a cross-disciplinary scoping review on the clinical utility of digital mobility outcomes. npj Digital Medicine. 2021;4(1):149. doi:10.1038/s41746-021-00513-5

#### Patient's Perspective and Expectations in Dupuytren's Disease Treatment

#### Sonja Pelzmann

<sup>1</sup>Paracelsus Medical University, Doctoral Programme in Medical Science, Salzburg, Austria: <sup>2</sup>Wiener Gesundheitsverbund Klinik Ottakring, Institute for Physical Medicine & Rehabilitation; Contact: sonja.pelzmann@stud.pmu.ac.at

#### Objective

Severity and duration of occurrence of the characteristic nodules and cords are important factors in the treatment of the neoplastic fibromatosis Dupuytren's disease (DD). Nevertheless, clinical practice shows and confirms that health-related aspects also significantly influence treatment and outcomes. It is necessary to include the patient's expectations and perspectives in the treatment process to achieve positive and realistic results. Although these aspects and their influence are always part of the treatment, they are rarely investigated in this progressive, non-curable disease, and there is no systematic overview on the available evidence. The aim of this study was to identify, classify and present the evidence base of patients' perspectives, expectations, and needs in DD treatment.

#### Methods

Using a systematic search strategy multiple electronic databases (PubMed, CINAHL) and hand specific journals were searched for relevant and available information. The studies were classified using the Oxford Level of Evidence (LoE) and their findings were linked to the International Classification of Functioning. Disability and Health (ICF) and classified into four for categories ("expectations of treatment", "expectations of results", "need during treatment" and "knowledge for the future") relevant for the treatment.

#### Results

Of the 1.604 publications found, five publications could be included in this review. The sample size of the five included studies ranged from seven to 24 patients and the LoE from IIIb to IV, respectively. The 34 findings were linked to the ICF-components "body function", "body structure", "activities" and "participation" as well as the ICF-domains "neuromusculoskeletal and movement-related functions", "structures related to movement", "sensory function and pain", "learning and applying knowledge", "communication", "self-care" and "interpersonal interactions and relationships". The results of this review show that aspects such as involvement in decisions, communication of knowledge about the disease, treatment options including possible outcomes are essential for patients in DD treatment.

#### Conclusions

The results demonstrate the inadequate evidence base of health-related aspects in DD treatment. In addition to physiologically measurable parameter, health-related aspects are an issue and expected by patients with DD. Nevertheless, it is difficult to determine the individual contribution of these topics and to assess their value to subsequently use them adequately for the benefit of patients in DD treatment or developing and improving health services.

- 1.
- 2.
- 3.
- 4.
- 5.
- Engstrand C, Krevers B, Nylander G, Kvist J. Hand Function and Quality of Life Before and After Fasciectomy for Dupuytren Contracture. J Hand Surg Am. 2014 Jul;39(7):1333-1343.e2. Townley WA, Baker R, Sheppard N, Grobbelaar AO. Dupuytren's contracture unfolded. BMJ. 2006;332:397–400. Benson LS, Williams CS, Kahle M, Otr L, Cooper AC. Dupuytren's contracture. Journal of the American Academy of Othopaedic Surgeons. 1998;6(1):24–35. Pratt AL, Byrne G. The lived experience of dupuytren's disease of the hand. J Clin Nurs. 2009;18:1793–802. Turesson C, Kvist J, Krevers B. Experiences of men living with Dupuytren's disease—Consequences of the disease for hand function and daily activities. Journal of Hand Therapy. 2020;33(3):386–93. Wilburn J, McKenna SP, Perry-Hinsley D, Bayat A. The impact of dupuytren disease on patient activity and quality of life. Journal of Hand Surgery [Internet]. 2013;38(6):1209–14. Available from: http://dx.doi.org/10.1016/j.ibsa.2013.03.036 6. http://dx.doi.org/10.1016/j.jhsa.2013.03.036

- http://dx.doi.org/10.1016/j.jhsa.2013.03.036
  Engstrand C, Kvist J, Krevers B. Patients' perspective on surgical intervention for Dupuytren's disease experiences, expectations and appraisal of results. Disabil Rehabil. 2016;38(26):2538–49.
  Winberg M, Turesson C. Patients' perspectives of collagenase injection or needle fasciotomy and rehabilitation for Dupuytren disease, including hand function and occupational performance. Disabil Rehabil. 2022;
  Beecher SM, Wilkinson JE, Cuggy C, O'Shaughnessy M. Patient's perspective of treatment in Dupuytren's Disease: collagenase versus limited fasciectomy. Journal of Hand Surgery: European Volume. 2022;
  Turesson C, Kvist J, Krevers B. Patients ' needs during a surgical intervention process for Dupuytren ' s disease. Disabil Rehabil [Internet]. 2019;41(6):666–73. Available from: https://doi.org/10.1080/09638288.2017.1402095
  Molineux ML, McCann A, Cox RJ, Cross SM. "Extremely Painful, but Worth It": The Lived Experience of Receiving Collagenase as Treatment for Dupuytren Disease. Plast Reconstr Surg. 2022 Apr 1;149(4):913–7.

#### BARRIEREN DES ZUGANGS ZU PALLIATIVE CARE IM KRANKENHAUS

<u>Stefan Pitzer</u><sup>1</sup>, Patrick Kutschar<sup>1</sup>, Piret Paal<sup>2</sup>, Patrick Mülleder<sup>1</sup>, Stefan Lorenzl<sup>2</sup>, Paulina Wosko<sup>3</sup>, Jürgen Osterbrink<sup>1</sup>, Johannes Bükki<sup>4</sup>

<sup>1</sup>Institut für Pflegewissenschaft und -praxis, Paracelsus Medizinische Privatuniversität, Salzburg, Österreich;

<sup>2</sup>Institut für Palliative Care, Paracelsus Medizinische Privatuniversität, Salzburg, Österreich;

<sup>3</sup>Gesundheit Österreich GmbH, Wien, Österreich;

<sup>4</sup>Helios-Kliniken Schwerin, Zentrum für Palliativmedizin, Schwerin, Deutschland;

Contact: stefan.pitzer@pmu.ac.at

#### Objective

Palliative Care kann Leiden verringern, die Lebensqualität von Patient\*innen und deren Angehörigen verbessern sowie die Kosten am Lebensende reduzieren. Diese Erkenntnisse förderten die Implementierung neuer Palliative Care-Angebote und erleichterten den Zugang. Im klinischen Alltag hängt dieser jedoch meist von der konkreten Zuweisungspraxis ab, welche wiederum von institutionellen Faktoren, individuellen Überzeugungen und soziokulturellen Variablen beeinflusst wird. Der Zugang zu Palliative Care ist damitnach wie vor kontingent und zahlreiche stationär behandelten Patient\*innen erhalten keinePalliativversorgung. Eine systematische Aufbereitung potenzieller und existierender Barrieren könnte die Entwicklung von Strategien zu deren Überwindung befördern.

#### Methods

Systematisches Mixed-Methods-Review unter Verwendung eines integrierten konvergenten Ansatzes und thematischer Synthese. Die Datenbanken Cochrane Library, MEDLINE, CINAHL und PsycINFO wurden nach Studien des Zeitraums 10/2003 (letztes Review zu Zugangsbarrieren im Krankenhaus) bis 12/2020 durchsucht. Studien jeglichen Designs, die Evidenz zu Hindernissen des Zugangs zur im Krankenhaus verfügbaren Palliativversorgung bieten, wurden inkludiert.

#### Results

Nach initialem Screening von 3359 Datenbankeinträgen und Prüfung von 555 Volltexten wurden 79 Studien eingeschlossen. Die thematische Synthese ergab 149 zugangsbezogene Phänomene in sechs Hauptkategorien: (1) Soziodemografische Merkmale, (2) Gesundheitsbezogene Merkmale von Patient\*innen, (3) Individuelle Überzeugungen und Haltungen, (4) Interindividuelle Kooperation und Unterstützung, (5) Verfügbarkeit und Zuteilung von Ressourcen sowie (6) Emotionale Herausforderungen. Während die Ergebnisse für die meisten soziodemografischen Faktoren nicht konklusiv waren, kristallisierten sich folgende Barrieren heraus: eine andere Erkrankung als Krebs oder geringe Symptomlast, ein generell kurativer therapeutischer Fokus zulasten palliativer Maßnahmen, die Nicht-Akzeptanz der bestehenden Prognose, negative Wahrnehmungen zu Palliative Care, missverständliche Kommunikation und widersprüchliche Versorgungspräferenzen, fehlende Ressourcen, schlechte Koordination, unzureichendeFachkenntnisse sowie emotionales Unbehagen des medizinischen Personals.

#### Conclusions

Die Ergebnisse des Reviews verdeutlichen, dass der Zugang zur Palliativversorgung in Krankenhäusern mit diversen mehrdimensionalen Barrieren verbunden ist. Der Vergleich mit früheren Reviews zeigt zudem, dass zahlreiche dieser Barrieren über Jahrzehnte hinweg persistieren. Es besteht somit nicht nur praktischer Bedarf an Abbau und Prävention von Barrieren, sondern auch an Forschungsprojekten, welche die Mechnismen der Aufrechterhaltung dieser Barrieren durchleuchten.

## The assessment of biological maturation in academy soccer players: a comparison of two methods

Maximilian Plachel<sup>1</sup>, James O'Brien<sup>1</sup>, Thomas Stöggl<sup>1</sup>, Florian Wichlas<sup>2</sup>

<sup>1</sup>Red Bull Athlete Performance Center, Salzburg, Austria;

<sup>2</sup>Department of Orthopaedics and Traumatology, Salzburg University Hospital, Paracelsus Medical University, Salzburg, Austria;

Contact: maximilian.plachel@stud.pmu.ac.at

#### Objective

The calculation of biological age and maturity status is highly relevant in the care of adolescent soccer players, as this information can have an im-pact on performance and injury risk. While various methods exist for assessing this, they are often expensive and challenging to implement (i.e., X-ray and MRI). The aim of this study was to assess an ultrasound-based examination method for evaluating skeletal age (SA), predicted adult height (PAH) and the biological maturation status (BMS, % of PAH) in a group of selected youth soccer players.

#### Methods

A total of n = 17 youth male soccer players ( $\emptyset$  13.21 years; range 12.65-13.62 years) of an Austrian Soccer Academy (U14) completed two meas-urements to assess biological maturity status outcomes at the start of the season 2023/24. Each player underwent an MRI and ultrasound diagnos-tics with the SonicBone device (from BAUSportTM) of the left wrist. A Bland-Altman analysis was conducted to evaluate potential differences. Furthermore, the level of agreement between the methods was assessed.

#### Results

The Bland-Altman analysis showed that more than 95 % of the values in the graph are within the limits of agreement for the mean value in the esti-mation of SA, PAH and BMS. Correlations of SA (r=0,84; p<0,001), PAH (r=0.87; p<0.001) and BMS (r=0.85; p<0.001) were highly significant.

#### Conclusions

The results highlight the measures usefulness and suggest that the SonicBone can serve as an alternative predictor of biological maturation within elite youth soccer. The biological maturity status should be taken into consideration in talent selection to avoid discrimination between early and late developers.

#### References

1. Swain, M., Kamper, S. J., Maher, C. G., Broderick, C., McKay, D., & Henschke, N. (2018). Relationship between growth, maturation and musculoskeletal conditions in adolescents: a systematic review. British journal of sports medicine, 52(19), 1246-1252.

#### Minimally invasive ascending aortic surgery: operative and mid-term outcomes

<u>Francesco Pollari</u><sup>1</sup>, Matthias Angerer<sup>1</sup>, Wolfgang Hitzl<sup>2</sup>, Lucia Weber<sup>1</sup>, Joachim Sirch<sup>1</sup>, Theodor Fischlein<sup>1</sup>

<sup>1</sup>Cardiac Surgery, Paracelsus Medical University – Klinikum Nürnberg, Nuremberg, Germany;
<sup>2</sup>Research and Innovation Management (RIM), Team Biostatistics, Paracelsus Medical University, Salzburg, Austria;
Contact: francesco.pollari@klinikum-nuernberg.de

#### Objective

Minimally invasive approaches, such as partial-sternotomy (PS), could reduce the surgical trauma and were proven to be safe for valve-related procedures [1]. We aimed to investigate the in-hospital and mid-term outcomes of patients undergoing ascending aortic surgery (AAS) through a partial or a full-sternotomy approach through a propensity matching analysis.

#### Methods

We retrospectively included all patients (n=167), who underwent elective AAS for aneurysm in our institution between 2013 and 2020. Patients, who received a surgical treatment in cases of emergency situations for aortic dissection were excluded. Study population was divided in twogroups according to the surgical access (n=40 in partial sternotomy or "PS", and n=127 in full sternotomy, or "FS"). Due to the signibcant differences between the groups, a propensity matching 1:3 was applied. Age, BMI, gender and EuroSCORE II were used as covariate variables and the propensity score was computed based on the combined aortic valve operation. The Mahalanobis distance including the propensity score was used as distancecalculation method. The order for matching was done at random. No maximum number of iterations were set for the optimization algorithm. After propensity matching, only the preoperative EF was signibcantly different between the two groups. Inhospital complications, survival and reoperation at follow-up were investigated.

#### Results

No operation started with a partial upper sternotomy had to be converted into a median full sternotomy during the operation. The majority of patients were operated with a brachiocephalic cannulation (PS=70% vs FS=61%, p=0.3) and selective cerebral perfusion (PS=65% vs FS=58%, p=0.3) in both groups. A combined aortic valve surgery was performed in the 92% of patients in the PS-group and 76% of patients in FS-group (p=0.06). PS groupshowed higher X-clamp and cardiopulmonary bypass times (94.2 min vs. 83 min and 164.2 min vs. 126.8 min). Moreover, the mean postoperative ventilation time was signibcantly higher in the PS group (41.5 hours±98.8 versus 22.5 hours±58.5), however not affecting the length ofstay in ICU (3.6 days±4.7 versus 2.9 days±3.3; p=0.1). The incidence of bleeding, stroke and inhospital mortality were similar between PS and FS group (11% vs. 3%, 3% vs. 6%, 5% vs 3%, respectively). After a median follow-up of 2±1.98 years, the Kaplan-Meier analysis showed notsignibcant differences between the PS and FS group (log-rank, p=0.17) in term of survival. Cardiac reoperations were observed in 2 cases (6%) of the PS-group and in 8 (7%) of the FS group. In the PS group the reason accounted for hemodynamic relevant pericardial effusion, that was successfully treated with a subxiphoidal drainage placement. In the FS-group patients were reoperated because of pericardial effusion (n=3), sternal complications (n=2), tricuspidal valve insujciency (n=1), aortic valve replacement (n=1) or tube prosthesis endocarditis by Streptococcus gallolyticus (n=1).

#### Conclusions

The surgical ascending aorta replacement through a partial sternotomy is associated with longer operative times, but this does not affect the early as well as the long-term follow-up. Surgical ascending aorta replacement can be safely performed through a minimally invasive approach.

#### Acknowledgements

Part of the present study was performed in fulfillment of the requirements for obtaining the doctoral degree of M. Angerer.

#### References

1. Paparella D, Malvindi PG, Santarpino G, Moscarelli M, Guida P, Fattouch K et al. Full sternotomy and minimal access approaches for surgical aortic valve replacement: a multicentre propensity-matched study. Eur J Cardiothorac Surg 2020;57:709–16.

#### ATR-FTIR as tool for quality control of extracellular vesicles

<u>Alexander Popoff</u><sup>1</sup>, Sophie Haller<sup>1</sup>, Noosha Riazy-Hossfeld<sup>1</sup>, Cristian-Tudor Matea<sup>2</sup>, Eva Klinglmayer<sup>2,3</sup>, Nicole Meisner-Kober<sup>2,3</sup>, Daniela Schuster<sup>1</sup>

<sup>1</sup>Department of Pharmaceutical and Medicinal Chemistry, Institute of Pharmacy, Paracelsus Medical University, Strubergasse 21, 5020 Salzburg, Austria; <sup>2</sup>Department of Biosciences and Medical Biology, Paris Lodron University Salzburg, Salzburg, Austria.; <sup>3</sup>Ludwig Boltzmann Institute for Nanovesicular Precision Medicine, Hellbrunnerstrasse 34, 5020 Salzburg, Austria.; Contact: Alexander.Popoff@pmu.ac.at

#### Objective

Extracellular vesicles (EVs) represent an exciting new research field with great therapeutic and diagnostic potential. Especially in the field of drug delivery, they open a promising field as novel vehicles for active pharmaceutical ingredients beside their inherent biological activity. Like any product for pharmaceutical purpose, a proper quality control is vital. In this context, ATR-FTIR can be a valuable tool. Here we evaluate the potential of ATR-FTIR techniques to determine important quality control parameters like lipid and protein content and composition as well as purity. For this purpose, we applied multivariate analysis methods for the acquired IR spectra in conjunction with classical colorimetric and biochemical techniques. As test subject, we used milk-derived vesicles that were isolated by tangential flow filtration due to their easy accessibility and high yield.

#### Methods

In order to asses the quality of milk derived extracellular vesicles, ATR-FTIR spectra of the vesicles were acquired in conjunction with colorimetric BCA and SPV assays to determine the protein and lipid content. In addition, an SDS PAGE and a LC-MS/MS analysis of excised geld bands was performed to analyze to protein content of EV samples with different purity grades. ATR-FTIR spectra were analyzed using a PLSR model. The EV samples itself were purified by tangential flow filtration and purity was determined by size exclusion chromatography.

#### Results

The protein content of EV samples with different purity grades is changing. Samples with low purity show an accumulation of BSA and keratin. Regarding the protein and lipid content, the protein/lipid ratio is decreasing with higher purity which demonstrates that the protein/lipid ratio can be used for determination of EV sample purity. A prediction of purity using simply IR spectra in conjunction with a PLSR model is a suitable alternative. The test set gave a coefficient of determination of 0.78.

#### Conclusions

The Protein/lipid ratio can be predictive for EV sample purity, but requires additional markers for higher accuracy. EV sample purity can also be predicted by IR spectroscopy in conjunction with a PLSR model. However, a more diverse training set and proper post processing of the spectra may improve performance of the model. In addition, the PLSR model revealed spectral markers of purity, some of known biomolecules, others still need to be assigned to a certain class of biomolecules

- Busatto S, Vilanilam G, Ticer T, Lin WL, Dickson DW, Shapiro S, Bergese P, Wolfram J. Tangential Flow Filtration for Highly Efficient Concentration of Extracellular Vesicles from Large Volumes of Fluid. Cells. 2018 Dec 16;7(12):273. doi: 10.3390/cells7120273.
- Visnovitz T, Osteikoetxea X, Sódar BW, Mihály J, Lőrincz P, Vukman KV, Tóth EÁ, Koncz A, Székács I, Horváth R, Varga Z, Buzás EI. An improved 96 well plate format lipid quantification assay for standardisation of experiments with extracellular vesicles. J Extracell Vesicles. 2019 Jan 29;8(1):1565263. doi: 10.1080/20013078.2019.1565263.
- extracellular vesicles. J Extracell Vesicles. 2019 Jan 29;8(1):1565263. doi: 10.1080/20013078.2019.1565263.
  Mihály J, Deák R, Szigyártó IC, Bóta A, Beke-Somfai T, Varga Z. Characterization of extracellular vesicles by IR spectroscopy: Fast and simple classification based on amide and CH stretching vibrations. Biochim Biophys Acta Biomembr. 2017 Mar;1859(3):459-466. doi: 10.1016/j.bbamem.2016.12.005.
- Harald Martens, Reliable and relevant modelling of real world data: a personal account of the development of PLS Regression, Chemometrics and Intelligent Laboratory Systems, Volume 58, Issue 2,2001, Pages 85-95, ISSN 0169-7439, https://doi.org/10.1016/S0169-7439(01)00153-8.

#### Ozanimod – Safety and Efficacy in a real-world-MS-cohort

<u>Richard Friedrich Radlberger</u><sup>1</sup>, Ferdinand Otto<sup>1</sup>, Michael Seiberl<sup>1</sup>, Lara Bieler<sup>1</sup>, Tobias Moser<sup>1</sup>, Andrea Harrer<sup>1,2</sup>, Peter Wipfler<sup>1</sup>

<sup>1</sup>Department of Neurology, Christian Doppler Clinic, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Dermatology and Allergology, Paracelsus Medical University, Salzburg, Austria; Contact: dr.radlberger@gmx.net

#### Objective

The shift towards "early intense therapies" for multiple sclerosis (MS) raises concerns about immune competence during pandemics as caused for example by the severe acute respiratory syndrome coronavirus type two (SARS-CoV-2) in 2019. So safety issues, especially concerning a potentially compromised defence, are becoming more and more relevant. We aimed to assess the baseline demographics and disease characteristics, the safety profile as well as efficacy of ozanimod, one of the newer, more selective, sphingosine 1-phosphate (S1P)1 and S1P5 receptor modulators, under real world conditions, in comparison to the SUNBEAM and RADIANCE approval studies.

#### Methods

We included 19 MS patients (range 22-45 years; 84% female) treated with ozanimod from our MS outpatient clinic of which all finished the projected one year observation period. Safety parameters including laboratory monitoring especially considering liver enzymes, lymphocyte counts, blood pressure, infections and expanded disability status scale (EDSS) were monitored at baseline/before treatment initiation (T0) and at follow-up control visits every three months (T3M-T12M), magnetic resonance imaging (MRI) control scan was performed at T12M.

#### Conclusions

In this real-life study, our patient cohort was comparable to that recruited in the pivotal studies. Accordingly, our results in terms of side effects as well as disease-suppressing effects match the published findings.

- Comi G, Kappos L, Selmaj KW, Bar-Or A, Arnold DL, Steinman L, Hartung HP, Montalban X, Kubala Havrdová E, Cree BAC, Sheffield JK, Minton N, Raghupathi K, Ding N, Cohen JA; SUNBEAM Study Investigators. Safety and efficacy of ozanimod versus interferon beta-1a in relapsing multiple sclerosis (SUNBEAM): a multicentre, randomised, minimum 12-month, phase 3 trial. Lancet Neurol. 2019 Nov;18(11):1009-1020. doi: 10.1016/S1474-4422(19)30239-X. Epub 2019 Sep 3. PMID: 31492651.
- Cohen JA, Comi G, Selmaj KW, Bar-Or A, Arnold DL, Steinman L, Hartung HP, Montalban X, Kubala Havrdová E, Cree BAC, Sheffield JK, Minton N, Raghupathi K, Huang V, Kappos L; RADIANCE Trial Investigators. Safety and efficacy of ozanimod versus interferon beta-1a in relapsing multiple sclerosis (RADIANCE): a multicentre, randomised, 24-month, phase 3 trial. Lancet Neurol. 2019 Nov;18(11):1021-1033. doi: 10.1016/S1474-4422(19)30238-8. Epub 2019 Sep 3. PMID: 31492652.

# The Influence of Chronotype on Interictal Epileptiform Discharges and Sleep Quality in Individuals with Epilepsy: A Study of Sleep-EEG Findings

<u>Fabrizio Rinaldi</u><sup>1,3</sup>, Bruna Nucera<sup>1,3</sup>, Davide Flore<sup>1</sup>, Sara Mariozzi<sup>1</sup>, Luigi Ferini-Strambi<sup>4</sup>, Eugen Trinka<sup>5</sup>, Kneginja Richter<sup>6,7,8</sup>

<sup>1</sup>Department of Neurology, Hospital of Merano (SABES-ASDAA), Merano, Italy; <sup>2</sup>Paracelsus Medical University, 90419, Nuermberg, Germany; <sup>3</sup>Paracelsus Medical University, 5020 Salzburg, Austria; <sup>4</sup>IRCCS San Raffaele Scientific Institute, Department of Clinical Neurosciences, Neurology-Sleep Disorders Centre, Milan, Italy; <sup>5</sup>Department of Neurology, Christian Doppler University Hospital, Member of European Reference Network EpiCARE, Paracelsus Medical University, Centre for Cognitive Neuroscience, Salzburg, Austria; <sup>6</sup>University Clinic for Psychiatry and Psychotherapy, Klinikum Nuernberg, Paracelsus Medical University, Nuremberg, Germany; <sup>7</sup>Technische Hochschule Nürnberg Georg Simon Ohm; <sup>8</sup>CuraMed Tagesklinik Nürnberg GmbH; Contact: rinaldi.fabrizio.riccardo@gmail.com

#### Objective

The aim of this study is to investigate the influence of sleep characteristics in patients with suspected first seizure on data obtained from sleep-deprivation EEG (S-EEG) recordings.

#### Methods

44 patients (45.5% female, mean age 49 years) were prospectively enrolled. The following data were collected: seizure semiology, demographics, neuroimaging, S-EEG results (sleep parameters, number and characteristics of IEDs). Patients also completed questionnaires to assess chronotype, insomnia, excessive daytime sleepiness (ESD).

#### Results

An increased number of IEDs was found in patients with a morning chronotype compared to a serotonin chronotype (13.86±11.21 vs. 2.62±3.62, p=0.013), with no differences in sleep parameters, including total sleep time (TST) ( $20\pm17.95$  min vs. 18.5±15.51 min, p=0.875).Patients with ESD show a non-significant increase in TST ( $26.78\pm21.87$  vs. 18.23±13.60, p=0.228), with no difference in the number of IEDs ( $3.14\pm6.38$  vs. 7.97±25.33, p=0.62).Patients with insomnia had a higher number of IEDs ( $8\pm18.1$  vs.  $6.5\pm3.9$ , p=0.015) with no difference in TST ( $20.1\pm19.30$  vs. 18.95±16.44, p=0.78).

#### Conclusions

This study demonstrates a correlation between patient chronotype and the number of IEDs in S-EEG, which is higher in patients with morning chronotype. Insomniac patients also show an increased number of abnormalities, unlike ESD patients. Finally, no correlation was found between sleep duration and the number of IEDs, suggesting that the usefulness of this examination may be independently related to sleep deprivation, as opposed to sleep attained during recording.

- Nobili L, Frauscher B, Eriksson S, Gibbs SA, Halasz P, Lambert I, Manni R, Peter-Derex L, Proserpio P, Provini F, de Weerd A, Parrino L. Sleep and epilepsy: A snapshot of knowledge and future research lines. J Sleep Res. 2022 Aug;31(4):e13622. doi: 10.1111/jsr.13622. Epub 2022 Apr 29. PMID: 35487880; PMCID: PMC9540671.
- Peltola, M. E., Leitinger, M., Halford, J. J., Vinayan, K. P., Kobayashi, K., Pressler, R. M., Mindruta, I., Mayor, L. C., Lauronen, L., & Beniczky, S. (2023). Routine and sleep EEG: Minimum recording standards of the International Federation of Clinical Neurophysiology and the International League Against Epilepsy. Epilepsia, 64(3), 602–618.

#### Risk factors for prolonged length of stay after first single-level lumbar microdiscectomy

<u>Leonard Ritter</u><sup>1</sup>, Adrian Liebert<sup>1</sup>, Thomas Eibl<sup>1</sup>, Barbara Schmid<sup>2</sup>, Hans-Herbert Steiner<sup>1</sup>, Ghassan Kerry<sup>1</sup>

<sup>1</sup>Department of Neurosurgery, Paracelsus Medical University, Nuremberg, Germany; <sup>2</sup>Department of Neurology, Paracelsus Medical University, Nuremberg, Germany; Contact: leonard.ritter@stud.pmu.ac.at

#### Objective

The objective is to identify risk factors that potentially prolong the hospital stay in patients after undergoing first single-level open lumbar microdiscectomy.

#### Methods

A retrospective single-centre study was conducted. Demographic data, medical records, intraoperative course, and imaging studies were analysed. The outcome measure was defined by the number of days stayed after surgery. A prolonged length of stay (LOS) stay was defined as a minimum of one additional day beyond the median hospital stay in our patient collective. Bivariate analysis and multiple stepwise regression were used to identify independent factors related to the prolonged hospital stay.

#### Results

Two hundred consecutive patients who underwent first lumbar microdiscectomy between 2018 and 2022 at our clinic were included in this study. Statistical analysis of factors potentially prolonging postoperative hospital stay was done for a total of 24 factors, seven of them were significantly related to prolonged LOS in bivariate analysis. Sex (p = 0.002, median 5 vs. 4 days for females vs. males) and age (rs = 0.35,  $p \le 0.001$ , N = 200) were identified among the examined demographic factors. Regarding preoperative physical status, preoperative immobility reached statistical significance ( $p \le 0.001$ , median 5 vs. 4 days). Diabetes mellitus (p = 0.043, median 5 vs. 4 days), anticoagulation and/or antiplatelet agents (p = 0.045, median 5 vs. 4 days), and postoperative narcotic consumption ( $p \le 0.001$ , median 5 vs. 4 days) as comorbidities were associated with a prolonged hospital stay. Performance of nucleotomy (p = 0.023, median 5 vs. 4 days) was a significant intraoperative factor. After linear stepwise multivariable regression, only preoperative immobility ( $p \le 0.001$ ) was identified as independent risk factors for prolonged length of postoperative hospital stay.

#### Conclusions

Our study identified preoperative immobility as a significant predictor of prolonged hospital stay, highlighting its value in preoperative assessments and as a tool to pinpoint at-risk patients. Prospective clinical trials with detailed assessment of mobility, including grading, need to be done to verify our results.

- 1. Kuo CC, Hess RM, Soliman MAR, Khan A, Pollina J, Mullin JP. Predicting prolonged length of stay in patients undergoing transforaminal lumbar interbody fusion. Acta Neurochir (Wien) 2022;164(10):2655-2665. DOI: 10.1007/s00701-022-05334-3.
- Fisher C, Noonan V, Bishop P, et al. Outcome evaluation of the operative management of lumbar disc herniation causing sciatica. J Neurosurg 2004;100(4 Suppl Spine):317-24. DOI: 10.3171/spi.2004.100.4.0317.
- 3. Dewing CB, Provencher MT, Riffenburgh RH, Kerr S, Manos RE. The outcomes of lumbar microdiscectomy in a young, active population: correlation by herniation type and level. Spine (Phila Pa 1976) 2008;33(1):33-8. DOI: 10.1097/BRS.0b013e31815e3a42.

# Burden of spinal cord injury, 1990 2019: a systematic analysis for the Global Burden of Disease Study 2019

Mahdi Safdarian, Eugen Trinka, Aljoscha Thomschewski

<sup>1</sup>Department of Neurology, Christian Doppler Klinik , Paracelsus Medical University; Contact: m.safdarian@salk.at

#### Objective

Spinal cord injuries contribute to a major burden of disabilities causing health loss due to premature mortality and residualdisability While spinal injuries are a public health issue, no comprehensive report on their global temporal, spatial and demographic patterns hasnot been published We aimed to measure the global, regional, and national incidence, prevalence, and years lived with disability ( of spinalinjuries from 1990 to 2019 using data from the Global Burden of Diseases, Injuries, and Risk Factors Study.

#### Methods

Data on incidence, prevalence, and YLDs of spinal injuries were derived systematically from the GBD 2019 study Using theframework of GBD 2019 we provide numbers and age standardized rate changes with 95 uncertainty intervals (for incidence, prevalence, and YLDs of spinal injuries at neck level and below neck level globally and for 21 GBD regions and 204 countries and territories, among all agegroups and both sexes from 1990 to 2019 We tried to detect and report meaningful trends based on location, age and gender.

#### Results

Globally, there were 20 6 million 95 UI 18 9 to 23 6 individuals living with SCI in 2019 The incidence of SCI was 0 9 million cases 95UI 0 7 to 1 1 with an estimated 6 2 million 95 UI 4 4 to 8 1 YLDs SCI rates increased considerably from 1990 to 2019 for global prevalence81 5 UI 74 2 87 1 incidence 52 7 UI 30 3 69 8 and YLDs 65 4 UI 56 3 76 0 ))(Figures 1 2 However, global age standardized rates per100 000 population showed slighter changes for prevalence 5 8 UI 2 6 to 9 5 incidence 6 1 UI 17 2 to 1 5 and YLDs 1 5 UI 5 5 to 3 2 The incidence, prevalence, and YLDs of SCI have consistently been higher in men globally, with a slight and steady increase observed for bothsexes from 1990 to 2019 (Figure 3 In 2019 the number of SCI increases by age until 15 19 years where it remains reasonably constant until 85years of age Prevalence and YLDs showed similar patterns with one peak at around 45 to 54 years (Figure 4).

#### Conclusions

While age standardized rates of incidence, prevalence, and YLDs for SCI changed slightly, absolute counts increased substantially from 1990 there is also a wide geographical heterogeneity in demographic, spatial and temporal patterns of SCI both at national and regionallevel, which should be considered by policy makers to reduce the burden of SCI locally

- 1. Hartkopp A, Brønnum Hansen H, Seidenschnur AM, Biering Sørensen F Survival and cause of death after traumatic spinal cord injury A long term epidemiological survey from Denmark Spinal cord 1997 35 2 76 85
- 2. Thietje R, Pouw MH, Schulz AP, Kienast B, Hirschfeld S Mortality in patients with traumatic spinal cord injury descriptive analysis of 62 deceased subjects The journal of spinal cord medicine 2011 34 5 482 7 3
- 3. Van den Berg ME, Castellote JM, de Pedro Cuesta J, Mahillo Fernandez I Survival after spinal cord injury a systematic review Journal of neurotrauma 2010 27 8 1517 28

# VALIDATION OF THE ARTHROPLASTY REGISTER OF SOUTH TYROL BETWEEN 2010-2020 – DATA ANALYSIS OF THE SABES HOSPITALS

Marco Santarelli<sup>1</sup>, Roberto Picus<sup>2</sup>, Michael Engl<sup>1</sup>

<sup>1</sup>Dept. Orthopaedic - Sabes Hospital Holding - Province of Bolzano, South Tyrol, Italy;

<sup>2</sup>Arthroplasty Register of Autonomous Province of Bolzano (Italy), Observatory for Health – Provincial Government South Tyrol;

Contact: marco.santarelli@stud.pmu.ac.at

#### Objective

To validate an Arthroplasty Register (AR) means to proof its robustness in producing complete and correct data. Existing literature showed missing revision rates from 12% to 20% and data entry error of 2%. The AR of South Tyrol, nested in the Italian one, began in year 2010. Its completeness was 98% for hips and 100% for the knees. Every year reports are published, but correctness of those data has to be checked. Aim of this work is to discuss the validation project

#### Methods

Retrospective analysis of register data of hip and knee arthroplasty between 2010 and 2020 has been performed (around 20,000 procedures, in seven public hospitals). 5% of primary and 10% of revision procedures were checked. Missing records checked as well. Particular attention has been made by registered aseptic loosening procedures.

#### Results

We found a general data entry error rate of 2 to 4%. Higher rates were related to revision procedures. Data entry errors were divided in tipping- and syntax ones, with a clear reduction during time, due to learning curve. False coding for aseptical revisions was about 7% of all loosening revisions. (2% of all revision cases). This was due to lacking correction of the data already entered in register, after receiving positive antibiogram. Missing cases are 3% for primary procedure and 14% for revision ones.

#### Conclusions

The resulted error rate after 10-year arthroplasty register in South Tyrol is aligned with previous validated national register around the world. Further correction steps needed to be undertaken to reduce overall error in all categories. Therefore, SABES implemented a novel internal AR-database with open-access to every competent authority. Preventive medicine must rely on complete and corrected register data.

- Arthursson, Astvaldur & Furnes, Ove & Espehaug, Birgitte & Havelin, Leif & Söreide, Jon. (2005). Validation of data in the Norwegian Arthroplasty Register and the Norwegian Patient Register: 5,134 Primary total hip arthroplasties and revisions operated at a single hospital between 1987 and 2003. Acta orthopaedica. 76. 823-8. 10.1080/17453670510045435.
- Pugely, A. J., Martin, C. T., Harwood, J., Ong, K. L., Bozic, K. J., & Callaghan, J. J. (2014). Database and registry research in orthopaedic surgery: Part 2: Clinical registry data. Journal of Bone and Joint Surgery - American Volume, 97(21), 1799–1808. https://doi.org/10.2106/JBJS.O.00134

## How primary care physicians deal with the climate crisis: preliminary results of a scoping review

#### Johanna Schauer-Berg<sup>1</sup>, Andrea Stitzel<sup>2</sup>

<sup>1</sup> Institute of General Practice, Family Medicine and Preventive Medicine, Paracelsus Medical University, Salzburg, Austria <sup>2</sup> Department of Health Sciences and Social Work, Carinthia University of Applied Sciences, Villach, Austria Contact: Johanna.Schauer@pmu.ac.at

#### Objective

In 2009 a LANCET commissions called the climate crisis "the biggest threat for health in the 21st century". Physicians in primary care are increasingly confronted with its direct and indirect health impacts e.g. on infectious diseases, allergies, mental health and heat-related diseases. At the same time healthcare contributes significantly to greenhouse gas emissions - in Austria: 7% of national carbon emissions.<sup>1</sup> A transformation of patient care is needed in order to reduce its footprint (climate mitigation) and to provide a proactive response to the unavoidable climate-health consequences (climate adaptation).

#### Methods

A scoping review is performed according to the Joanna Briggs Institute (JBI) approach<sup>2</sup> to map available evidence on climate change mitigation and adaptation in primary care. Reference search was conducted via PubMed, Medline and Web of Science and a hand search applied.

References were reviewed by an interdisciplinary research team. Only references reporting interventions from EU countries the UK, Norway and Switzerland, published between January 2012 and May 2023 were included. With regard to the sources, an open format was chosen. Reporting is done according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR)<sup>3</sup> scheme.

#### Results

As this is an ongoing research project only preliminary results can be presented. 2.769 references were screened. Only six references included for final analysis were relevant to primary care physicians. All of those articles covered aspects of climate mitigation (travel&transport (5), pharmaceuticals (2), circular healthcare & waste management (2), electricity (2), greater health system effectiveness (2) food/nutrition (1) and buildings & Infrastructure (1)) but not climate adaptation.

#### Conclusion

Preliminary results show that there is very little evidence on actual implementation of climate mitigation and adaptation in primary care practices esp. with regard to validation of effectiveness of measures. However, the provision of sound evidence-based guidance is essential in order to increase the climate resilience of patient care and thus the population. Therefore, in depth research is urgently needed.

- 1. Weisz U, Pichler PR, Jaccard IS, Haas W, Matej S, Nowak P, Bachner F, Lepuschütz L, Windsperger A, Windsperger B, Weisz H. Der Carbon Fußabdruck des österreichischen Gesundheitssektors. Endbericht. Klima- und Energiefonds, Austrian Climate Research Programme. 2019
- 2. Erik von Elm, Gerhard Schreiber, Claudia Cornelia Haupt. Methodische Anleitung für Scoping Reviews (JBI-Methodologie). Z Evid Fortbild Qual Gesundhwes. 2019 Jun;143:1-7. doi: https://doi.org/10.1016/j.zefq.2019.05.004
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ, Horsley T, Weeks L, Hempel S, Akl EA, Chang C, McGowan J, Stewart L, Hartling L, Aldcroft A, Wilson MG, Garritty C, Lewin S, Godfrey CM, Macdonald MT, Langlois EV, Soares-Weiser K, Moriarty J, Clifford T, Tunçalp Ö, Straus SE. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. Ann Intern Med. 2018 Oct 2;169(7):467-473. doi: 10.7326/M18-0850. Epub 2018 Sep 4. PMID: 30178033.

#### SEX DIFFERENCES IN PAIN COURSES IN PATIENTS WITH HIP OR KNEE OSTEOARTHRITIS AFTER MULTIMODAL SPA THERAPY IN THE AUSTRIAN VALLEY OF GASTEIN

<u>Moritz Schröter</u><sup>1,2</sup>, Kathrin Bogensberger<sup>1,2,3</sup>, Julia Fuchs<sup>1,2,3</sup>, Sonja Wildburger<sup>1,2,3</sup>, Markus Ritter<sup>1,2,3,4</sup>, Antje van der Zee-Neuen<sup>1,2,3,5</sup>

<sup>1</sup>Institute for Physiology and Pathophysiology, Center for Physiology, Pathophysiology and Biophysics, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Gastein Research Institute, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Ludwig Boltzmann Institute for Arthritis and Rehabilitation, Vienna, Austria; <sup>4</sup>Kathmandu University School of Medical Sciences, Dhulikhel, Nepal; <sup>5</sup>Center for Public Health and Health Care Research, Paracelsus Medical University, Salzburg, Austria; Austria; Contact: moritz.schroeter@stud.pmu.ac.at

#### Objective

Sex-differences in perceived pain have been discussed widely throughout the literature pointing, among others, towards the role of intervention type in the association of sex and perceived pain. Therefore, pain outcomes after interventions should be evaluated separately for men and women enabling determination of adequate treatment intervals and sex-specific modulation of interventions. The current study aimed to shed light on sex-differences in the course of pain in patients with hip or kneeosteoarthritis (OA) after multimodal spa-therapy in the Austrian valley of Gastein.

#### Methods

Longitudinal analyses of prospectively collected registry data from the ongoing "Radon indication registry" collecting data directly before (baseline), directly after and 3; 6 and 9 months after multimodal spa-therapy were conducted. Multilevel mixed-effects linear regression models were computed to explore the association of timepoint of measurement with a) pain in motion and b) pain in rest (both measured on an 11-point numeric rating scales; 0=no pain, 10=worst pain imaginable) separately for men and women while adjusting for age and body mass index (BMI) and accounting for individual differences through inclusion of patient identification number as random effect parameter.

#### Results

The sample consisted of 34 women and 52 men with a mean age of 59 (SD 6.4) & 58 (SD 7.9) years and a mean BMI of 26.0 (SD 4.3) & 28.3 (SD 3.4), respectively. Both sexes showed significant improvements in pain in rest and motion until up to 9 months after treatment when compared to baseline. However, improvements were larger and in case of pain in rest sustained longer in women than in men.

#### Conclusions

These results point towards the relevance of individualized treatment intervals while accounting for sex and might give rise to treatment plan modulation to support sustained treatment results specifically for men with OA who receive multimodal spa-therapy in the Austrian Gastein valley. They also support previous findings that emphasize the necessity of sex specific intervention evaluation and development.

#### Comparison of naturally produced EVs and artificial CDVs

Pauline Ley<sup>1</sup>, <u>Negar Sedghi Aminabad</u><sup>1</sup>, Antri Stefani<sup>1</sup>, Martin Wolf<sup>1</sup>, Thomas Heuser<sup>2</sup>, Nicole Maeding<sup>1</sup>, Sarah Hochmann<sup>1</sup>, Dirk Strunk<sup>1</sup>

<sup>1</sup>Institute of Experimental and Clinical Cell Therapy, Paracelsus Medical University (PMU), Salzburg, Austria; <sup>2</sup>Vienna Biocentre Core Facilities, Vienna, Austria; Contact: negar.sedghi@pmu.ac.at

#### Objective

In the development of a Parkinson's Disease (PD) vaccine, a biomarker to identify patients at risk is crucial. Currently, extracellular vesicles (EVs) stand out as promising candidates. This demands the production of model vesicles as reference material in large quantities. Isolation methods such as TFF are time-intensive, requiring weeks to collect sufficient cell supernatant due to a limited number of 100-1000 EVs released by cells daily. In comparison, an extrusion could significantly speed up the process. Here, cells are pressed through an extruder membrane, which produces a high number of extracellular particles, as depicted in figure 1. Therefore, this experiment sets out to analyse the characteristics of such CDVs to determine if they have comparable properties to secreted EVs. Due to its relevance for PD research, this experiment focused on one fluorescent neuroblastoma cell line (SH-SY5Y GFP) which expresses GFP-labelled  $\alpha$ -syn, a key protein for PD development and progression. Two control lines HEK CD63 mNEON and HEK-WT were included for comparison.

#### Methods

CDVs were extruded from three cell lines (SH-SY5Y GFP, HEK CD63 mNEON, HEK-WT) comparing two extruder models (Extr. a / Extr. b). Resulting particles were imaged through cryo-transmission electron microscopy (cryo-TEM), counted by tunable resistive pulse sensing (TRPS) and further evaluated by fluorescence nanoparticle tracking analysis (f-NTA). Dot blots confirmed the EV identity by tetraspanin expression and presence of  $\alpha$ -syn. TFF-EVs served as comparison material.

#### Results

Characterisation of extruded CDVs: Three different cell lines were used (SH-SY5Y, HEK CD63 mNEON, HEK-WT) to produce CDVs comparing two extruders (Extr. a / Extr. b). Graphs are formed from the mean of the results of three extrusions. For all experiments no difference between the two extruder models could be detected. It could be confirm by Cryo-TEM that the produced particles were EVs and CDVs with a double membrane.Particle count: Size distribution and concentration of CDVs and EVs, both measured by TRPS. CDVs from all three cell lines show a similar size distribution with a mean size of about 120 nm (mode size 90 nm). This is slightly bigger than reported values of naturally produced EVs from HEK cells (mean size 90 nm) (Görgens et al., 2022). The CDVs were concentrated at 1011 particles/mL which is slightly higher than TFF-EVs (at 1010).EV identity: Dot blots of tetraspanin and  $\alpha$ -syn at 1 second exposure time confirmed the expression of all three tested tetraspanins (CD9, CD63, CD81) in all cell lines and the expression of  $\alpha$ -syn in SH-SY5Y.Fluoresence Analysis: Fluorescence positivity of CDVs measured by f-NTA. The fluorescence analysis via f-NTA showed that CDVs were less fluorescent than EVs. While approximately 80% of the HEK CD63 mNEON EVs were fluorescent, only 10% of the CDVs expressed mNEON. SH-SY5Y-EVs and -CDV fluorescence was <1%. (g) Double-positive HEK CD63 mNEON EVs and CDVs for CD63 and mNEON measured by f-NTA. In the HEK CD63 mNEON samples, mNEON was expressed by 60% of the CD63+ EVs and 20% of the CD63+ CDVs.

#### Conclusions

Extruded CDVs have a high concentration and show similarities with EVs in terms of size. They also express tetraspanins and in case of the SH-SY5Y cells  $\alpha$ -syn. However, there seem to be major losses of fluorescence during cell extrusion, as CDVs were 10-fold less fluorescent than EVs. To better understand the mechanisms behind cell extrusion, we could conduct additional experiments to test extrusion parameters such as membrane pore size, stabilizers, cell numbers, temperature etc. Additional filtration and concentration steps could be added to further improve the quality of the samples.

- 1. Le QV, Lee J, Lee H, Shim G, Oh YK. Cell membrane-derived vesicles for delivery of therapeutic agents. Acta Pharmaceutica Sinica B. 2021 Aug 1;11(8):2096-113
- 2. Wen Y, Fu Q, Soliwoda A, Zhang S, Zheng M, Mao W, Wan Y. Cell-derived nanovesicles prepared by membrane extrusion are good substitutes for natural extracellular vesicles. Extracellular vesicle. 2022 Dec 1;1:100004.
- Alter CL, Detampel P, Schefer RB, Lotter C, Hauswirth P, Puligilla RD, Weibel VJ, Schenk SH, Heusermann W, Schürz M, Meisner-Kober N. High efficiency preparation of monodisperse plasma membrane derived extracellular vesicles for therapeutic applications. Communications biology. 2023 May 3;6(1):478.

#### Physicians' Perspectives, Well-being, and Fulfilment in Telemedicine

#### Cristina Sicorschi Gutu<sup>1,2</sup>, Krisztina Schmitz-Grosz<sup>1</sup>, Carsten Sommer<sup>1</sup>, Günter Niklewski<sup>2</sup>

<sup>1</sup>Medgate AG, Basel, Switzerland;

<sup>2</sup>University Clinic for Psychiatry and Psychotherapy, Paracelsus Medical University, Nuremberg, Germany.; Contact: crsicorschi@gmail.com

#### Objective

Telemedicine, a safe way of medical care appreciated by patients, has proven to have different benefits for our healthcare systems. Studies on patient experiences and outcomes in telemedicine are thriving. However, few studies have explored the possible associations between telemedicine and physician work-life balance or mental health.

#### Methods

We conducted a cross-sectional survey for physicians who practiced in Switzerland. Physicians who participated answered validated questions about work-life balance, professional satisfaction, emotional exhaustion, and depression. There were three groups: a control group with physicians who did not practice telemedicine (n=30), an experimental group with physicians who practice telemedicine (n=32), and a third group with physicians who left telemedicine (n=31). The 42-question survey was open between June and August of 2023. Participation was optional, anonymous, and voluntary.

#### Results

In total, 93 participants completed the questionnaire. In terms of work-life balance, the best results were observed in the telemedicine group (4.63 vs 4.35 in the control group). A high level of exhaustion was observed in 26.7% of the control group, compared to 12.5 % of the telemedicine group. Low professional fulfilment was present in 62.5% of participants in the telemedicine group vs. 46.7 % in the control group. Among the telemedicine physicians, 21.9% were suspected of having depression.

#### Conclusions

Practicing telemedicine as a profession may have a positive impact on physicians' work-life balance and burnout. We found no significant statistical differences among groups regarding professional fulfillment and depression. Further research is necessary to explore long-term effects of telemedicine on physicians' mental health.

- Zhang A, Wang J, Wan X, Zhang Z, Zhao S, Guo Z, Wang C. A Meta-Analysis of the Effectiveness of Telemedicine in Glycemic Management among Patients with Type 2 Diabetes in Primary Care. Int J Environ Res Public Health. 2022 Mar 31;19(7):4173. doi: 10.3390/ijerph19074173. PMID: 35409853; PMCID: PMC8999008.
- Choi JS, Kim JH, Park S, Lin M, Abdur-Rahman F, Mack WJ, Volker CCJ. Telemedicine in Otolaryngology During COVID-19: Patient and Physician Satisfaction. Otolaryngol Head Neck Surg. 2022 Jul;167(1):56-64. doi: 10.1177/01945998211041921. Epub 2021 Sep 7. PMID: 34491856.

#### Characterisation of variants in the SLC26A4 gene identified in a Slovak cohort with nonsyndromic enlarged vestibular aqueduct and Pendred syndrome

<u>Marek Sklenár</u><sup>1</sup>, Emanuele Bernardinelli<sup>2</sup>, Silvia Borecká<sup>1</sup>, Lukáš Varga<sup>1,3</sup>, Diana Ugorová<sup>3</sup>, Daniela Gašperíková<sup>1</sup>, Silvia Dossena<sup>2</sup>

<sup>1</sup>Institute of Experimental Endocrinology, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovakia; <sup>2</sup>Institute of Pharmacology and Toxicology and FIZ - RM & NT, Paracelsus Medical University, Salzburg, Austria; <sup>3</sup>Department of Otorhinolaryngology-Head and Neck Surgery, Faculty of Medicine and University Hospital, Comenius University Bratislava, Bratislava, Slovakia;

Contact: marek.sklenar@savba.sk

#### Objective

Pathogenic variants in the SLC26A4 gene, encoding for CI-/HCO3- and I- anion transporter pendrin, are one of the most common causative factors of hereditary hearing loss worldwide and can be associated with non-syndromic hearing loss with enlarged vestibular aqueduct (NSEVA) or Pendred syndrome (PDS), manifested by thyroid dysfunction and goiter (1). The aim of this study was to characterize the functional and molecular features of five pendrin variants identified in Slovak patients with sensorineural hearing loss (NSEVA or PDS) and assess a correlation with the patient's clinical symptoms.

#### Methods

Pendrin variants were introduced into SLC26A4-encoding mammalian expression vectors by site-directed mutagenesis and expressed in HEK293 Phoenix cells and HeLa cells. The ion transport efficiency of pendrin variants was determined by fluorometric analysis by using the iodide-sensitive EYFP variant H148Q;I152L to monitor iodide influx. Quantitative confocal imaging was performed to determine the total expression level of pendrin variants. The subcellular localisation of pendrin variants was determined by analysing the co-localisation with plasma membrane (PM) or endoplasmic reticulum (ER) markers.

#### Results

Functional tests showed that five out of six studied pendrin variants (p.G139R, p.M147I, p.Y530S, p.D754IfsX5, and p.F161I) exhibit significantly reduced function compared to wild-type pendrin, while the function of the variant p.R47Q was preserved. By quantitative confocal microscopy, we confirmed a massively reduced expression of pendrin variants p.G139R, p.M147I, p.Y530S, and p.F161I compared to wild-type. The p.R47Q variant showed a wild-type-like expression. Moreover, variants p.G139R and p.Y530S showed retention in the ER in comparison to the wild-type protein and severely diminished co-localization with the PM. Variants p.M147I and p.F161I showed moderate, but significant retention in ER compared to wild-type pendrin and significantly reduced PM targeting. Variant p.R47Q was identical to the wild-type.

#### Conclusions

Our results demonstrated reduced function and impaired subcellular localization of variants p.G139R, p.M147I, p.Y530S, p.D754IfsX5, and p.F161I. The experiments confirmed the potentially pathogenic character of these five variants and their association with symptoms of NSEVA/PDS in affected patients. On the other hand, the variant of unknown significance p.R47Q was reclassified as benign and its role in the pathogenesis of mild hearing loss has not been confirmed.

#### Acknowledgements

Supported by grants APVV-20-0236, VEGA 1/0572/21

#### References

1. Khan MR, Bashir R, Naz S. SLC26A4 mutations in patients with moderate to severe hearing loss. Biochem Genet. 2013 Aug;51(7-8):514-23. PMID: 23504402.

# Prognostic and predictive factors (clinical data, histological and molecular data, laboratory values) under palliative first-line therapy for squamous cell carcinoma of the lung with chemotherapy plus immunotherapy

Johanna Sodmann<sup>1</sup>, Dieter Würflein<sup>1</sup>, Fabian Reich<sup>1</sup>, Wolfgang Brückl<sup>1</sup>, Joachim Ficker

<sup>1</sup>Department of Respiratory Medicine, Allergology and Sleep Medicine / Nuremberg Lung Cancer Center, Paracelsus Medical University, Nuremberg, Germany; Contact: johanna.sodmann@stud.pmu.ac.at

#### Objective

Patients with squamous cell carcinoma of the lung have low 5-year survival rates (2). This study examines various prognostic and predictive factors (clinical data and blood values) in patients with chemotherapy plus immune checkpoint inhibitors in palliative first-line therapy.

#### Methods

Data collected from 61 patients (2018 to 2023) were analyzed retrospectively. All Patients suffered on a squamous cell lung cancerg and received a combination of at least one cycle of the ICI pembrolizumab and chemotherapy (carboplatin and (nab-)paclitaxel) as part of palliative therapy. Significant parameters were identified using log-rank test and chi-square test: Cox regression was applied.

#### Results

The 61 patients examined had a median age of 65 years and 72.1% were male. The median overall survival was 19.9 months and the median progression-free interval was 9.9 months. In the multivariate analysis, age  $\geq$  65 years (p=0.001) and a body mass index value  $\leq$  25 (p=0.01) were significant independent prognostic parameters for longer median overall survival. Age  $\geq$  65 years (p=0.007) and PD-L1 status > 0 (p=0.039) are significant for progression-free survival. An increased CRP value before the second cycle ("dynamic change") showed predictive relevance for a poorer prognosis in the univariate analysis. A PD-L1 status > 0 (p = 0.016) and normal G-GT value before the start of immunochemotherapy (p = 0.008) are positive predictors of treatment response.

#### Conclusions

Clinical and laboratory parameters can give an outlook on the response and course of therapy of patients with squamous cell carcinoma of the lung with immunochemotherapy.

- 1. Derman BA, Mileham KF, Bonomi PD, Batus M, Fidler MJ. Treatment of advanced squamous cell carcinoma of the lung: A review. Transl Lung Cancer Res. 2015;4(5). doi:10.3978/j.issn.2218-6751.2015.06.07
- 2. Santos ES, Rodriguez E. Treatment Considerations for Patients With Advanced Squamous Cell Carcinoma of the Lung. Clin Lung Cancer. 2022;23(6). doi:10.1016/j.cllc.2022.06.002
- Novello S, Kowalski DM, Luft A, et al. Pembrolizumab Plus Chemotherapy in Squamous Non-Small-Cell Lung Cancer: 5-Year Update of the Phase III KEYNOTE-407 Study. Journal of Clinical Oncology. 2023;41(11). doi:10.1200/JCO.22.01990

#### Metabolic Footprint of Ketosis-inducing Diets in Mice

Julia Tevini<sup>1</sup>, Daniela D Weber<sup>1</sup>, <u>Victoria E Stefan<sup>1,2</sup></u>, Thomas K Felder<sup>3,4</sup>, Rohit Dnyansagar<sup>1</sup>, Barbara Kofler<sup>1</sup>, Kalina Duszka<sup>5</sup>

<sup>1</sup>Research Program for Receptor Biochemistry and Tumor Metabolism, Department of Pediatrics, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Biosciences and Medical Biology, University of Salzburg, Austria; <sup>3</sup>Department of Laboratory Medicine, Paracelsus Medical University, Salzburg, Austria; <sup>4</sup>Institute of Pharmacy, Paracelsus Medical University, Salzburg, Austria; <sup>5</sup>Department of Nutritional Sciences, University of Vienna, Vienna, Austria; Contact: v.stefan@salk.at

#### Objective

Emerging evidence associates diet-induced ketosis with several beneficial effects on health and disease, inter alia reduced inflammation, improved lipid profiles, weight loss, and improved diabetes management (1). However, little is known about the effect of different ketosis-inducing dietary approaches on the metabolome. Therefore, we aimed to characterize the metabolic profile of multiple matrices, namely plasma, liver, spleen, white adipose tissue, duodenum, and feces of mice fed with different ketosis-inducing diets, including caloric restriction (CR), intermittent fasting (IF), fasting-mimicking diet (FMD), ketogenic diet (KD), and regular mouse chow.

#### Methods

Male C57BL/6 mice were subjected to the following dietary regimes: CR (14 days 80% kcal of regular ad libitum chow intake), IF (28 days 100% food restriction every other day followed by 24h ad libitum refeeding), FMD (28 days, 3 cycles of 4 days fasting with 50% kcal of regular ad libitum chow intake on day 1, 10% kcal on days 2-4 with 7 days of refeeding), KD (28 days ad libitum ketogenic ratio 4:1), and regular chow (28 days ad libitum) (2). Samples were collected at a mouse age of 16 weeks for all dietary intervention groups. A comprehensive metabolomics analysis using the MxP® Quant 500 Kit (biocrates life sciences AG) enabled the quantification of 26 compound classes, such as amino acids, amino acid-related metabolites, bile acids, free fatty acids, cholesterol esters, acylcarnitines, sphingolipids, glycerophosphocholines, and glycerolipids.

#### Results

In the dietary groups, lipid and polar metabolite classes varied substantially in plasma and tissues. For instance, CR, IF, FMD, and KD reduced glycerolipid levels (di- and triacylglycerides) in plasma by 38%, 56%, 90%, and 49%, respectively, compared to regular chow. Moreover, two metabolites, namely proline betaine and trigonelline, were consistently reduced in plasma and in all organs, while 3-methylhistidine was consistently increased in all matrices except feces by the ketosis-inducing diets.

#### Conclusions

Our analysis provides a comparable multi-tissue metabolic footprint for each of the diets as a basis for understanding the various effects attributed to ketosis-inducing diets on a metabolic level. Further analysis will reveal the association of specific metabolic changes with diet-induced ketosis and its health benefits (e.g. association with inflammatory markers).

- 1. Puchalska P, Crawford PA. Metabolic and Signaling Roles of Ketone Bodies in Health and Disease. Annu Rev Nutr. 2021 Oct 11;41:49-77. doi: 10.1146/annurev-nutr-111120-111518. PMID: 34633859; PMCID: PMC8922216.
- Gregor A, Huber L, Auernigg-Haselmaier S, Sternberg F, Billerhart M, Dunkel A, Somoza V, Ogris M, Kofler B, Longo VD, König J, Duszka K. A Comparison of the Impact of Restrictive Diets on the Gastrointestinal Tract of Mice. Nutrients. 2022 Jul 29;14(15):3120. doi: 10.3390/nu14153120. PMID: 35956298; PMCID: PMC9370610.

#### Theranostic EV biomarkers from non-invasive liquid biopsies

<u>Antri Stefani</u><sup>1</sup>, Martin Wolf<sup>1</sup>, Sarah Hochmann<sup>1</sup>, Nicole Maeding<sup>1</sup>, Anna Raninger<sup>1</sup>, Anna Steinhuber<sup>1</sup>, Rodolphe Poupardin<sup>1</sup>, Dirk Strunk<sup>1</sup>

<sup>1</sup>Cell Therapy Institute, Paracelsus Medical University, Salzburg, Austria; Contact: antri.stefani@pmu.ac.at

#### Objective

Extracellular vesicles (EVs) possess key roles in cell-to-cell communication in health and disease (1). Their presence in most tissues and body fluids makes them an excellent target for biomarker discovery. Here we are developing a theranostic Parkinson's disease (PD) biomarker to select candidate patients for vaccination and monitor their response. We selected, PD driving oligomeric  $\alpha$ -synuclein ( $\alpha$ -syn) as primary diagnostic target. Not knowing the location of  $\alpha$ -syn inside EVs or as surface cargo, we apply methods preserving natural EV corona enabling extended downstream biomarker analysis.

#### Methods

EVs were enriched from healthy donor urine (uEV) via tangential flow filtration (TFF). Influence of storage at - 80°C on EV quantity and size was analysed by tunable resistive pulse sensing (TRPS), and quality by tetraspanin+ biomarker expression in dot plots and super-resolution microscopy

#### Results

Small uEVs with a mode size 89.75±0.59 nm (mean±SEM) were isolated by TFF. EV CD9/63/81 tetraspanin expression did not show significant differences before/after cryo-storage despite 63-86% loss of particle number from 3.0±0.5x1010 to 1.6±1.0x1010 particles/mL (mean±SEM) after thawing. Spiking a-syn-EVs as a representative biomarker into uEVs enabled comparing both sources for setting up a sensitive detection assay. Titration of a-Syn-EVs will determine optimum pre-analytic conditions for subsequent biomarker detection based on ISEV uEV task force recommendations.

#### Conclusions

Developing new theranostic biomarker analysis methods from easy accessible non-invasive sources is important for patient acceptance and study performance. We develop a standardized protocol regarding sample retrieval, pre-processing, storage and EV isolation for biomarker studies

#### References

1. Pupardin et al., 2023, PMID: 38270559

#### Nanoindentation on Tendon Tissue

Herbert Tempfer<sup>1,2</sup>, Thomas Lettner<sup>1,2</sup>, Bettina Faustini<sup>1,2</sup>, Andreas Traweger<sup>1,2</sup>

<sup>1</sup>Institute of Tendon and Bone Regeneration, Paracelsus Madical University, Salzburg, Austria; <sup>2</sup>Austrian Cluster for Tissue Regeneration, Vienna, Austria; Contact: herbert.tempfer@pmu.ac.at

#### Objective

Biomechanical testing of tendons is commonly performed by uniaxial tensile stretching along collagen fibre orientation. So far, only little is known about tendon biomechanical response to compression on a nanoscale, which may provide information on tissue quality not achievable by tensile testing.

#### Methods

Cryosections of unfixed healthy rat patella tendons derived from male SD rats and from tendons 2, 4 and 6 weeks after creating a 2 mm punch defect were examined with a Chiaro Nanoindenter (Optics11 B.V.). Measurements were performed at several time points after immersion, either in Phosphate Buffered Saline (PBS) or 8% polyethylene glycole (PEG) in PBS. For indentation, a cantilever with a stiffness of 0.5 N/m and a sphere tip with a radius of 52 µm was used. Stiffness was calculated using the Hertzian model.

#### Results

In healthy tendons, E(eff) declined from  $23.4\pm3.6$  kPa after 5 minutes to  $11.3\pm2.9$  kPa after 60 minutes, stiffness of the tendon defect area only declined from  $4.9\pm1.4$  kPa after 5 Minutes to  $3.6\pm1.1$  kPa after 60 minutes (Figure 1). Immersion in 8% PEG after 5 minutes hydration in PBS lead to an average stiffness of  $135\pm34.1$  kPa after 10 minutes.

#### Conclusions

The observed decline in elasiticity modulus due to PBS immersion needs to be considered for longer lasting measurements like surface mapping studies. Moreover, the use of PEG to avoid tissue swelling as it has been shown for tensile testing (1) of tendons is not suitable for tendon nanoindentation. As nanoindentation measurements can only be performed in aqueous solutions, the "in vivo degree of hydration" remains an unknown variable, favouring semiquantitative, whithin- experiment analyses.

#### References

 Safa BN, Meadows KD, Szczesny SE, Elliott DM. Exposure to buffer solution alters tendon hydration and mechanics. J Biomech. 2017 Aug 16;61:18-25. doi: 10.1016/j.jbiomech.2017.06.045. Epub 2017 Jul 6. PMID: 28720201; PMCID: PMC5659756.

#### The clinical relevance of low-grade infected nonunion

<u>Katharina Trenkwalder<sup>1,2</sup></u>, Sandra Erichsen<sup>1,2</sup>, Ferdinand Weisemann<sup>3</sup>, Peter Augat<sup>1,2</sup>, Simon Hackl<sup>2,3</sup>

<sup>1</sup>Institute for Biomechanics, BG Unfallklinik Murnau, Germany;

<sup>2</sup>Institute for Biomechanics, Paracelsus Medical University, Salzburg, Austria;

<sup>3</sup>Department of Trauma Surgery, BG Unfallklinik Murnau, Germany; Contact: katharina.trenkwalder@bgu-murnau.de

#### Objective

Treatment algorithms for fracture nonunion depend on the presence or absence of a bacterial infection. However, the clinical presentation of nonunion varies and an underlying infection is not always obvious. While patients with manifest infection present with clinical signs of infection, patients with low-grade infected nonunion present similarly to patients with aseptic nonunion. Only microbiology and histopathology of intraoperative obtained samples can reveal a low-grade infection. The aim of this study was to evaluate the clinical relevance of low-grade infection in the development and management of nonunion.

#### Methods

A prospective, multicenter clinical study enrolled patients with femoral or tibial shaft nonunion and regular healed fractures. Preoperatively, serum markers including C-reactive protein (CRP), leukocytes, and procalcitonin were determined, clinical infection signs were recorded, and a suspected 'septic' or 'aseptic' diagnosis was made. During nonunion revision or routine implant removal, tissue samples were collected for microbiology and histopathology, and osteosynthesis material for sonication. Nonunion patients were followed for 12 months. Definitive diagnosis of 'septic' or 'aseptic' nonunion was made according to diagnostic criteria for fracture-related infection (1), considering the follow-up period.

#### Results

110 patients with nonunion and 34 patients with regularly healed fractures were included. 27 nonunions were suspected to have an underlying infection, which was confirmed in 22 cases. 83 nonunions were assumed to be aseptic, but an underlying low-grade infection was diagnosed in 23 cases. Three cases with an unclear diagnosis were excluded from statistical analyses. Sensitivity and specificity for the suspected diagnosis were 49% (34-64% CI) and 95% (86-99% CI), respectively. The low-grade infection rate in presumed aseptic nonunion was 28%. CRP was the only serum blood marker with mean values above the reference range that was significantly higher in all nonunion groups compared with the regular healer group. Manifest and low-grade infected nonunions were commonly caused by coagulase-negative staphylococci (59% and 74%) and Cutibacterium acnes (23% and 52%). The latter was detected significantly more often in low-grade infections compared to manifest infections (p=0.042). Patients with low-grade infections were treated with systemic antibiotics significantly less frequently (55% vs. 96%, p=0.002) and for a significantly shorter duration than patients with manifest infections (6±3 vs. 11±5 weeks, p=0.005). Eight nonunions in the low-grade infection group consolidated during follow-up period without systemic antibiotics. The overall healing rate of low-grade infected nonunion (83%) was statistically comparable to that of manifest infected nonunion (62%) and aseptic nonunion (90%).

#### Conclusions

Our findings demonstrate a significant role for low-grade infection in the development of nonunion. The only difference between aseptic and low-grade infected nonunion was the intraoperative detection of bacteria and the associated need for systemic antibiotics. Combined with an effective surgical revision, a high healing rate could be achieved for low-grade infected nonunion.

#### References

1. Metsemakers WJ, Morgenstern M, McNally MA, Moriarty TF, McFadyen I, Scarborough M, et al. Fracture-related infection: A consensus on definition from an international expert group. Injury. 2018;49(3):505-10.

#### Bioanalytic methods in psychosomatic medicine

#### Sophia Urbanczyk

<sup>1</sup>Department/Institute of Psychosomatic Medicine, Paracelsus Medical University, Nuremberg, Germany; Contact: sophia.urbanczyk@klinikum-nuernberg.de

#### Objective

Psychological and emotional stress triggers a cascade of physiological responses, including the activation of the hypothalamic-pituitary-adrenal (HPA) axis and the sympathetic-adrenal-medullary (SAM) axis. These processes can lead, among other things, to the increased production of reactive oxygen species (ROS). Chronic states of increased ROS levels are referred to as "oxidative stress" and are characterized by lipid, protein and DNA damage. Oxidative stress contributes to a myriad of pathologies, such as cancer, autoimmunity, cardiovascular diseases and neurological disorders.1)2)

#### Methods

Quantification of ROS levels: Electron Paramagnetic Resonance (EPR)EPR is a spectroscopic method based on the interaction between unpaired electrons and an external magnetic field. For the detection of ROS special membrane-permeabel spinprobes are employed which produce less reactive and more stable species that can be readily detected by EPR.2)Quantification of DNA damage: Single-cell electrophoresis ("Comet Assay")In the comet assay, cells are embedded in agarose on a microscope slide, lysed to remove cellular proteins, and subjected to alkaline denaturation with subsequent electrophoresis. During electrophoresis, DNA fragments migrate away from the nucleus towards the anode according to their size, forming a comet-like tail. The quantification of DNA damage is achieved through analysis of the tail moment, which is the product of the tail length and the fraction of DNA within the tail.Quantification of mitochondrial oxygen consumption rates: Oxygraph O2K "Oroboros"The Oroboros oxygraph operates on the principle of high-precision polarography, where the oxygen concentration is detected amperometrically by a Clark-type electrode. The application of "SUIT" (substate, uncoupler, inhibitor, titration) protocols allow the real-time monitoring of oxygen flux in various experimental conditions, thereby enabling the analysis of mitochondrial respiration and function of cells and organs.

#### Results

Preliminary data from the MALT study, which analysed the effectiveness of art therapy after a misscarriage or stillbirth, showed significantly increased levels of ROS in the blood and slightly but not significantly increased tail moment, indicative of DNA damage, in the study group compared to the control group. The participants were analysed under resting conditions (T-10), experimental stress (T0) and after regeneration (T+20). A Social Evaluated Cold Pressor Test (SECPT) was employed to induce a reliable stress response.

#### Conclusions

Finding biological marker for psychological stress may represent promising tools for future research and treatment.

#### Acknowledgements

Luis Gerber, Sofie Roecken, Andrea Dorenkamp, Alexandra Hass

- 1. "Reactive oxygen species: from health to disease", Brieger et al., Swiss Med Wkly. 2012
- 2. "Detection of Reactive Oxygen and Nitrogen Species by Electron Paramagnetic Resonance (EPR) Technique", Suzen et al., Molecules, 2017

### Detecting high risk neighborhoods and socioeconomic determinants for common oral diseases in Germany with geostatistics and machine learning models

<u>Sebastian Völker<sup>1,2</sup></u>, Antje van der Zee-Neuen<sup>3,4</sup>, Alexander Rinnert<sup>5</sup>, Jessica Hanneken<sup>5</sup>, Tim Johansson<sup>2</sup>

<sup>2</sup>Institute of General Practice, Family Medicine and Preventive Medicine, Program Medical Science, Paracelsus Medical University, Salzburg, Austria;

<sup>1</sup>Data Science Center of Excellence, BFS|arvato|Bertelsmann, Dortmund, Germany;

<sup>3</sup>Center for Physiology, Pathophysiology and Biophysics, Institute for Physiology and Pathophysiology/ Gastein Research Institute/Center for Public Health and Healthcare Research, Paracelsus Medical University, Salzburg, Austria;

<sup>4</sup>Ludwig Boltzmann Institute for Arthritis and Rehabilitation, Salzburg, Austria;

<sup>5</sup>Investment & Strategy Management, BFS|arvato|Bertelsmann, Dortmund, Germany;

Contact: sebastian.voelker@stud.pmu.ac.at

#### Objective

Ideally, health services and interventions to improve dental health should be tailored to local target populations. But this is not the standard. Little is known about risk clusters in dental health care and their evaluation based on small-scale, spatial data. To investigate this, i) analyses of the spatial distribution of oral diseases (periodontitis, severe caries, irreversible pulpitis, tooth loss), ii) identification of risk clusters, and iii) the description of these different risk clusters based on socioeconomic determinants are necessary.

#### Methods

A retrospective cohort study was performed to calculate the age- and sex-standardized incidence rate of oral diseases in a study population of privately insured and self-pay patients in Germany who received dental treatment between 2016 and 2021. This was based on anonymized claims data from the BFS health finance|arvato|Bertelsmann (BFS). The disease history of individuals was recorded and aggregated at the ZIP code 5 level (n = 8871). Statistically significant, spatially compact clusters and relative risks (RR) of incidence rates were identified. By linking disease and socioeconomic databases on the ZIP-5 level, local risk models for each disease were estimated based on spatial-neighborhood variables using different machine learning models.

#### Results

We found that dental diseases were spatially clustered among privately insured and self-payer patients in Germany. The relative risks (RR) for a new dental disease in primary risk clusters were 1.3-2.7, depending on the disease. Incidence rates within clusters were significantly elevated compared to incidence rates outside clusters. The methodology presented here enables the identification of disease clusters of greatest demand, which would allow implementing more targeted approaches and improve access to quality care where they can have the most impact.

#### Conclusions

Our results show that private claims data can be used to highlight locations and variables relevant to oral healthcare. A network-based, data-driven approach, that includes the use of non-traditional data sets, holds great potential in supporting resource-based management in the health system. The findings of our study can inform policymakers and researchers in focusing on oral disease incidence and socioeconomic predictors to mitigate disease risk and improve oral health.

- 1. Peres MA, Macpherson LMD, Weyant RJ, Daly B, Venturelli R, Mathur MR, Listl S, Celeste RK, Guarnizo Herreño CC, Kearns C, et al. Oral diseases: a global public health challenge. The Lancet (2019) 394:249 260. doi : 10.1016/S0140 6736(19)31146 8
- 2. Winkelmann J, Rossi JG, van Ginneken E. Oral health care in Europe: Financing, access and provision. Health systems in transition (2022) 24:1 176.
- Jordan AR, Micheelis W. Fünfte Deutsche Mundgesundheitsstudie --(DMS IV). Deutscher Zahnärzte Verlag DÄV Köln (2016)

#### The Impact of Educational Status on the Occurrence of Colonic Diverticula

<u>Andreas Völkerer</u><sup>1</sup>, Sarah Wernly<sup>1</sup>, Georg Semmler<sup>2</sup>, Maria Flamm<sup>3</sup>, Konrad Radzikowski<sup>1</sup>, Leonora Datz<sup>1</sup>, Elmar Aigner<sup>4</sup>, Christian Datz<sup>1</sup>, Bernhard Wernly<sup>3,1</sup>

<sup>1</sup>Department of Internal Medicine, General Hospital Oberndorf, Teaching Hospital of the Paracelsus Medical University, Salzburg, Austria;

<sup>2</sup>Division of Gastroenterology and Hepatology, Department of Medicine III, Medical University of Vienna, Vienna, Austria;
 <sup>3</sup>Institute of general practice, family medicine and preventive medicine, Paracelsus Medical University, Salzburg, Austria;
 <sup>4</sup>University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria;

Contact: a.voelkerer@kh-oberndorf.at

#### Objective

The level of education often serves as an indirect indicator of socioeconomic status.(1) Preliminary investigations have suggested that individuals from lower socioeconomic backgrounds may be more susceptible to diverticulosis.(2) Based on data obtained from the United States, diverticular disease has emerged as a growing health problem with significant financial consequences.(3) Considering these circumstances, our research aimed to investigate the association between educational attainment and the occurrence of colonic diverticula.

#### Methods

We conducted a study involving 5,532 asymptomatic patients in Austria who were undergoing screening colonoscopy. The purpose of our investigation was to examine the connection between educational attainment and diverticulosis. To assess the educational level obtained through the questionnaire, we classified patients into three categories (low, medium, and high) using the updated Generalized International Standard Classification of Education (GISCED). To ascertain the relationship between educational status and the detection of diverticulosis, we employed multilevel logistic regression models, controlling for variables such as age, sex, metabolic syndrome, dietary habits, and physical activity. The local ethics committee for the province Salzburg approved the study protocol (approval no. 415-E/1262).

#### Conclusions

Individuals with lower educational status were more likely to have colon diverticula. People with a moderate level of education exhibited a lower incidence of diverticulosis, even after adjusting for relevant factors, with no apparent correlation to the distribution across different segments of the colon. This trend was also observed among individuals with higher levels of education. Implementing targeted strategies that consider socioeconomic conditions could potentially reduce the treatment costs associated with this condition.

- 1. Zajacova A, Lawrence EM. The relationship between education and health: Reducing disparities through a contextual approach. Annu Rev Public Health. 2018;39: 273–289.
- 2. Lukosiene JI, Reichert MC, Lammert F, Schramm C, Goeser T, Kiudelis G, et al. Environmental and dietary risk factors for colonic diverticulosis and diverticulitis. J Gastrointestin Liver Dis. 2021;30: 66–72.
- 3. Peery AF, Crockett SD, Barritt AS, Dellon ES, Eluri S, Gangarosa LM, et al. Burden of gastrointestinal, liver, and pancreatic diseases in the United States. Gastroentero. 2015;149: 1731–1741.e3.
#### Optimised intraoperative radiotherapy treatment workflow using machine learning methods

<u>Sara Vockner</u><sup>1</sup>, Markus Stana<sup>1</sup>, Christoph Gaisberger<sup>1</sup>, Franz Zehentmayr<sup>1</sup>, Klarissa Ellmauer<sup>1</sup>, Josef Karner<sup>1</sup>, Matthias Mattke<sup>1</sup>, Elvis Ruznic<sup>1</sup>, Falk Röder<sup>1</sup>

<sup>1</sup>Department of Radiotherapy and Radio-Oncology, Paracelsus Medical University, Salzburg, Austria; Contact: sara.vockner@stud.pmu.ac.at

#### Objective

Intraoperative radiotherapy (IORT) complements teletherapy in a broad spectrum of applications in the treatment of cancer by delivering radiation directly to the tumor during surgery. This has a number of advantages i.e. sparing of organs at risk and a precise localization and targeting of the tumor bed, allowing for the application of high doses in one fraction. On the other hand, advances in treatment planning in teletherapy, especially using voxel-based 3D models, have far surpassed IORT dose calculation concepts. Leaving aside the lack of appropriate imaging devices until recently, one of the main reasons for this discrepancy is the tight time schedule in an operative setting. This project aims to address the challenges related to the laborious and time-consuming nature of treatment planning, the need for precise applicator positioning and the necessity of optimal placement of radioprotection shielding.

#### Methods

This project introduces new algorithms, capitalizing on the synergy of artificial intelligence (AI), advanced image processing and computational methods. The first major objective is to develop and validate an artificial neural network to automatically contour organs at risk and the applicator on Cone Beam Computed Tomography (CBCT) scans, that were acquired with the mobile Imaging Ring-m during IORT procedures. In recent years, U-Net models were the most popular architectures in the domain of medical image segmentation. However, transformer network architectures became the new trend for deep learning models due to their capability to learn long-range dependencies and therefore handling global context. Although, it is essential to emphasize that transformers may not perform as effectively in capturing the finer spatial details in images, which is an area where U-Net excels. Consequently, the idea is to use a combination of the UNet and Transformer architecture to enhance the segmentation accuracy on CBCT images. The algorithm is trained and validated using a dataset consisting of 69 CBCT scans. This dataset is obtained from breast cancer patients undergoing IORT in the university clinic of radiotherapy in Salzburg. Building on the Al-based contouring model, an algorithm to accurately position the radiotranslucent applicator within the treatment planning software RADIANCE (by GMV) can be implemented. In addition to this, the orientation of radioprotection shields with respect to the applicator can be automatically verified. Ensuring that the shields are positioned perpendicular to the applicator, minimizes the exposure of organs at risk to radiation. The accuracy and effectiveness of the algorithm is evaluated through rigorous testing and comparison with established manual placement techniques.

#### Results

Implementation of state-of-the-art deep learning algorithms for automatic segmentation will help save time to allow for the adoption of 3D model based treatment planning in an operative setting. In a first step, automated precise applicator positioning will result in improved treatment planning accuracy, minimizing the risk of radiation-related complications and enhancing patient safety.

#### Conclusions

Closing the gap in treatment planning between teletherapy and IORT has the potential to optimise the quality of IORT treatments and ultimately improve patient outcomes.

- 1. A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. Gomez, et al. Attention Is All You Need, 2023, arXiv:1706.03762v7
- Ronneberger, O., Fischer, P., Brox, U-Net: Convolutional Networks for Biomedical Image Segmentation. In: Navab, N., Hornegger, J., Wells, W., Frangi, A. (eds) Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015. MICCAI 2015

# PKC activators orchestrate neuronal immune modulation: Unveiling microglial dynamics in NF-KB activation and phagocytosis

<u>Sofia Chanel Weinbender<sup>1,2,3</sup></u>, Marco Zattoni<sup>1,2</sup>, Bruno Benedetti<sup>2,4</sup>, Moritz Brosch<sup>2,3,4</sup>, Viktor Lippert<sup>2,4</sup>, Ariane Benedetti<sup>2,4</sup>, Sabine Bernegger<sup>1,2</sup>, Sebastien Couillard-Després<sup>2,4</sup>, Ludwig Aigner<sup>1,2</sup>

<sup>1</sup>Institute of Molecular Regenerative Medicine, Paracelsus Medical University, Salzburg, Austria;

<sup>2</sup>Austrian Cluster for Tissue Regeneration, Vienna, Austria;

<sup>3</sup>Department of Biosciences and Medical Biology, University of Salzburg, Salzburg, Austria;

<sup>4</sup>Institute of Experimental Neuroregeneration, Paracelsus Medical University, Salzburg, Austria; Contact: sofia.weinbender@pmu.ac.at

#### Objective

Chronic neuroinflammation, associated with central nervous system (CNS) pathologies, involves persistent microglial activation. Misfolded proteins in Alzheimer's disease and mechanical CNS lesions initiate prolonged immune responses, contributing to neuronal damage. Recent literature suggests the induction of controlled immune modulation to foster neuro-restoration. We propose the use of protein kinase C (PKC) activators to modulate CNS innate immune cells (i.e., microglia) for regenerative therapeutic approaches.

#### Methods

Therefore, we examined the molecular and functional effect of PKC activators (Prostratin and Bryostatin) in immortalized BV-2 cells and primary murine microglia.

#### Results

Western Blot (WB) and immunocytochemistry (ICC) analyses revealed significant increases in NF-KB phosphorylation (pNF-KB), one of the pathways downstream to PKC, in Prostratin-treated BV-2 cells. ICC analysis confirmed Prostratin-induced nuclear translocation of pNF-KB in BV-2 and primary murine microglia. Morphological alterations were investigated by quantifying microglial cell area and phagocytic activity was measured to explore microglial functionality. PKC activation increased microglial cell area associated with enhanced phagocytic activity. Finally, gene-expression and flow cytometry analysis were used to investigate the polarization of BV-2 microglia cells upon PKC activation. The absence of pro-inflammatory marker expression, together with the reduction of CD40 expression in lipopolysaccharide-exposed BV-2 cells, suggests potential neuroregenerative phenotypes for PKC-activated microglia cells.

#### Conclusions

Prostratin and Bryostatin's impact on microglial signaling reveals intricate complexities, with NF- KB activation hinting at pro-inflammatory responses. However, unexpected CD40 downregulation and absence of cytokine overexpression suggest diverse, unresolved pathways. The use of PKC activators holds therapeutic promise, offering transformative potential for neuroregenerative medicine by targeting microglial responses.

#### References

1. Ahmad MA, Kareem O, Khushtar M, Akbar M, Haque MR, Iqubal A, et al. Neuroinflammation: A Potential Risk for Dementia. Int J Mol Sci. 2022;23(2).

## Preliminary Data of a 3d Single-Limb-Squat Assessment of Laterality in Exercise-proficient Physiotherapists

Klaus Widhalm, Sebastian Durstberger, Peter Putz

<sup>1</sup>FH Campus Wien University of Applied Sciences, Health Sciences, 1100 Vienna, Austria; <sup>2</sup>Paracelsus Medical University, Institute of Biomechanics, 5020 Salzburg, Austria; Contact: klaus.widhalm@stud.pmu.ac.at

#### Objective

Single-Limb-Squat (SLS) is a common functional test and exercise in conditions such as ACL rupture, and degenerative knee or hip pathologies [1,2]. Lower limb valgus has been assessed two-dimensionally in numerous studies [3]. Although SLS is a multidimensional and neuromuscularly highly demanding task [4], only a few studies assessed its kinematics and kinetics three-dimensionally [2]. Previous pertinent studies differed in terms of sample size [5], non-consideration of the role of gender, or the role of health status and exercise proficiency on the SLS execution quality. So we investigated as a benchmark of performance quality if the dominant side differs from the non-dominant one regarding hip and knee frontal moments (fM) as well as frontal Range of Motion (fRoM) in performing SLS by a healthy exercise-proficient sample.

#### Methods

Eighteen (9 female / 9 male) healthy physiotherapists aged 19 to 62 (mean 34, sd 9.6) years performed five SLS with the right and left leg each. Data were captured using an optoelectronic movement analysis system with the modified Cleveland Clinical Markerset and a three-axis force plate. Processed trials were timenormalized for the down-phase and up-phase. Hip and knee fM, as well as fRoM were tested for difference between the dominant and non-dominant leg using one-dimensional parametric statistical mapping (1dSPM) with paired t-tests with alpha set at 0.05.

#### **Results**

The 1dSPM analyses did result in no statistically significant differences between the dominant and nondominant leg in any of the parameters for the lower limb. The null hypothesis was thus not rejected by the sample, although the trajectories show a clear deviation in the knee fM.

#### Conclusions

To our knowledge this is the first study providing three-dimensional reference data of SLS performed by a healthy exercise proficient sample. Laterality seems to be relevant for knee fM, when defining feedback thresholds. Parameters characterizing the quality of exercise execution related to the control of dynamic leg alignment show smaller values compared to prior studies. Gender differences are subject to the analysis of the complete dataset. As a limitation, squatting velocity and depth was monitored only verbally, which however did not affect the homogeneity of relevant biomechanical outcomes. Based on well-executed exercises, the physiological range of kinematic and kinetic parameters' cut-offs shall be established, by opposing the data of the exercise proficient sample to that of less experienced exercisers.

- 1. Jamaludin NI, Sahabuddin FNA, Rasudin NS, Shaharudin S. The Concurrent Validity and Reliability of Single Leg Squat Among Physically Active Females with and without Dynamic Knee Valgus. Int J Sports Phys Ther. 2022;17(4):574-584.
- 2022;17(4):574-584.
  Alenezi F, Herrington L, Jones P, Jones R. The reliability of biomechanical variables collected during single leg squat and landing tasks. Journal of Electromyography and Kinesiology. 2014;24(5):718-721.
- and landing tasks. Journal of Electromyography and Kinesiology. 2014;24(5):718-721.
  Lopes TJA, Ferrari D, Ioannidis J, Simic M, Azevedo FMd, Pappas E. Reliability and Validity of Frontal Plane Kinematics of the Trunk and Lower Extremity Measured With 2-Dimensional Cameras During Athletic Tasks: A Systematic Review With Meta-analysis. Journal of Orthopaedic & Sports Physical Therapy. 2018;48(10):812-822.
- Nguyen AD, Shultz SJ, Schmitz ŘJ, Luecht RM, Perrin DH. A preliminary multifactorial approach describing the relationships among lower extremity alignment, hip muscle activation, and lower extremity joint excursion. J Athl Train. 2011;46(3):246-256.
- Rossom Sv, Smith CR, Thelen DG, Vanwanseele B, Assche DV, Jonkers I. Knee Joint Loading in Healthy Adults During Functional Exercises: Implications for Rehabilitation Guidelines. Journal of Orthopaedic & Sports Physical Therapy. 2018;48(3):162-173.

# The degree of cyclic stretching influences lapine Achilles tenocytes and hMSCs in 2D culture and under 3D conditions on a Dura mater cell carrier

<u>Clea Wiedmann-Kline</u><sup>1</sup>, Clemens Gögele<sup>1</sup>, Jens Konrad<sup>2</sup>, Bernd Hoffmann<sup>2</sup>, Christian Werner<sup>1</sup>, Kerstin Schäfer-Eckart<sup>3</sup>, Andreas Traweger<sup>4</sup>, Andreas Teuschl-Woller<sup>5</sup>, Gundula Schulze-Tanzil<sup>1</sup>

<sup>1</sup>Institut für Anatomie und Zellbiologie, Paracelsus Medizinische Universität, Salzburg und Nürnberg, Nürnberg, Deutschland; <sup>2</sup>Institut für Biologische Informationsverarbeitung: IBI-2 Forschungszentrum Jülich, Deutschland; <sup>3</sup>Klinik für Onkologie/Hämatologie, Klinikum Nürnberg Nord, Nürnberg, Deutschland; <sup>4</sup>Institut für Sehnen und Knochenregeneration, Paracelsus Medizinische Universität, Salzburg und Nürnberg, Salzburg, Österreich; <sup>5</sup>Department Life Science Engineering, Fachhochschule Technikum Wien, Wien, Österreich; Contact: c.kline@gmx.net

#### Objective

Mobility can be severely impaired after tendon injuries. The self-healing and regenerative capacities of tendons are generally low. In the future, however, tissue- engineered constructs could be used to reconstruct injured tendons in order to prevent serious consequences. In the proposed basic research project, the effect of cyclic stretching on the formation of tendon tissue from both tenocytes and human mesenchymal stem cells (hMSCs) after 2D and 3D stimulation will be investigated in detail.

#### Methods

For the 2D culture approach (1), silicone chambers (4 cm2) were seeded with Lapine Achilles Tenocytes (LAT) on the one hand and human hMSCs (6000 cells/cm2) on the other. After 24 hours of adherence, the stimulated chambers were exposed to either moderate or strenuous exercise, while the control chambers were not exposed to stimulation. For the 3D culture approach, decellularized dura mater was recellularized with either LAS or hMSCs analogous to the 2D experiments and then subjected to moderate or strenuous exercise. At the end of each training session, viability was qualitatively assessed with a viability assay and DNA and sulfated glycosaminoglycan (sGAG) content was quantitatively measured. In addition, RNA was isolated to enable future gene expression analyzes of the most important tendon-associated matrix components and thus better characterize tendon development.

#### Results

The viability test showed that the cells were viable due to the careful preparation of the dura and that there were only a few dead cells in the tissue. The viability was over 50%. Histological staining showed that the architecture of the tissue had been preserved, and thus a suitable matrix for revitalization was present. The viability test showed that both LAT and hMSCs predominantly survived not only the moderate but also the strenuous training in 2D and 3D culture. A first trend in the alignment of the cells after stimulation is also recognizable. Quantitative analyzes confirm that moderate stretching has a greater influence on the sGAG synthesis of hMSCs than on LAT.

#### Conclusions

The porcine dura halves were carefully removed as a whole, so that a sufficiently large "scaffold matrix" was available for subsequent experiments. Initial differences in cell viability have already been observed, not only between LAT and hMSCs, but also between training programs. The effects of 3D stimulation on cell behavior are greater than those of 2D stimulation.

#### Acknowledgements

The work is financially supported by the Nürnberger Versicherung and the Kerscher Diabetes Foundation.

#### References

1. Goegele C, Hoffmann C, Konrad J, Merkel R, Schwarz S, Tohidnezhad M, et al. Cyclically stretched ACL fibroblasts emigrating from spheroids adapt their cytoskeleton and ligament-related expression profile. Cell Tissue Res. 2021;384(3):675-90. PMID: 33977325

#### Lessel-Kubisch syndrome as a model to decipher the Achilles heel of p53

Jana Willim<sup>1</sup>, Davor Lessel<sup>2</sup>

<sup>1</sup>Department of human genetic PMU, University Hospital Salzburg, Austria; <sup>2</sup>Department of human genetic PMU, University Hospital Salzburg, Austria; Contact: jana.willim@stud.pmu.ac.at

#### Objective

Inducing cellular senescence of cancer cells is considered a potentially effective strategy against cancer. Cellular tumor antigen p53 or just p53 is a regulatory protein and a transcription factor. It plays a key role in balancing cellular destiny between many different molecular pathways involved in cell division, senescence, DNA damage response and repair, and programmed cell death (apoptosis) when DNA damage is irreparable. p53 is activated by stress, however in normal conditions, p53 levels are kept low through a continuous degradation. This is achieved by protein interaction with MDM2, which translocates p53 from the nucleus to the cytosol and ubiq-uitilates p53, therefore marking it for proteasomal degradation.p53 is coded by TP53 gene, and variants in TP53 are associated with various cancer types, highlighting its significance in maintaining cellular integrity and preventing malignant transformation. Somatic TP53 loss-of-function variants are the most common alterations in cancer and a major cause of chemo-resistance. In addition to its role in cancer prevention, p53 contributes significantly to the ageing process by influencing several features of ageing.To gain a deeper insight in the p53-regulated balance between cancer and ageing, we aim to identify p53 down-stream effectors through transcriptome analysis. We hypothesize that this could bring us to identify novel p53 downstream targets, that could be used in personalized cancer therapy, where p53 is dysregulated.

#### Methods

Primary dermal fibroblasts, lymphoblastoid (LCL) cell lines, human osteosarcoma (U2OS) and non-small cell lung cancer (H1299) cell lines are used in the functional analyses including cellular localization studies, cytokinesis block micro-nucleus assay, chromosome analysis and stability assays. Transcriptome analyses are used to identi-fied p53 dependent dysregulated genes that can be use in therapeutic approaches.

#### Results

Previously, we discovered a homozygous anti-terminal mutation c.1492T>C in MDM2 that leads to Lessel-Kubisch syndrome (LSKB), characterized by premature ageing phenotype without tumor predisposition. On cellular level this specific homozygous C-terminal MDM2 variant leads to an increased p53 protein levels, increased p53 stability, and localization of p53 in the nucleolus, ultimately leading to an increased genome stability and decreased cell proliferation rate as well as a premature senescence phenotype. Transcriptome analysis of primary LSKB fibroblasts revealed 481 upregulated gene and 450 downregulated genes under normal conditions. From this, 35 up-regulated and 75 down-regulated were associated with senes-cence. We hypothesize that from these genes we could further select potential new targets for cancer treat-ment. Likewise, the identified genes will be classified in the cellular context and a signaling pathway analysis will be carried out. Top three hits for up- and downregulated genes will be further analyzed functionally

#### Conclusions

p53 as a tumor suppressor has been excessively studied as well how loss of p53 activity contributes to human cancers. In LSKB fibroblasts we observed increased p53 activation, along with cellular senescence as well as chromosomal stability. Therefore, we believe that LSKB are an excellent cellular model for the deeper characterization the p53 function, which may lead to new therapeutic options in oncology.

#### References

1. Lessel et al. Dysfunction of the MDM2/p53 axis is linked to premature aging. 2017, J Clin Invest. PMID: 28846075 PMCID: PMC5617664 DOI: 10.1172/JCI92171

#### The effect of UVC irradiation on germ density on breast implants and epidermis

Florian Wimmer<sup>1,2</sup>, Fabian Köninger<sup>1</sup>, Karl Schwaiger<sup>1</sup>, Gottfried Wechselberger<sup>1</sup>, Elisabeth Russe<sup>1</sup>

<sup>1</sup>Department of Plastic, Reconstructive and Aesthetic surgery, Hospital of the Brothers of St. John of God Salzburg, Salzburg, Austria; <sup>2</sup>Research Program of Medical Sciences, Paracelsus Medical University, Salzburg, Austria; Contact: florian.wimmer@stud.pmu.ac.at

#### Objective

Bacterial colonization of breast implants is a side effect of their implementation. This can lead to infection or capsular contracture, which is the most common long-term complication (1). Since these adverse events can lead to reoperation, prevention of bacterial colonization is essential. The germicidal effect of conventional 254 nm UVC light is already known (2), but presents a risk for human skin or eyes due to its mutagenic and cytotoxic effect. In contrast, previous studies showed, that far-UVC light (207-222 nm) efficiently inactivates germs (3) without representing a health risk to human skin cells because of its limited penetration ability (4). Therefore, we aim to prove that 222 nm UVC light irradiation of sterile implants and surrounding skin can lead to a significant reduction of germ density.

#### Methods

Five samples were obtained from each patient (n = 18) undergoing breast reduction surgery. Pre-incision samples (Sample I-III) were collected using sterile implant shells (Polytech Health & Aesthetics GmbH, Germany), applied to the patient's submammary fold during surgery following standard sterile procedures. Implant shells underwent 222 nm UVC irradiation (Disinfection cabinet, Sterilsystems GmbH, Austria) either before (Sample I) or after patient contact (Sample II), while one sample remained untreated (Sample III). After surgical skin removal, resected skin was irradiated with UVC light (Sample V) or left untreated (Sample IV) and pressed against an implant shell. Consequently, the sterile breast implant shells were then pressed onto agar contact slides (HYCON® Contact Slides, Merck Millipore, Germany) and incubated at 35°C for 24-48 hours. Microbiological analysis determined germ density and diversity of isolated microorganisms. All statistical analyses were conducted using NCSS (NCSS 2022, NCSS, LLC, Kaysville, UT).

#### Results

In the analysis of pre-incision samples, 17 out of 18 cases revealed either lower or absent germ presence in the UVC-treated Sample II compared to Samples I (p-value 0.06) and III (p-value 0.128). Subsequently, post-incision samples exhibited significantly reduced germ density in UVC-irradiated Sample V versus non-UVC-irradiated Sample IV (p-value = 0.037). Staphylococcus species were the most frequently identified bacterial species.

#### Conclusions

The study demonstrates the germicidal efficacy of 222 nm UVC light, indicating its potential role in reducing microbial contamination during surgery and consequently decreasing the risk of associated surgical side effects in implant-based breast surgery. However, further research is necessary to strengthen these preliminary findings.

- 1. Headon H, Kasem A, Mokbel K. Capsular Contracture after Breast Augmentation: An Update for Clinical Practice. Archives of Plastic Surgery. 2015;42(5):532. PMID: 26430623.
- 2. Miller SL, Linnes J, Luongo J. Ultraviolet Germicidal Irradiation: Future Directions for Air Disinfection and Building Applications. Photochemistry and Photobiology. 2013 May 20;89(4):777–81. PMID: 23581680.
- 3. Buonanno M, Welch D, Shuryak I, Brenner DJ. Far-UVC light (222 nm) efficiently and safely inactivates airborne human coronaviruses. Scientific Reports. 2020 Jun 24;10(1):1–8. PMID: 32581288.
- 4. Buonanno M, Ponnaiya B, Welch D, Stanislauskas M, Randers-Pehrson G, Smilenov L, et al. Germicidal Efficacy and Mammalian Skin Safety of 222-nm UV Light. Radiation research. 2017;187(4):483–91. PMID: 28225654.

#### DIAFILD - a study aiming to expedite DIAgnosis of Fibrosing Interstitial Lung Disease

<u>Mirja Wirtz</u><sup>1</sup>, Marie-Louise Geißler<sup>1</sup>, Jan Boor<sup>1</sup>, Stefan Zembacher<sup>1</sup>, Christina Imlinger<sup>1</sup>, Bernd Lamprecht<sup>2</sup>, Bernhard Kaiser<sup>2</sup>, David Lang<sup>2</sup>, Klaus Hackner<sup>3</sup>, Florentine Moazedi-Fuerst<sup>4</sup>, Benedikt Hergan<sup>5</sup>, Michael Studnicka<sup>1</sup>

<sup>1</sup>Department of Pulmonary Medicine, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Pulmonology, Kepler University Hospital, Linz, Austria; <sup>3</sup>Department of Pneumology, University Hospital Krems, Karl Landsteiner University of Health Sciences, Krems, Austria; <sup>4</sup>Division of Rheumatology and Immunology, Department of Internal Medicine, Medical University of Graz, Austria; <sup>5</sup>Department of Diagnostic and Interventional Radiology, Hospital Klagenfurt am Woerthersee, Klagenfurt, Austria; Contact: mirja.wirtz@stud.pmu.ac.at

#### Objective

Fibrosing interstitial lung diseases (FILD) are characterized by significant morbidity and mortality (1). Accurate and timely diagnosis of FILD is clinically challenging yet crucial to improve patient outcomes by early initiation of disease-specific anti-fibrotic therapy. This study was designed to evaluate the significance of combining readily available non-invasive diagnostic methods with the aim to identify FILD patients early.

#### Methods

DIAFILD is a prospective observational study targeted to evaluate the cumulative diagnostic value of surveys assessing FILD-specific symptoms – namely the University of California San Diego (UCSD) shortness of breath questionnaire and the Cough Assessment Test (COAT) – in combination with electronic lung auscultation (ELAUS) and thoracic ultrasound (THUS) in patients with (cases) and without (controls) FILD. Presence of fibrotic indicators on auscultation (disease-characteristic findings called Velcro crackles) and thoracic ultrasound (B-lines and pathological pleural lines) were assessed and correlated to presence of lung fibrosis on high-resolution CT scan (HRCT, gold standard of diagnosing FILD).

#### Results

Here, we present preliminary results featuring characteristics of 19 consecutive FILD patients (case group) who were enrolled in the study between February and October 2021. Females (n = 10) and males were equally represented, median age at the time of study enrollment was 77.0 years (range 60.0 to 85.0 years) and median time since diagnosis and the time of study enrollment was 3.3 years (range 0.3 to 43.5 years). At the time of the initial study visit, about half of the patients (52.6%) were receiving antifibrotic treatment. Mean pulmonary function tests at the time of the initial study visit were generally well-preserved with a FEV1%FVC of 0.86 (SD +/- 0.2), a predicted FVC of 86.4% (SD +/- 19.1), a predicted TLC of 97.8% (SD +/- 21.7) and a predicted DLCO of 67.6% (SD +/- 20.1). Both symptom-assessing surveys indicated a mildly symptomatic patient cohort (median scores of UCSD and COAT were 16 (range 0 to 45) and 3 (range 0 to 16), respectively). On ELAUS, Velcro crackles were found in most patients (n = 14; 73.7%). On THUS, most patients were found to have B-lines (n = 16; 84.2%) and pathological pleural lines (n = 17; 89.5%). Presence of pathological pleural lines on THUS and fibrotic changes on HRCT correlated strongly (Spearman's  $\rho$  = 0.561, p = 0.013), however, no correlation was found between presence of Velcro crackles and fibrotic findings on HRCT (Spearman's  $\rho$  = 0.001, p = 0.999).

#### Conclusions

FILD-characteristic diagnostic patterns were identified using non-invasive methods in the case group. Interim analysis showed a strong correlation between indicators of lung fibrosis on THUS and radiological presence of lung fibrosis on HRCT. Comparison to the corresponding findings of the control group has yet to take place.

#### References

 Nasser M, Larrieu S, Boussel L, Si-Mohamed S, Bazin F, Marque S, et al. Estimates of epidemiology, mortality and disease burden associated with progressive fibrosing interstitial lung disease in France (the PROGRESS study). Respir Res. 2021 May 24;22(1):162. PMID: 34030695

# A new approach for management of aneurysmal subarachnoid hemorrhage based on Manual and Artificial Intelligence blood volume measurement in Non-Contrast Head CT scans

<u>Melina Wirtz<sup>1,7</sup></u>, Saif Salman<sup>1</sup>, Yujia Wei<sup>2</sup>, Vishal Patel<sup>3</sup>, Rohan Sharma<sup>1</sup>, Vikash Gupta<sup>4</sup>, Qiangqiang Gu<sup>5</sup>, Benoit Dherin<sup>6</sup>, Sanjana Reddy<sup>6</sup>, Rabih Tawk<sup>1</sup>, William David Freeman<sup>1</sup>

<sup>1</sup>Departments of Neurological Surgery, Neurology and Critical Care, Mayo Clinic, Jacksonville, FL, 32224;

<sup>2</sup>Artificial Intelligence (AI) Laboratory, Department of Radiology, Mayo Clinic, Rochester, MN 55905;

<sup>4</sup>Center for Augmented Intelligence, Department of Radiology, Mayo Clinic, Jacksonville, FL 32224;

<sup>5</sup>Research Collaborator | Mayo Clinic College of Medicine and Science;

<sup>6</sup>Google, Inc. Mountain View, CA, 94043;

<sup>7</sup>Paracelsus Medical University, Nuremberg, Germany;

Contact: melina.wirtz@stud.pmu.ac.at

#### Objective

Utilize Artificial Intelligence (AI) and Machine-Learning (ML) methods to compute the volume of Subarachnoid hemorrhage blood (SAHV) and generate a 3D brain map from non-contrast CT scans. Compared to a Manual Method (MM), this approach aims to assess the precise volume of SAH blood more quickly, establish a potentially new imaging biomarker for just-in-time treatment options, and accelerate patient recovery.

#### Methods

The novel scan method, SAH Volumetric AI (SAHVAI-3D), was applied to a cohort of 10 patients (92 NCCT scans, Æ182 slices each) in direct comparison to a Manual Measurement Method (MM). A proprietary AI-ML algorithm generated SAH Volumetric AI (SAHVAI) values. In both methods (AI-ML, MM), a 3D SAH brain map was created for each patient. The SAHVs were analyzed and compared to neurological outcomes and visual areas of thickest blood concentration concerning vasospasm.

#### Results

SAHVAI quantified the volume of SAH blood in 42 seconds and demonstrated superior speed to the MM (>60 minutes). The SAHVAI showed an accuracy of 99.8%, Dice of 0.701, FPR=0.0005, and NPV=0.999. The absolute difference of SAHVs between both methods was 5.67ml. The SAHVAI-3D brain map and volume at admission appeared highly associated with neurological outcomes (inverse with Glasgow Coma Scale, linear with length of stay in hospital), especially in areas of thick blood concentration.

#### Conclusions

The introduction of SAHVAI-3D and the resultant "SAHVAI-3D Brain map" represent a new neurologic imaging biomarker. This innovation fills a significant gap in current SAH imaging techniques, enabling rapid evaluation and targeted interventions to enhance patient care.

#### Acknowledgements

Saif Salman MD, Yujia Wei MD, Vishal Patel MD PhD, Rohan Sharma MBBS, Vikash Gupta PhD, Qiangqiang Gu Phd, Benoit Dherin PhD, Sanjana Reddy MS, Rabih Tawk MD, William David Freeman MD

- 1. Salman S, Gu Q, Dherin B, et al. Hemorrhage Evaluation And Detector System for Underserved Populations: HEADS UP. Neurology; 2023. doi:10.1101/2023.06.25.23291870
- 2. Salman S, Gu Q, Sharma R, et al. Artificial Intelligence and Machine Learning in Aneurysmal Subarachnoid Hemorrhage: Future Promises, Perils, and Practicalities. Neurology; 2023. doi:10.1101/2023.07.18.23292822

<sup>&</sup>lt;sup>3</sup>Division of Neuroradiology, Department of Radiology, Mayo Clinic, Jacksonville, FL 32224;

# Wound healing and self organization of skin cells is guided by protein corona-bearing extracellular vesicles

<u>Martin Wolf</u><sup>1</sup>, Fausto Gueths Gomes<sup>1,3</sup>, Thomas Kocher<sup>5</sup>, Patricia Ebner-Peking<sup>1</sup>, Rodolphe W. Poupardin<sup>1</sup>, Anna M. Raninger<sup>1</sup>, André Cronemberger Andrade<sup>1</sup>, Nicole Maeding<sup>1</sup>, Constantin Blöchl<sup>2</sup>, Astrid Obermayer<sup>2</sup>, Balazs Vari<sup>1</sup>, Essi Eminger<sup>1</sup>, Heide-Marie Binder<sup>1</sup>, Sarah Hochmann<sup>1</sup>, Gabriele Brachtl<sup>1</sup>, Andreas Spittler<sup>4</sup>, Thomas Heuser<sup>6</sup>, Racheli Ofir<sup>7</sup>, Christian G. Huber<sup>2</sup>, Ulrich Koller<sup>5</sup>, Johann Bauer<sup>5</sup>, Zami Aberman<sup>7</sup>, Katharina Schallmoser<sup>3</sup>, Hans-Dieter Volk<sup>8</sup>, Dirk Strunk<sup>1,3</sup>

<sup>1</sup>Institute for experimental and clinical cell therapy, Paracelsus Medical University (PMU), Salzburg, Austria;

<sup>2</sup>Department of Biosciences, Paris Lodron University Salzburg, Austria;

<sup>3</sup>Department of Transfusion Medicine and SCI-TReCS, PMU, Salzburg, Austria;

<sup>4</sup>Anna Spiegel Center of Translational Research, Medical University, Vienna;

<sup>5</sup>EB House Austria, Research Program for Molecular Therapy of Genodermatoses, Department of Dermatology and Allergology, University Hospital of the Paracelsus Medical University Salzburg, Salzburg, Austria;

<sup>6</sup>Vienna Biocenter Core Facilities, Medical University, Vienna;

<sup>7</sup>Pluristem Ltd., Haifa, Israel;

<sup>8</sup>BCRT & Institute of Medical Immunology, Charite - Universitätsmedizin Berlin, Germany;

Contact: martin.wolf@pmu.ac.at

#### Objective

Transport of functional protein cargo via extracellular vesicles (EVs) is an important mechanism in cell communication. We aim to understand the distribution of active cargo proteins between EV's inside and outside and the implication for therapeutic applications in wound healing.

#### Methods

Here we used skin organoids and a human-on-mouse skin model to investigate the contribution of EVs to this process. EVs from placenta-derived stromal (PLX) cell-conditioned medium or from human platelet lysate were enriched by tangential flow filtration (TFF) optionally followed by ultracentrifugation or size exclusion chromatography (SEC) to obtain EV preparations with different degrees of protein corona preservation. Self organization capacity in presence or absence of different EV preparations was analyzed in a high throughput organoid assay. Contribution of EVs to wound healing was assessed in an in vivo model transplanting human skin cells on NSG mice1,2.

#### Results

Skin organoid formation by keratinocytes together with endothelial cells and keratinocytes or spheroid formation of human skin fibroblasts was significantly superior in the presence of TFF-purified EVs bearing a functional protein corona. Soluble factors alone or corona-depleted EVs after SEC did not initiate 3D organization. In vivo, TFF-purified EVs were essential for proper stratified organization of human skin cells in wound areas and provided sufficient vascular suport.

#### Conclusions

This findings allowed us to develop a self-organizing human skin model in mice and show the important role of corona proteins for EV function. The surface-to-bulk partition of EV cargo for small EVs < 180 nm is in favor of surface cargo loading supporting the new concept of a functional EV corona.

- 1. Wolf, M. et al. (2022) 'A functional corona around extracellular vesicles enhances angiogenesis, skin regeneration and immunomodulation', Journal of Extracellular Vesicles. John Wiley and Sons Inc, 11(4). doi: 10.1002/jev2.12207
- 2. Ebner-Peking, P. et al. Self-assembly of differentiated progenitor cells facilitates spheroid human skin organoid formation and planar skin regeneration. Theranostics 11, 8430–8447 (2021)

# Beyond Stress: The Role of Attachment and Mentalizing for the Well-being among Elementary Educators

Carmen Wusatiuk<sup>1,2</sup>, Antonia Dinzinger<sup>1,2</sup>, Beate Priewasser<sup>1,2</sup>

<sup>1</sup>Institute for Early Life Care, Paracelsus Medical University, Salzburg, Austria; <sup>2</sup>Department of Pediatrics, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria; Contact: carmen.wusatiuk@pmu.ac.at

#### Objective

The capacity to mentalize (1), in addition to the educator-child relationship (2), appears to function as a protective factor against stress, thereby enhancing the well-being of elementary educators, given their high workload. Educators who have positive mental models of relationships experience more positive affect in their interactions with children, which promotes their own well-being (3). Moreover, an appropriate perception of mental states allows for a more constructive approach to stressful situations (4), even in the absence of positive mental models of relationships, because negative affect can be identified and regulated. The aim of this study was to assess whether the ability to mentalize exerts a stronger influence on stress and well-being in an elementary educational context than attachment behavior.

#### Methods

To address the research question, we collected questionnaire data from N = 216 prospective elementary educators to examine their mentalizing capacity (MZQ), attachment behavior (ASQ), stress levels (TICS-SSCS), and well-being (SOC).

#### Results

Data revealed the anticipated correlations between attachment behavior and mentalization ability with wellbeing and stress levels. Additionally, mediation analyses showed that mentalization capacity partially explained the influence of attachment behavior on stress and well-being.

#### Conclusions

These findings indicate that attachment behavior plays a subordinate role in the context of stress and wellbeing, as its influence is partially explained by the ability to mentalize. Mentalizing can be trained and learned in adulthood (5), suggesting that mentalization training could support well-being and improve the abilities of elementary educators under stressful conditions.

- 1. Schwarzer, N. & Gingelmaier, S. (2020). Mentalisieren als schützende Ressource bei angehenden Erzieherinnen und Erziehern. Frühe Bildung, 9(3), 144–152. https://doi.org/10.1026/2191-9186/a000485
- Milatz, A., Lüftenegger, M. & Schober, B. (2015). Teachers' Relationship Closeness with Students as a Resource for Teacher Wellbeing: A Response Surface Analytical Approach. Frontiers in Psychology, 6. https://doi.org/10.3389/fpsyg.2015.01949
- 3. Spilt, J. L., Koomen, H. M. & Thijs, J. (2011). Teacher Wellbeing: The Importance of Teacher–Student Relationships. Educational Psychology Review, 23(4), 457–477. https://doi.org/10.1007/s10648-011-9170-y
- 4. Schwarzer, N., Nolte, T., Fonagy, P. & Gingelmaier, S. (2022). Self-Rated Mentalizing Mediates the Relationship Between Stress and Coping in a Non-Clinical Sample. Psychological Reports, 125(2), 742–762. https://doi.org/10.1177/0033294121994846
- Luyten, P., Campbell, C., Allison, E. & Fonagy, P. (2020). The Mentalizing Approach to Psychopathology: State of the Art and Future Directions. Annual Review Of Clinical Psychology, 16(1), 297–325. https://doi.org/10.1146/annurevclinpsy-071919-015355

# Short-term CD8+ T cells ablation reduces microgliosis in the hippocampus of old APP/PS1 animals

<u>Marco Zattoni</u><sup>1</sup>, Sophie-Marie Rieder<sup>1</sup>, Sabine Bernegger<sup>1</sup>, Barbara Altendorfer<sup>1</sup>, Ariane Benedetti<sup>2</sup>, Heike Mrowetz<sup>1</sup>, Sofia Weinbender<sup>1</sup>, Jennifer Forster<sup>3</sup>, Michael Stefan Unger<sup>4</sup>, Ludwig Aigner<sup>1,5</sup>

<sup>1</sup>Institute of Molecular Regenerative Medicine, Paracelsus Medical University, Salzburg, Austria;

<sup>2</sup>Institute of Experimental Neuroregeneration, Paracelsus Medical University, Salzburg, Austria;

<sup>3</sup>Salzburg Cancer Research Institute, Salzburg, Austria;

<sup>4</sup>Department of Biosciences and Medical Biology, University of Salzburg, Salzburg, Austria;

<sup>5</sup>Austrian Cluster for Tissue Regeneration, Salzburg, Austria;

Contact: marco.zattoni@pmu.ac.at

#### Objective

Alzheimer's disease (AD) progression has been recently associated to the accumulation of CD8+ T cells into disease-affected brain parenchyma, where they tightly associate with microglial and neuronal structures (1, 2). However, the functional role of CD8+ T cells in the AD brain is still elusive. Therefore, we investigated the impact of short-term CD8+ T cell ablation on neuroinflammation in transgenic AD mice (APP/PS1).

#### Methods

CD8+ T cells ablation was performed via intraperitoneal injections of anti-CD8 antibody for three days in twoyear-old APP/PS1 and wild-type male mice. Control groups received the respective isotype antibody (four animals per group). Heart-collected blood samples underwent flow cytometry analysis. After transcardiac perfusion, the brain tissue was processed for immunohistochemistry analysis.

#### Results

As expected, anti-CD8 injections resulted in a substantial decrease of CD8+ T cells from the blood circulation as well as brain parenchyma. We observed a slight reduction in amyloid plaque pathology upon anti-CD8 treatment and a statistically significant reduction in the percentage area of Iba1+ cells in the hippocampus of APP/PS1 mice treated with anti-CD8 antibody compared to isotype-treated mice.

#### Conclusions

Short-term CD8+ T cell ablation revealed a consistent reduction of microglial cell staining area in the hippocampus of treated animals. These results suggest that short-term CD8 ablation could influence microglia-driven neuroinflammation.

#### Acknowledgements

We thank the Microscope Core facility and the Animal facility at PMU for their support of this work.

- Unger MS, Li E, Scharnagl L, Poupardin R, Altendorfer B, Mrowetz H, et al. CD8 + T-cells infiltrate Alzheimer's disease brains and regulate neuronal- and synapse-related gene expression in APP-PS1 transgenic CD8 + T-cells infiltrate Alzheimer's disease brains and regulate neuronal- and synapse-related gene expression in APP-PS1 transgenic mice. Brain Behav Immun. 2020;89:67-86
- 2. Gate D, Saligrama N, Leventhal O, Yang AC, Unger MS, Middeldorp J, et al. Clonally expanded CD8 T cells patrol the cerebrospinal fluid in Alzheimer's disease. Nature. 2020;577(7790):399-404.

# 3D Scanning and Printing Technology for Breast Reconstruction: Increased Aesthetic Outcomes and Quality of Life

Maximilian Zaussinger<sup>1,2</sup>,Karl Schwaiger<sup>3</sup>, Gurdrun Ehebruster<sup>1</sup>, Manfred Schmidt<sup>1</sup>

<sup>1</sup> Section of Plastic and Reconstructive Surgery, Kepler University Hospital Linz Krankenhausstrasse 9, 4020 Linz, Austria;
 <sup>2</sup> Paracelsus Medical University, Doctoral Degree Program in Medical Science, Strubergasse 21, 5020 Salzburg, Austria;
 <sup>3</sup> Department of Plastic, Aesthetic and Reconstructive Surgery Hospital of the Barmherzige Brüder, Kajetanerplatz 1, 5010 Salzburg, Austria;

Contact: m.zaussinger@hotmail.com

#### Objective

Sculpting a flap of autologous tissue into an aesthetically pleasing breast represents an integral challenge in modern breast reconstruction.(1) Several methods of assessment have been investigated to deviate from individual artistic skills towards objective breast shaping. Among them, 3D technology might provide the most promising support but needs validation in applicability and accuracy. Hence, the aim of this study was to evaluate the potential influence of 3D technology on autologous breast reconstruction with regards on postoperative breast appearance and quality of life.

#### Methods

Within this study, outcomes of 24 female patients who underwent autologous breast reconstruction assisted with 3D technology were analysed and compared with previous similar patients without the use of 3D technology. Based on individual 3D scans and simulation, patient-specific templates were used for reconstruction. Quality of life assessment and aesthetic evaluation was performed and compared with the control group. Appropriate statistical analysis was carried out.

#### Results

A total of 24 female patients met inclusion criteria and underwent autologous breast reconstruction using 3D technology. Quality of life scores showed increased values, postoperatively as well as compared with the control group. Overall aesthetic evaluation revealed notable improved outcomes on breast appearance. A critical appraisal of external plastic surgeons showed statistically significant better outcomes on breast shape. All patients would recommend and repeat breast reconstruction using 3D technology.

#### Conclusions

Using 3D dimensional patient specific templates represents a practical method facilitating personalized flap planning and shaping. The results of this study demonstrate that 3D technology enhances outcomes in breast reconstruction concerning aesthetics as well as postoperative quality of life.

#### References

1. Hunsinger V, Hivelin M, Derder M, Klein D, Velten M, Lantieri L. Long-Term Follow-Up of Quality of Life following DIEP Flap Breast Reconstruction. Plast Reconstr Surg. 2016 May;137(5):1361–71. PMID: 26796375

# Impact of 2-hydroxypropyl- $\beta$ -cyclodextrin inclusion complex formation on dopamine receptor-ligand interaction – a case study

<u>Lukas Zell</u><sup>1</sup>, Thomas S. Hofer<sup>3</sup>, Mario Schubert<sup>4, 5</sup>, Alexander Popoff<sup>1</sup>, Anna Höll<sup>1</sup>, Moritz Marschhofer<sup>1</sup>, Petra Huber-Cantonati<sup>2</sup>, Veronika Temml<sup>1</sup>, Daniela Schuster<sup>1</sup>

<sup>1</sup>Department of Pharmaceutical and Medicinal Chemistry, Institute of Pharmacy, Paracelsus Medical University, 5020 Salzburg, Austria; <sup>2</sup>Department of Pharmaceutical Biology, Institute of Pharmacy, Paracelsus Medical University, 5020 Salzburg, Austria; <sup>3</sup>Institute of General, Inorganic and Theoretical Chemistry, Center for Biochemistry and Biomedicine, University of Innsbruck, 6020 Innsbruck, Austria; <sup>4</sup>Department of Biosciences and Medical Biology, University of Salzburg, 5020 Salzburg, Austria; <sup>5</sup>Department of Chemistry, Freie Universität Berlin, 14195 Berlin, Germany; Contact; lukas.zell@pmu.ac.at

#### Objective

The octanol-water distribution coefficient (logP), used as a measure of lipophilicity, plays a major role in the drug design and discovery processes. While average logP values remain unchanged in approved oral drugs since 1983, current medicinal chemistry trends towards increasingly lipophilic compounds that require adapted analytical workflows and drug delivery systems. Solubility enhancers like cyclodextrins (CDs), especially 2-hydroxypropyl- $\beta$ -CD (2-HP- $\beta$ -CD), have been studied in vitro and in vivo investigating their ADMET (adsorption, distribution, metabolism, excretion and toxicity)-related properties. However, data is scarce regarding the applicability of CD inclusion complexes (ICs) in vitro compared to pure compounds.

#### Methods

In this study, dopamine receptor (DR) ligands were used as a case study, utilizing a combined in silico / in vitro workflow. Media-dependent solubility and IC stoichiometry were investigated using HPLC. NMR was used to observe IC formation-caused chemical shift deviations while in silico approaches utilizing basin hopping global minimization were used to propose putative IC binding modes. A cell-based in vitro homogeneously time-resolved fluorescence (HTRF) assay was used to quantify ligand binding affinity at the DR subtype 2.

#### Results

While all ligands showed increased solubility using 2-HP- $\beta$ -CD, they differed regarding IC stoichiometry and receptor binding affinity. This case study shows that IC-formation was ligand-dependent and sometimes altering in vitro binding.

#### Conclusions

Therefore, IC complex formation can't be recommended as a general means of improving compound solubility for in vitro studies as they may alter ligand binding.

- 1. Gleeson, M.P., Hersey, A., Montanari, D. and Overington, J., Probing the links between in vitro potency, ADMET and physicochemical parameters. Nat Rev Drug Discov, 2011. 10(3): p. 197-208. DOI: 10.1038/nrd3367. PMID: 21358739.
- Jun, S.W., et al., Preparation and characterization of simvastatin/hydroxypropyl-beta-cyclodextrin inclusion complex using supercritical antisolvent (SAS) process. Eur J Pharm Biopharm, 2007. 66(3): p. 413-21. DOI: 10.1016/j.ejpb.2006.11.013. PMID: 17240129.
- 3. Duchene, D. and Bochot, A., Thirty years with cyclodextrins. Int J Pharm, 2016. 514(1): p. 58-72. DOI: 10.1016/j.ijpharm.2016.07.030. PMID: 37765313.
- 4. Higuchi, T. and Connors, K.A., Phase Solubility Techniques. Adv. Anal. Chem. Instrum., 1965. 4: p. 117-212. DOI: 10.1016/S0076-6879(82)87029-8.

#### BEATS – MIGRAINE: Biomarker Establishment of Activated Targets in Skin for Migraine

<u>Janine Zwicklhuber</u><sup>1</sup>, Sarah Hochmann<sup>1</sup>, Antri Stefani<sup>1</sup>, Martin Wolf<sup>1</sup>, Nicole Maeding<sup>1</sup>, Rodolphe W. Poupardin<sup>1</sup>, Stefan Leis<sup>2</sup>, Daniela D. Weber<sup>3</sup>, Julia Tevini<sup>3</sup>, Barbara Kofler<sup>3</sup>, Alena Rückl<sup>4</sup>, Mark Rinnerthaler<sup>5</sup>, Elisabeth Russe<sup>4</sup>, Dirk Strunk<sup>1</sup>

<sup>1</sup>Institute of Experimental and Clinical Cell Therapy, PMU, Salzburg, Austria; <sup>2</sup>Dept. of Neurology, Christian Doppler University Hospital of the PMU, Salzburg, Austria; <sup>3</sup>Research Program for Receptor Biochemistry and Tumor Metabolism, Dept. of Pediatrics, University Hospital of the PMU, Salzburg, Austria; <sup>4</sup>Dept. of Plastic, Aesthetic and Reconstructive Surgery, Barmherzige Brüder, Salzburg, Austria; <sup>5</sup>Dept. of Biosciences and Medical Biology, University of Salzburg, Austria; Contact: janine.zwicklhuber@pmu.ac.at

#### Objective

Migraine is a common neurological disease affecting about one in five women under 50, making it the third most common worldwide (1). Until now, migraine diagnosis is based on clinical criteria as there is still no reliable test or any biomarker to validate the diagnosis (2). Temperature perception, olfactory sensitivity and osmophobia have been shown to play a considerable role in migraine (3, 4). Growing evidence shows that migraine pain may be caused by irritation of peripheral extracranial nerves, vessels or other soft tissues which led to innovative treatment options. Surgical extracranial trigger deactivation by decompression or removal of nerves and/or vessels in affected areas for severe migraineurs is a promising surgical treatment option (5-7). This procedure is accompanied by a simultaneous upper eyelid blepharoplasty. Immunohistochemistry (IHC) of these biopsies will reveal the spatial distribution of biomarkers and provide information on differences in expression levels. Extracellular vesicles (EVs) are key players in cell-to-cell communication in health and disease (8). Isolated EVs from blood samples will be the basis for proteome analysis to gain further insight into potential differences between healthy and migraine. Establishment of an EV-migraine signature allows validation of migraine diagnosis and will help to identify further treatment options for refractory patients. We hypothesize that spatial distribution, gene expression and EV signature in skin differs in patients diagnosed with migraine and migraine-free controls. Our goals are therefore to discover a migraine signature allowing for biomarker identification of activated receptors and other genes in patient skin and the establishment of skin as a testing system for migraine interventions and/or drug screening.

#### Methods

The department of Plastic, Aesthetic and Reconstructive Surgery at the Barmherzige Brüder in Salzburg offers the above-mentioned surgical treatment for migraineurs and is therefore eligible to collect skin samples for further examination. To achieve the stated goals we use qPCR, next generation sequencing (NGS), western blot, histology and immunocytochemistry, flow cytometry, establishment of tissue culture models as well as metabolomics. For isolation of patient EVs we follow our recently established protocol (9).

#### Results

Collection of skin biopsies and blood from migraine patients and migraine-free controls started with the beginning of May. Primary keratinocytes and fibroblasts have already been isolated from several biopsies and skin sections have been embedded in paraffin for subsequent IHC staining.

#### Conclusions

The establishment of a migraine signature will allow validation of migraine diagnosis and will probably help in the identification of further treatment options for refractory patients.

#### References

 Lampl C. et al. 2003; (2) Headache Classification Committee of the International Headache Society 2018; (3) Sjöstrand C. et al. 2010; (4) Gossrau G. et al. 2022; (5) Burstein R. et al. 2017; (6) Guyuron B. et al. 2014; (7) Perry CJ. et al. 2016; (8) Wolf M. et al. 2022; (9) Stefani A. et al. 2024.

### Liste der Einrichtungen (korrespondierende Autor\*innen)

(in alphabetischer Reihenfolge)

- Ankara University, Ankara
- Berufsgenossenschaftliche Unfallklinik Murnau Lehrkrankenhaus der PMU Salzburg
- Kepler Universitätsklinikum
- Ludwig Boltzmann Institut für Traumatologie
- Paracelsus Medizinische Privatuniversität Nürnberg / Klinikum Nürnberg
- Paracelsus Medizinische Privatuniversität Salzburg / Universitätsklinikum Salzburg
- Paris Lodron Universität Salzburg
- Red Bull Athlete Performance Center
- Südtiroler Sanitätsbetrieb SABES, Brixen
- Slovak Academy of Sciences, Bratislava
- Technischen Hochschule Nürnberg Georg Simon Ohm
- Technische Universität München Universität Wien
- University College London
- ZURZACH Care Klinik für Schlafmedizin Bad Zurzach

### Verzeichnis aller Autor\*innen

(fett gedruckt = Erstautor\*in)

Autor*in	Abstract Nr.	В
Α		B
Abed, Selim	1	В
Aberman, Zami	97	В
Ablinger, Celina	2	B
Acker, Jells Georg	<b>3</b> 15 57 88	D
Aigner, Lindi Aigner, Ludwig	40, 07, 00 90, 99	B
Alberti Paola	8	B
Altendorfer, Barbara	99	B
Ammon, Josephine	11	В
Angerer, Matthias	47, 69	В
Augat, Peter	5, 58, 85	
Aurnhammer, Felix	62	C
Avramovska, Irina	29	C
		C
B Barth Andrá A	15 40	
Bauer Andre	40	C
Bauer, Florian	4	C C
Bauer, Johann	97	Č
Baumeister, Dirk	5	
Beckert, Manuela	6	D
Beil-Hildebrand, Margitta	7, 46	D
Benedetti, Ariane	90, 99	D
Benedetti, Bruno	90	D
Bergamo, Mattia	8	D
Berger, Ursula	32 0 27 49 61 90	U D
Bernarder Sabine	9, 37, 40, 01, 00	ם ח
Bernik Rok	90, 99 47	ם
Berns, Maresa D.	15	D
Bieler, Lara	71	D
Billner, Moritz	<b>10</b> , 15, 40	D
Binder, Heide-Marie	97	D
Birkholz, Lorri	7	D
Bitti, Roberta	11	D
Blöchl, Constantin	97	D
Bock, Marlene	32	D
Bogensperger, Kathrin	// E	D
Boor lan	5 95	
Door, Jan	30	

Borecká, Silvia Brachtl, Gabriele Brandstötter, Christina Braun, Tobias Breidung, David Breitwieser, Martin Brisch, Karl Heinz Brosch, Moritz Brückl, Wolfgang Büchner, Susanne Bükki, Johannes Bürgler, Elisabeth Büscher, Andreas	<b>12,</b> 10,	<b>13,</b> <b>15</b> , 33	80 97 34 40 <b>16</b> ,39 90 81 <b>17</b> 67 24 46
<b>C</b> Caciagli, Lorenzo Cadamuro, Janne Cesur, Nevra Pelin Chaves, Gustavo Couillard-Després, Seba Crespo Pimentel, Berna Cronemberger, André	astien rdo	19,	20 41 <b>18</b> 53 90 <b>20</b> 97
<b>D</b> Datz, Christian Datz, Leonora Dauriz, Marco Deisl, Christine Devlitšarov, Daniel Delavari, Sarina Diener, Markus Dherin Benoit		45, 29,	88 88 57 55 <b>21</b> 15 30 96
Dinzinger, Antonia Dnyansagar, Rohit Dono, Fedele Dorenkamp, Andrea Dossena, Silvia Dubecz, Attila Duncan, John Durstberger, Sebastian Duszka, Kalina	9, 37, 48, 49,	31, 61, 29,	39 82 64 59 80 30 20 91 82

#### Ε Gantschnigg, Antonia 34 Ebbert, Ronald 22 Gašperíková, Daniela 80 Ebner-Peking, Patricia 97 Gehwolf, Renate 18 73 Geißler. Marie-Louise 95 Eibl, Thomas Edo, Alejandro Marti 15 Gensluckner, Sophie 45.57 Egle, Alexander 43 Gerber, Luis 59 100 Ehebruster, Gurdrun Ghiani, Alessandro 65 10, 15, 40 Girotto, Giorgia 48 Ehrl. Denis 75 Giulini, Luca **29**, 30 Engl, Michael Eisendl. Agatha 17 Glowalla, Claudio 58 Ellmauer, Klarissa 89 Gögele, Clemens 12, 13, 14, 27, 28, 92 Gosch, Markus Eminger, 32, 52 Emmanuel, Katja 16 Grabmer, Christoph 42 Emmanuel, Klaus 30 Grechenig, Michael 30 Enders, Bernadette 59 Greif, Elke 31 Erichsen, Sandra 85 Greil. Richard 43 Ernst, Anna Maria 55 Grieb, Gerrit 15 Eschermann, Kirsten 23 Gröger, Michael 59 Esenay, Isik Figen 24 Gu, Qiangqiang 96 Guenzler, Valentin 47 F Gueths Gomes, Fausto 97 Farmaki, Filitsa 30 Gupta, Vikash 96 Faustini, Bettina 84 Felder, Thomas K 82 н Ferini-Strambi, Luigi 72 Haas, Christoph 26 32 81 Habboub, Basel Ficker, Joachim Fischer, Florian 25 Hackl, Simon 85 Fischer. Thorsten 8 Hackner, Klaus 95 Fischlein, Theodor 13, 47, 69 Hantinger, Marion 33 Flamm, Maria 45,88 Haller, Sophie 70 12.28 Hanneken, Jessica 87 Fleischmann, Nils Flore, Davide 72 Harrer. Andrea 71 26 Föhrenbach, Kim 43 Hartl, Arnulf 42 Haslauer, Theresa 34 Föttinger, Fabian 26 59 Foisner, Vera Hass, Alexandra 99 Heger, Patrick Forster, Jennifer 30 Fuchs, Julia 77 Hergan, Benedikt 95 Herrmann, Sven 58 Franco, Valentina 64 Freeman, William David 96 Hertle, Christine 36 Heuser, Thomas Freude, Thomas 16 78,97 Frey, Vanessa 44 Hitzl, Wolfgang 15, 29, 30, 35, 69 Fricke, Angela 36 Hochmann, Sarah 78, 83, 97, 102 Fusch, Christoph 22 Höfler, Julia 20 Höhl. Rainer 35 G Höll, Anna 101 Gaisbauer, Stefanie 34 Höllerer, Raffael 2 Gaisberger, Christoph 89 Hoffmann, Bernd 92 36 Gantenbein, Andreas 3 Hofmann, Hanna

XII

Huber, Christian G.      97      Kohl, Thomas      62        Huber, Christisoph      26      Kokozidou, Maria      12, 13, 14        Huber, Florian      37, 48, 61      Koller, Ulrich      97        Huber, Sara      34      Konrad, Jens      92        Huber, Axel      65      Kubitz, Jens      35        Huten, Clemens      35      Kuchkhidze, Giorgi      20        Hüttner, Kelk      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungrer, Sven      58      Krautk-Köstler, Eva      36        Iglseder, Bernhard      45      L      1        Ismair, Selina      39      Lang, David      95        Jager, Simon      65      Lamer-Plamberger, Sandra      42        Jakas, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        K      Leigname, Giuseppe      49        Kaiser, Bernhard      95      Leis, Stefan      102        Kaiser, Gurfun      64	Hofmann, Sabrina	22	Kofler, Barbara	34, 82, 102
Huber, Christoph      26      Kokozidou, Maria      12, 13, 14        Huber, Forian      37, 48, 61      Kolrer, Ulrich      97        Huber, Sara      34      Konrad, Jens      92        Huber-Cantonati, Petra      2, 56, 101      Koutny, Florian      45, 57        Huber, Axel      65      Kubitz, Jens      35        Hütther, Felix      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hunger, Christian      95      Lamprecht, Bernd      95        Igleder, Bernhard      45      L      1        Inlinger, Christina      95      Lamprecht, Bernd      95        Jager, Simon      65      Langr, H.      33        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Leigner, Giuseppe      49        Kaier, Bernhard      95      Leigner, Giuseppe      49        Kaisinki, Jurij Matija      47      Leigner, Stonas      102	Huber, Christian G.	97	Kohl, Thomas	62
Huber, Florian      37, 48, 61      Koller, Ulrich      97        Huber, Sara      34      Konrad, Jens      92        Huber, Axel      65      Kubitz, Jens      35        Huteher, Kelik      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungrer, Sven      58      Krauk-Köstler, Eva      36        Iglseder, Bernhard      45      L      1        Ismair, Selina      39      Lang, David      95        Jager, Simon      65      Lamprecht, Bernd      95        Jager, Simon      65      Langthaler, Patrick      45        Jager, Simon      67      Lauth, Wanda      42        Lagmane, Giuseppe      49      49        Kaiser, Bernhard      95      Lauth, Wanda      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Kaiser, Bernhard      95      Leit, Mic	Huber, Christoph	26	Kokozidou, Maria	12, 13, 14
Huber, Sara      34      Konrad, Jens      92        Huber-Cantonati, Petra      2, 56, 101      Kouthy, Florian      45, 57        Hueber, Axel      65      Kubitz, Jens      35        Hütther, Feix      29, 30      Kundt, Firuzan      36        Hüttner, Feix      29, 30      Kundt, Firuzan      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungrer, Sven      58      Krauß-Köstler, Eva      36        I      Krutschar, Patrick      7, 67        Inlinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, H.      33        J      Langthaler, Patrick      45        Jäger, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Leit, Stefan      102        Kaiser, Bernhard      95      Leis, Michael      41        Kaiser, Bernhard      95      Leit, Michael      42        Jakab, Martin      9, 37, 61      La	Huber, Florian	<b>37</b> , 48, 61	Koller, Ulrich	97
Huber-Cantonati, Petra      2, 56, 101      Koutny, Florian      45, 57        Hueber, Axel      65      Kubtz, Jens      35        Hüttenbrink, Clemens      35      Kuchukhidze, Giorgi      20        Hütten, Felix      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungerer, Sven      58      Krauß-Köstler, Eva      36        I      I      I      I      I        Iglseder, Bernhard      45      L      I        Imlinger, Christina      39      Lang, David      95        Ismair, Selina      39      Lang, H.      33        J      Lang, H.      33      J        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Jahansson, Tim      87      Leigname, Giuseppe      49        Kaiser, Bernhard      95      Leis, Stefan      102        Kaises, Gudrun      64      Lell, Michael      11        Kamabusen, Anne      35      Lenhart, Armin      27, 28        Karat, Jorkan      4	Huber, Sara	34	Konrad, Jens	92
Hueber, Axel      65      Kubitz, Jens      35        Hüttner, Keinx      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Iglseder, Bernhard      45      L      1        Imlinger, Christina      95      Lang, David      95        Jager, Simon      65      Langr, H.      33        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Legmer, Giuseppe      49      47      Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      11      1      Karner, Kornad      10, 15, 40      Lessel, Davor      93        Karner, Josef      89	Huber-Cantonati, Petra	2, 56, 101	Koutny, Florian	<b>45</b> , 57
Hüttenbrink, Clemens      35      Kuchukhidze, Giorgi      20        Hüttner, Felix      29, 30      Kundt, Firuzan Sari      7        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungerer, Sven      58      Krauß-Köstler, Eva      36        I      Iglseder, Bernhard      45      L        Imlinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, David      95        Jäger, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Kaiser, Bernhard      95      Leiler, Spela      47        Kaisers, Gudrun      64      Lell, Michael      10        Kaisers, Gudrun      64      Lell, Michael      11        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Kart, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Ork	Hueber, Axel	65	Kubitz, Jens	35
Hüther, Felix      29, 30      Kundt, Firuzan Sari      7, 67        Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungerer, Sven      58      Krauß-Köstler, Eva      36        Iglseder, Bernhard      45      L        Inlinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, David      95        Jager, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kalisnik, Jurij Matija      47      Leisch, Michael      41        Karls, Gudrun      64      Leil, Michael      41        Karbar, Jorkan      42      Liebert, Adrian      73        Kera	Hüttenbrink, Clemens	35	Kuchukhidze, Giorgi	20
Hummel, Christian      38      Kutschar, Patrick      7, 67        Hungerer, Sven      58      Krauß-Köstler, Eva      36        I      I      I      I        Iglseder, Bernhard      45      L      I        Inlinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, David      95        J      Langthaler, Patrick      45        Jäger, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Kaser, Bernhard      95      Leis, Stefan      1002        Kaiser, Bernhard      95      Leis, Stefan      102        Kaiser, Gurun      64      Lell, Michael      41        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Karz, Konrad      10, 15, 40      Lessel, Davor      93        Kart, Orkan      42      Liebert, Adrian      73        Kerner, Josef      89      Ley, Pauline <t< td=""><td>Hüttner, Felix</td><td>29, 30</td><td>Kundt, Firuzan Sari</td><td>7</td></t<>	Hüttner, Felix	29, 30	Kundt, Firuzan Sari	7
Hungerer, Sven    58    Krauß-Köstler, Eva    36      I    Iglseder, Bernhard    45    L      Iglseder, Bernhard    95    Lamprecht, Bernd    95      Ismair, Selina    39    Lang, David    95      J    Lang, David    95    Lang, David    95      Jäger, Simon    65    Laner-Plamberger, Sandra    42      Jakab, Martin    9, 37, 61    Lapinskas, Simonas    15      Johansson, Tim    87    Lauth, Wanda    42      K    Legname, Giuseppe    49      Kaiser, Bernhard    95    Leis, Stefan    102      Kalss, Gudrun    64    Leil, Michael    11      Kamphausen, Anne    35    Lenhart, Armin    27, 28      Kart, Tanja    41    Lether, Thomas    84      Karmer, Josef    89    Ley, Pauline    78      Kartal, Orkan    42    Liebent, Adrian    73      Kiener, Teresa    2    Liuni, Raffaella    37, 48, 49      Kiener, Jorensa    84    Ling, Raffaella    37, 48, 49      Kiener, Presa    2	Hummel, Christian	38	Kutschar, Patrick	7, 67
Krutter, Simon46IIImlinger, Christina95Ismair, Selina39Lang, H.33Jager, Simon65Jäger, Simon65Jakab, Martin9, 37, 61Lapinskas, Simonas15Johansson, Tim87Lauth, Wanda42Leginskas, Simonas15Johansson, Tim87KLeiler, SpelaKLeiler, SpelaK47Kaiser, Bernhard95Kaiser, Bernhard95Lauth, Wanda42Kalas, Gudrun64Kalas, Gudrun64Karner, Josef89Karner, Josef89Karner, Josef89Karner, Josef89Kerry, Chassan73Kindinger, Barbel62Liobmann, Sebastian50, 51Kiener, Teresa2Liuni, Raffaella37, 48, 49Kiener, Teresa2Liuni, Raffaella37, 48, 49Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Teresa2Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Teresa2Kiensi, Teresa2Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jochen65Kiensi, Jo	Hungerer, Sven	58	Krauß-Köstler, Eva	36
IIglseder, Bernhard45LIndinger, Christina95Lamprecht, Bernd95Ismair, Selina39Lang, David95Jager, Simon65Laner-Plamberger, Sandra42Jakab, Martin9, 37, 61Lapinskas, Simonas15Johansson, Tim87Lauth, Wanda42Kaiser, Bernhard95Leis, Stefan1002Kaiser, Bernhard95Leis, Stefan102Kaisnik, Jurij Matija47Leisch, Michael43Kalssik, Surdurun64Lell, Michael41Karaz, Konrad10, 15, 40Lessel, Davor93Kartal, Orkan41Lettner, Thomas84Kartal, Orkan42Liebert, Adrian73Kerry, Ghassan73Lindlbauer, Nadja42Kiener, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kienslich, Tobias56Lorenzl, Stefan67Kinglingayer, Eva70MKabirik, Joria53Kocher, Thomas97Maeding, Nicole78, 83, 97, 102Knab, Katja22Maeding, Nicole <td>-</td> <td></td> <td>Krutter, Simon</td> <td>46</td>	-		Krutter, Simon	46
Iglseder, Bernhard      45      L        Imilinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, David      95        Jager, Simon      65      Langrhaler, Patrick      45        Jäger, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Legname, Gluseppe      49      49        K      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kaiser, Gursphausen, Anne      35      Lenhart, Armin      27, 28        Kardz, Konrad      10, 15, 40      Lessel, Davor      93        Kart, Tanja      41      Lether, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kerny, Ghassan      73      Lindblauer, Nadja      42        Kieninger, Bärbel      62      Lobmann, Sebastian      50, 51	I			
Imlinger, Christina      95      Lamprecht, Bernd      95        Ismair, Selina      39      Lang, David      95        Jager, Simon      65      Langr. Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Legname, Giuseppe      49      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Kartal, Orkan      41      Letther, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kiem, Dorninik      43      Lippert, Viktor      90        Kiener, Teresa      2      Liuni, Raffaella      37, 48, 49        Kieninger, Bärbel      62      Lobman, Sebastian      50, 51        Kiesslich, Tobias      56      Lorenzl, Stefan      67	lglseder, Bernhard	45	L	
Ismarr, Selina    39    Lang, David    95      Jager, Simon    65    Langthaler, Patrick    45      Jakab, Martin    9, 37, 61    Lapinskas, Simonas    15      Johansson, Tim    87    Lauth, Wanda    42      K    Legname, Giuseppe    49      K    Leiler, Spela    47      Kaiser, Bernhard    95    Leis, Stefan    102      Kalass, Gudrun    64    Lell, Michael    11      Kamphausen, Anne    35    Lenhart, Armin    27, 28      Karcz, Konrad    10, 15, 40    Lessel, Davor    93      Karl, Tanja    41    Lettner, Thomas    84      Karner, Josef    89    Ley, Pauline    73      Kerny, Ghassan    73    Lindbauer, Nadja    42      Kiem, Dominik    43    Lippert, Viktor    90      Kiener, Teresa    2    Liobman, Sebastian    50, 51      Kiensley, Jochen    65    Lutz, Nina    65      Kiewull, Lorenz    44    Lubar, Francesca    52      Kienner, Friederike    65    Lutz, Nina    65 <td>Imlinger, Christina</td> <td>95</td> <td>Lamprecht, Bernd</td> <td>95</td>	Imlinger, Christina	95	Lamprecht, Bernd	95
J      Lang, H.      33        Jäger, Simon      65      Langthaler, Patrick      45        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Legname, Giuseppe      49      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kanzs, Gudrun      64      Lell, Michael      11        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Karcz, Konrad      10, 15, 40      Lessel, Davor      93        Karl, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kiener, Melissa      30      Liesenberg, Tanja      28        Kerry, Ghassan      73      Lindlbauer, Nadja      42        Kiener, Teresa      2      Lioni, Raffaella      37, 48, 49        Kiener, Friederike      65	Ismair, Selina	39	Lang, David	95
J      Langthaler, Patrick      45        Jäger, Simon      65      Laner-Plamberger, Sandra      42        Jakab, Martin      9, 37, 61      Lapinskas, Simonas      15        Johansson, Tim      87      Lauth, Wanda      42        Legname, Giuseppe      49      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kalss, Gudrun      64      Lell, Michael      11        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Kart, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kemeter, Melissa      30      Liesenberg, Tanja      28        Kerry, Ghassan      73      Lindibauer, Nadja      42        Kiesnik, Jochen      65      Lorenzl, Stefan      67        Kieninger, Bärbel      62      Lobmann, Sebastian      50, 51        Kiesslich, Tobias			Lang, H.	33
Jäger, Simon65Laner-Plamberger, Sandra42Jakab, Martin9, 37, 61Lapinskas, Simonas15Johansson, Tim87Lauth, Wanda42Legname, Giuseppe49Leiler, Spela47Kaiser, Bernhard95Leis, Stefan102Kaissik, Jurij Matija47Leisch, Michael43Kalss, Gudrun64Lell, Michael11Kamphausen, Anne35Lenhart, Armin27, 28Karcz, Konrad10, 15, 40Lessel, Davor93Kartal, Orkan42Liebert, Adrian73Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kienn, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kienslich, Tobias56Lutz, Nina65Klenner, Friederike65Lutz, Nina65Klenner, Friederike65Lutz, Nina63Kochen65Kutz, Nina5354Kocher, Thomas97Maeding, Nicole78, 83, 97, 102Knab, Katja22Maeding, Nicole78, 83, 97, 102Knab, Katja32Mahar, Mirjana21, 55Köhler, Paul36Malara, Mirjana21, 55Köhninger, Fabian94Malsagova, Asja T.15Köbp, Matthias20Manava, Panagiota11<	J		Langthaler, Patrick	45
Jakab, Martin9, 37, 61Lapinskas, Simonas15Johansson, Tim87Lauth, Wanda42Legname, Giuseppe49KLeiler, Spela47Kaiser, Bernhard95Leis, Stefan102Kalisnik, Jurij Matija47Leisch, Michael43Kalss, Gudrun64Lell, Michael11Kamphausen, Anne35Lenhart, Armin27, 28Karcz, Konrad10, 15, 40Lessel, Davor93Karl, Tanja41Lether, Thomas84Karner, Josef89Ley, Pauline78Kartal, Orkan42Liebert, Adrian73Kerry, Ghassan73Lindbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kienslich, Tobias56Lorenzl, Stefan67Kiunl, Lorenz44Lubian, Francesca52Klenne, Friederike65Lutz, Nina65Klenne, Friederike65Lutz, Nina65Klenne, Friederike32Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koche, Thomas97Maieron, Andreas45Köbler, Paul36Malar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köninger, Fabian94Malsagova, Asja T.15	Jäger, Simon	65	Laner-Plamberger, Sandra	42
Johansson, Tim      87      Lauth, Wanda      42        Legname, Giuseppe      49        Kaiser, Bernhard      95      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kaisnik, Jurij Matija      47      Leisch, Michael      43        Kalss, Gudrun      64      Lell, Michael      11        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Karcz, Konrad      10, 15, 40      Lessel, Davor      93        Karl, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kerry, Ghassan      73      Lindlbauer, Nadja      42        Kien, Dominik      43      Lippert, Viktor      90        Kiener, Teresa      2      Liuni, Raffaella      37, 48, 49        Kieninger, Bärbel      62      Lobmann, Sebastian      50, 51        Kieslich, Tobias      56      Lorezl, Stefan      67        Kiwull, Lorenz      44      Lubian, France	Jakab, Martin	9, 37, 61	Lapinskas, Simonas	15
KLegname, Giuseppe49Kaiser, Bernhard95Leis, Stefan102Kaissr, Burnhard95Leis, Stefan102Kaissr, Gudrun64Lell, Michael43Kamphausen, Anne35Lenhart, Armin27, 28Karcz, Konrad10, 15, 40Lessel, Davor93Karl, Tanja41Lettner, Thomas84Karner, Josef89Ley, Pauline78Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiunul, Lorenz44Lubian, Francesca52Klenk, Jochen65Lutz, Nina65Klinglmayer, Friederike65Lutz, Nina65Kinglmayer, Fraiderike32Madorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15	Johansson, Tim	87	Lauth, Wanda	42
K      Leiler, Spela      47        Kaiser, Bernhard      95      Leis, Stefan      102        Kalisnik, Jurij Matija      47      Leisch, Michael      43        Kalss, Gudrun      64      Lell, Michael      11        Kamphausen, Anne      35      Lenhart, Armin      27, 28        Karcz, Konrad      10, 15, 40      Lessel, Davor      93        Karl, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Lieberh, Adrian      73        Kemeter, Melissa      30      Liesenberg, Tanja      28        Kerry, Ghassan      73      Lindlbauer, Nadja      42        Kieen, Dominik      43      Lippert, Viktor      90        Kiener, Teresa      2      Liuni, Raffaella      37, 48, 49        Kieninger, Bärbel      62      Lobmann, Sebastian      50, 51        Kieninger, Friederike      65      Lutz, Nina      65        Klenner, Friederike      65      Lutz, Nina      65        Klenner, Friederike      65 </td <td></td> <td></td> <td>Legname, Giuseppe</td> <td>49</td>			Legname, Giuseppe	49
Kaiser, Bernhard    95    Leis, Štefan    102      Kalisnik, Jurij Matija    47    Leisch, Michael    43      Kalss, Gudrun    64    Lell, Michael    11      Kamphausen, Anne    35    Lenhart, Armin    27, 28      Karcz, Konrad    10, 15, 40    Lessel, Davor    93      Karl, Tanja    41    Lettner, Thomas    84      Karner, Josef    89    Ley, Pauline    78      Kartal, Orkan    42    Liebert, Adrian    73      Kemeter, Melissa    30    Liesenberg, Tanja    28      Kerry, Ghassan    73    Lindibauer, Nadja    42      Kienen, Teresa    2    Liuni, Raffaella    37, 48, 49      Kieninger, Bärbel    62    Lobmann, Sebastian    50, 51      Kiesslich, Tobias    56    Lorenzl, Stefan    67      Kilnglmayer, Eva    70    M    78      Kinab, Katja    22    Maeding, Nicole    78, 83, 97, 102      Kneskeler, Guntje    30    Majeed, Yasir    53      Kocher, Thomas    97    Maeiron, Andreas    45      Kobler,	к		Leiler. Spela	47
Kalisnik, Jurij Matija    47    Leisch, Michael    43      Kalss, Gudrun    64    Lell, Michael    11      Kamphausen, Anne    35    Lenhart, Armin    27, 28      Karcz, Konrad    10, 15, 40    Lessel, Davor    93      Karl, Tanja    41    Lettner, Thomas    84      Karner, Josef    89    Ley, Pauline    78      Kartal, Orkan    42    Liebert, Adrian    73      Kemeter, Melissa    30    Liesenberg, Tanja    28      Kerry, Ghassan    73    Lindlbauer, Nadja    42      Kienn, Dominik    43    Lippert, Viktor    90      Kiener, Teresa    2    Liuni, Raffaella    37, 48, 49      Kieninger, Bärbel    62    Lobmann, Sebastian    50, 51      Kiesslich, Tobias    56    Lorenzl, Stefan    67      Kiunglmayer, Eva    70    M    53      Kochen    65    Lutz, Nina    65      Klinglmayer, Eva    70    M    53      Kochen    52    Mahorivska, Iryna    53      Koch, Oliver    30    Majeed	Kaiser, Bernhard	95	Leis, Stefan	102
Kalss, Gudrun    64    Lell, Michael    11      Kamphausen, Anne    35    Lenhart, Armin    27, 28      Karcz, Konrad    10, 15, 40    Lessel, Davor    93      Karl, Tanja    41    Lettner, Thomas    84      Karner, Josef    89    Ley, Pauline    78      Kartal, Orkan    42    Liebert, Adrian    73      Kemeter, Melissa    30    Liesenberg, Tanja    28      Kerry, Ghassan    73    Lindlbauer, Nadja    42      Kiem, Dominik    43    Lippert, Viktor    90      Kiener, Teresa    2    Liouni, Raffaella    37, 48, 49      Kieninger, Bärbel    62    Lobmann, Sebastian    50, 51      Kiesslich, Tobias    56    Lorenzl, Stefan    67      Kiwull, Lorenz    44    Lubian, Francesca    52      Klenk, Jochen    65    Lutz, Nina    65      KlingImayer, Eva    70    M    53      Koch, Oliver    30    Majeed, Yasir    54      Kocher, Thomas    97    Maieron, Andreas    45      Köbler, Paul    36	Kalisnik. Jurii Matiia	47	Leisch, Michael	43
Kamphausen, Anne      35      Lenhart, Armin      27, 28        Karcz, Konrad      10, 15, 40      Lessel, Davor      93        Karl, Tanja      41      Lettner, Thomas      84        Karner, Josef      89      Ley, Pauline      78        Kartal, Orkan      42      Liebert, Adrian      73        Kemeter, Melissa      30      Liesenberg, Tanja      28        Kerry, Ghassan      73      Lindlbauer, Nadja      42        Kiem, Dominik      43      Lippert, Viktor      90        Kiener, Teresa      2      Liuni, Raffaella      37, 48, 49        Kieninger, Bärbel      62      Lobmann, Sebastian      50, 51        Kiesslich, Tobias      56      Lorenzl, Stefan      67        Kiunglmayer, Eva      70      M      M        Kinab, Katja      22      Maeding, Nicole      78, 83, 97, 102        Knab, Katja      22      Maeding, Nicole      78, 83, 97, 102        Knab, Katja      22      Maeding, Nicole      78, 83, 97, 102        Knab, Katja      22      Maeding, Nicole      78, 83, 97, 102	Kalss, Gudrun	64	Lell, Michael	11
Karcz, Konrad10, 15, 40Lessel, Davor93Karl, Tanja41Lettner, Thomas84Karner, Josef89Ley, Pauline78Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kiensiger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kivull, Lorenz44Lubian, Francesca52Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kamphausen, Anne	35	Lenhart, Armin	27, 28
Karl, Tanja41Lettner, Thomas84Karner, Josef89Ley, Pauline78Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenk, Jochen65Lutz, Nina65Klenk, Jochen65Kitja70Knab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köpp, Matthias20Manava, Panagiota11	Karcz, Konrad	10, 15, <b>40</b>	Lessel, Davor	93
Karner, Josef89Ley, Pauline78Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenk, Jochen65Klenk, Jochen65Kinab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Karl, Tanja	41	Lettner, Thomas	84
Kartal, Orkan42Liebert, Adrian73Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Karner, Josef	89	Lev, Pauline	78
Kemeter, Melissa30Liesenberg, Tanja28Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klenk, Jochen65Kinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kartal, Orkan	42	Liebert, Adrian	73
Kerry, Ghassan73Lindlbauer, Nadja42Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kemeter, Melissa	30	Liesenberg, Tanja	28
Kiem, Dominik43Lippert, Viktor90Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kerry, Ghassan	73	Lindlbauer, Nadia	42
Kiener, Teresa2Liuni, Raffaella37, 48, 49Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Kochr, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kiem, Dominik	43	Lippert, Viktor	90
Kieninger, Bärbel62Lobmann, Sebastian50, 51Kiesslich, Tobias56Lorenzl, Stefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klenk, Jochen65Kinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kiener. Teresa	2	Liuni, Raffaella	37, <b>48, 49</b>
Kiesslich, Tobias56Lorenzl, Štefan67Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klenk, Jochen65Kinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kieninger, Bärbel	62	Lobmann, Sebastian	50, 51
Kiwull, Lorenz44Lubian, Francesca52Klenner, Friederike65Lutz, Nina65Klenk, Jochen65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kiesslich, Tobias	56	Lorenzl. Stefan	67
Klenner, Friederike65Lutz, Nina65Klenk, Jochen65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kiwull. Lorenz	44	Lubian. Francesca	52
Klenk, Jochen65Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Klenner. Friederike	65	Lutz. Nina	65
Klinglmayer, Eva70MKnab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Klenk, Jochen	65	,	
Knab, Katja22Maeding, Nicole78, 83, 97, 102Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Klinglmayer, Eva	70	Μ	
Kneiseler, Guntje32Mahorivska, Iryna53Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Knab. Katia	22	Maeding, Nicole	78, 83, 97, 102
Koch, Oliver30Majeed, Yasir54Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kneiseler, Guntie	32	Mahorivska, Iryna	53
Kocher, Thomas97Maieron, Andreas45Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Koch. Oliver	30	Maieed, Yasir	54
Köbler, Paul36Malnar, Mirjana21, 55Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Kocher, Thomas	97	Maieron, Andreas	45
Köninger, Fabian94Malsagova, Asja T.15Köpp, Matthias20Manava, Panagiota11	Köbler, Paul	36	Malnar, Miriana	21.55
Köpp, Matthias 20 Manava, Panagiota 11	Köninger, Fabian	94	Malsagova. Asia T.	15
	Köpp, Matthias	20	Manava, Panagiota	11

### XIV

Mani, Rakesh	55	Oesterreich, Alina	24
Mantegna, Gianluca	49	Ombres, Laura	42
March, Albert	52	Oostingh, Gertie J.	41
Marino, Angela	49	Ortner, Felix	15
Mariozzi, Sara	72	Österbrink, Jürgen	67
Markova, Gabriela	24.39	Otto, Ferdinand	71
Marr. Lisa	62	0,	
Marschhofer Moritz	101	Р	
Matea Cristian-Tudor	70	Paal Piret	67
Matthe Matthias	89	Paar Vera	14
Mavr. Christian	56	Pachmayr Johanna	2
Mayr, Onnetian Mayr, Daniela	56	Pahernik Sascha	35
Mega Andrea	45 <b>57</b>	Panebianco Luca	64
Megas Joannis Eivos	40, <b>01</b> 15	Panzer Simone	65
Meisner Keber Nicola	70	Parizer, Simone Datal Visbal	05
Melehardt Thomas	10	Palei, Visilai Douluchor, Pornhord	90
Metzger Illrike	40	Paulweber, Derinaru Dolzmon, Sonio	40
Minnish Bornd	24	Peizinan, Sonja Dieus, Deberte	<b>00</b> 75
Madak Julia	27	Picus, Roberto Diravai Niek	75
Magedi Everet Eleventing	13	Pilavej, Nick Divich Christian	4
	95	Pinch, Christian Ditear, Ctafar	2
Mock, Jonas	5	Pitzer, Stefan	67
Morabito, Rossana	49	Pizzadili, Gina	28
Moser, Tobias	/1	Plachel, Maximilian	68
Mostacci, Barbara	64	Pleyer, Lisa	43
Motloch, Lukas	14	Pollari, Francesco	69
Mrazek, Cornelia	42	Popotf, Alexander	<b>70</b> , 101
Mrowetz, Heike	99	Poupardin, Rodolphe	83, 97, 102
Mühling, Mischa	58	Prader, Sonia	8
Mülleder, Patrick	67	Priewasser, Beate	24, 31, 39
Müller, Markus	36, <b>59</b>	Plattner, B.	33
Müller, Silvana	27, 28	Prusik-Lut, Olaf	59
Musset, Boris	50, 51, 53, <b>60</b>	Pruszak, Jan	21, 55
		Püski, Tamas	15
N		Putz, Peter	91
Nasser, Houssein	37, <b>61</b>		
Neumann, Bernd	62	R	
Neureiter, Daniela	56	Radermacher, Peter	36,59
Neurohr, Claus	65	Radlberger, Richard Friedrich	71
Niemann, Stefan	62	Radzikowski, Konrad	88
Niklewski, Günter	79	Rainer, Lucas	20
Nguyen, Mai Quyen	63	Raninger, Anna	83, 97
Nguyen, Nhat	5	Rath, Anca	62
Nucera, Bruna	<b>64</b> , 72	Reddy, Sanjana	96
,	,	Reich, Fabian	81
0		Reichert, Bert	15
Obermaver. Astrid	27.97	Reinhardt, Alexander	12
O'Brien, James	68	Remigante, Alessia	49
Ofir. Racheli	97	Rendl. Gundula	2
, i donon	01		2

Riazy-Hossfeld, Noosha	70
Rieder, Sophie-Marie	99
Richter, Kneginja	3, 72
Rinaldi, Fabrizio	64, <b>72</b>
Rinnert, Alexander	87
Rinnerthaler, Gabriel	43
Rinnerthaler, Mark	102
Ritter, Leonard	73
Ritter, Markus	56, 77
Rochow, Niels	22
Roecken, Sofie	59
Röder, Falk	89
Rösch, Sebastian	9, 37, 48, 61
Rohde, Eva	42
Rohleder, Michael	59
Rosenberg-Granzner, Adrian	22
Rückl, Alena	102
Rüdig, Vivienne	12
Russe, Elisabeth	94, 102
Rupp, Bernhard	16
Ruznic, Elvis	89

#### S

Safdarian, Mahdi	74
Salman, Saif	96
Sandriesser, Sabrina	58
Santarelli, Marco	75
Santoro, Robin	14
Sauerbeck, Annette	4
Schauer-Berg, Johanna	76
Schäfer-Eckart, Kerstin	27, 28
Schider, Nicole	21
Schmid, Barbara	73
Schrader, Eva	53
Schubert, Mario	101
Seiberl, Michael	71
Sluet, Patrick	47
Schallmoser, Katharina	97
Schmid, Barbara	11
Schmidbauer, Josef	4
Schmidt, Manfred	100
Schmitz-Grosz, Krisztina	79
Schneider-Brachert, Wulf	62
Schäfer-Eckart, Kerstin	92
Schröter, Moritz	77
Schropp, Jonas	39
Schulze-Tanzil, Gundula	12, 13, 14, 27, 28
Schumann, Lena	15

Schulze-Tanzil, Gundula Schuster, Anja Schuster, Daniela Schwaiger, Karl Schweizer, Matthias Sedghi Aminabad, Negar Seitelberger, Rainald Semmler, Georg Sharma, Rohan Sicorschi Gutu, Cristina Singler, Katrin Sirch, Joachim Sklenár, Marek Sodmann, Johanna Sörgel, Fritz Sommer, Carsten Somogyi, Krisztina Speer, Robert Spinelli, Sara Spittler, Andreas Stana, Markus Stangassinger, Lea M. Stefan, Victoria E Stefani, Antri Stein, Barbara Stein, Stefan Steiner, Hans-Herbert Steinhuber, Anna Steinmann, Jörg Stitzel, Andrea Stöggl, Thomas Strubegger, Barbara Strunk, Dirk Studnicka, Michael	92 41 25, 70, 101 94, 100 53 <b>78</b> 13 88 96 <b>79</b> 32 69 <b>80</b> <b>80</b> <b>81</b> 35 79 22 32 69 <b>80</b> <b>81</b> 35 79 22 32 49 97 89 41 <b>82</b> 78, <b>83</b> , 102 36, 59 53 73 83 62 76 68 24 78, 83, 97, 102 95
T Tawk, Rabih Temml, Veronika Tempfer, Herbert Teuschl-Woller, Andreas Tevini, Julia Thomschewski, Aljoscha Thumfart, Lucas Tomson, Torbjörn Traweger, Andreas Temml, Veronika Trenkwalder, Katharina	96 25 <b>84</b> 92 82, 102 74 29, 30 64 18, 84, 92 101 <b>85</b>

Trinka, Eugen

20, 45, 64, 72, 74

#### XVI

Tsolova, Dayana Tutmann-Scheld, Michaela	15 15	Weinbender, Sofia Chanel Weisemann, Ferdinand	<b>90</b> , 99 85
	10	Werner Christian	12 14 28 92
u –		Wernly Bernhard	45 88
Ugorová Diana	80	Wernly, Bernhald Wernly, Sarah	-10, 00
Unger Michael Stefan	99	Wichlas Florian	68
Urbanczyk Sonhia	59 86	Widhalm Klaus	91
orbanozyk, oopnia	00, 00	Wiedmann-Kline Clea	92
V		Wildburger Sonia	77
Vanacore Nicola	52	Willim Jana	93
Van der Zee-Neuen Antie	77 87	Wiltzsch Sven	27 28
Varga Lukáš	80	Wimmer Florian	94
Vari, Balazs	97	Winkler, Martin	5
Vittadello, Fabio	57	Wipfler, Peter	71
Vockner. Sara	89	Wipplinger, Anna	56
Völker. Sebastian	87	Wirtz. Miria	95, 96
Völkerer. Andreas	88	Wolf. Martin	78, 83, <b>97</b> , 102
Volk, Hans-Dieter	97	Wosko, Paulina	67
Vollmar, Christian	20	Würflein, Dieter	81
Von Imhoff, Phillip	15	Wusatiuk, Carmen	98
W		Х	
Wagner, Andrea	18	Xiao, Fenglai	20
Waller, Christiane	36, 59		
Wandschneider, Britta	20	Z	
Wanninger, Anke	15	Zattoni, Marco	90, <b>99</b>
Weber, Daniela D.	82, 102	Zaussinger, Maximilian	100
Weber, Lucia	69	Zehentmayr, Franz	89
Wechselberger, Gottfried	94	Zell, Lukas	101
Weghuber, Daniel	45	Zembacher, Stefan	95
Wehrmann, Julius	12	Zimmermann, Georg	42, 64
Wei, Yujia	96	Zink, Fabian	59
		Zwicklhuber, Janine	102

# **#SGT**



Paracelsus Medizinische Privatuniversität Forschungsmanagement | Salzburg Strubergasse 21, 5020 Salzburg, Austria +43 (0)662 2420-80281 www.pmu.ac.at

ISBN: 978-3-9505441-2-1